
Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.







En83:B

.

9







U.S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

OF

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED. BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

BY

SAMUEL HENSHAW.

Parts I, II, and III.

THE MORE IMPORTANT WRITINGS

BENJAMIN DANN WALSH

AND
CHARLES VALENTINE RILEY.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1890.



3523 32

U.S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

of

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

BY

SAMUEL HENSHAW.

PART I.

THE MORE IMPORTANT WRITINGS

 \mathbf{OF}

BENJAMIN DANN WALSH.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1889.

FORTY-SEVENTH CONGRESS, FIRST SESSION.

CONGRESS OF THE UNITED STATES,
IN THE HOUSE OF REPRESENTATIVES,

June 24, 1882.

Resolved, By the House of Representatives (the Senate concurring) that there be printed, for the use of the Department of Agriculture, one thousand copies of a special report, entitled "Bibliography of Economic Entomology."

Attest:

EWD. McPherson, Clerk.

Passed the Senate July 6, 1882.

F. E. Shober,

Acting Secretary.

A true copy.

EWD. McPherson, Clerk.

2

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
DIVISION OF ENTOMOLOGY,
Washington, D. C., June 12, 1889.

SIR: In the absence of Professor Riley, who is now at the Paris Exposition in official capacity, I beg to submit for publication Parts I, II, and III of the Bibliography of American Economic Entomology, which was ordered by Congress in July, 1882. Owing to various unforeseen delays this work has not yet been completed. Its preparation was originally assigned by Professor Riley to Mr. B. Pickman Mann, formerly of this Division, and upon the severance of his connection with the Department, in the fall of 1885, it was turned over in an incomplete and fragmentary condition to Mr. Samuel Henshaw, of Boston, who has added greatly to it and practically rewritten the portions already prepared.

The extent of the work renders it advisable to publish it in several parts, and as there is unquestionably a more urgent need on the part of working economic entomologists for a bibliography of the writings of B. D. Walsh and C. V. Riley than of others, on account of their great number and value, these portions have been finished first. Their immediate publication will probably antecede the publication of the final part only by a few months.

Very respectfully, yours,

L. O. HOWARD,

Acting Entomologist.

Hon. J. M. Rusk, Secretary of Agriculture.



TABLE OF CONTENTS TO PARTS I, II, AND III.

	Page.
Introduction	., 7
The writings of B. D. Walsh (1860–1873)	9
The joint writings of B. D. Walsh and C. V. Riley (1868-1869)	51
The writings of C. V. Riley (1863-1888)	97
Systematic index of the new names:	
By B. D. Walsh and by Walsh and Riley	373
By C. V. Riley	377
Index	



INTRODUCTION TO PARTS I, II, AND III.

The object of the present lists is to record the principal entomological writings of B. D. Walsh and C. V. Riley. Few will be likely to criticize the association of the names of Walsh and Riley in a list of this kind, so largely devoted to economic entomology, or to question the desirability of collating their writings in advance of others.

As recognized authorities upon entomology, especially from an economic standpoint, their writings have been extensively quoted sometimes with, but frequently without, credit in the agricultural and horticultural journals of the day. These secondary references, though of a certain value when the original is inaccessible, have, as a rule, been omitted. In citing such as are given I have used my personal judgment. To have given all would have increased the size of the list considerably with but a slight increase in its usefulness.

Occasionally, a publisher to meet some popular demand puts the writings of an author into book form without the knowledge of the latter. Such an occurrence has happened more than once with Dr. Riley's writings. To mention but one: In 1877 Messrs. George Rutledge & Sons, of London, published "The Colorado beetle, with suggestions for its repression and methods of destruction," with Dr. Riley's name upon the title-page as author. This is a partial reprint of Potato Pests entirely unsanctioned and published without knowledge of the author. (See London Times, October 17, 1887.) These publishers' reprints are omitted from this list. The "S.-b." after the reference of many of the articles from the agricultural and other journals refers to the series of scrap-books used in divisional work.

A biographical sketch of Dr. Walsh is given in the second volume of the *American Entomologist*, Vol. II, No. 3, pp. 65-68, January, 1870.

The chief facts in Dr. Riley's life may be found in the "Commonwealth of Missouri" (1875) and the National Farmer, September 20, 1883.

In the preparation of the list I have freely used all data accumulated by the Division, notably the work of Mr. B. P. Mann and a private list of Dr. Riley's.

Many persons, both within and without the Department, have kindly answered my inquiries.—S. H.



BIBLIOGRAPHY OF THE MORE IMPORTANT CONTRIBUTIONS TO AMERICAN ECONOMIC ENTOMOLOGY.

THE WRITINGS OF B. D. WALSH (1860-1873).

- 1. Walsh, B. D. Bark-lice. <Western Rural. S.-b. No. 3, p. 11. Habits of Lecanium compared with Aspidiotus [=Mytilaspis].
- 2. Walsh, B. D. Entomological notes. < Prairie Farmer, 17 May, 1860, [v. 21], n. s., v. 5, pp. 308-309, figs. S.-b. No. 1, pp. 42-43.
 - Notice of Coccus harrisii n. sp. [=Chionaspis furfurus]; figure of its scales; description and figure of Capsus oblineatus [=Lygus pratensis] infesting apple, quince, and pear trees; figure of larva case and imago and description of Phycita nebulo [=Acrobasis indiginella]; figures of imago and of injuries of Amphicerus bicaudatus; habits of the same and of Scolytus [=Xyleborus] pyri, Saperda bivittata [=candida] and Chrysobothris femorata; means against the last.
- 3. Walsh, B. D. "Cutworms." < Prairie Farmer, 9 August, 1860, [v. 22], n. s., v. 6, pp. 82–83, fig. S.-b. No. 1, pp. 84–85.
 - Account of the larvæ of Lachnosterna quercina [=fusca] of Elateridæ of Agrotidæ and of Pangus [=Harpalus] caliginosus; figures of larvæ and imagos of the same or allied species; also of Heteronychus [=Ligyrus] relictus, Lopha [=Bcmbidium] 4-maculatum and Agonoderus pallipes.
- 4. Walsh, B. D. [*Ips quadrisignatus*, Say.] < Prairie Farmer, 1860, [v. 22], n. s., v. 6, p. —.
 - Ips 4-signatus [=fasciatus] attacks growing ears of sweet corn.
- 5. Walsh, B. D.] "Insect Life." < Proc. Agric. Conv.; Ann. meeting Ill. Nat. Hist. Soc.; Comm. Exerc. State Normal Univ. [Bloomington, Ill.], 1860, pp. 11-12.
 - Report, compiled from Chicago papers of address delivered before the Illinois Natural History Society; comparison between injurious insects and an invading army; necessity of appropriations for carrying on entomological work; economic entomology in Europe; importance of beneficial insects.

- 6. Walsh, B. D. Insects injurious to vegetation in Illinois. < Trans. Ill. State Agric. Soc., September, 1861, v. 4, pp. 335-378, figs. 1-11. Separate: 1861, 43 pp., 1 pl. Reprint in part: < Trans. Ill. Nat. Hist. Soc., v. 1. Prairie Farmer, 6 December, 1861, [v. -], n. s., v. -, pp. 370-371.
 - Ravages of injurious insects; enemies of the same; notes on Cecidomyia destructor, Saperda bivittata [=candida], Chrysobothris femorata, Conotrachelus nemphar and Blissus leucopterus; natural history, ravages and means against Leucania unipuncta; figures its larva, pupa and imago; descriptions and figures primary and secondary parasites of the army-worm; figures Hippodamia (=Megilla) maculata, Coccinella munda [=sanguinea] and a larval coccinellid.
- 7. Walsh, B. D. From Benj. D. Walsh. < Illinois Farmer, October, 1861.
 - Denies statement attributed by C. Thomas (Illinois Farmer, September, 1861), that he (Walsh) believed in the hibernation of the pupa of *Leucania* unipuncta.
- 8. Walsh, B. D. The army-worm question. < Prairie Farmer, 5 December, 1861, [v. 24], n. s., v. 8, pp. 370-371. S.-b., No. 4, p. 22. Replies to arguments of C. Thomas (Prairie Farmer, 1861, v. 8, pp. 306-307),

on the hibernation of Leucania unipuncta.

- 9. Walsh, B. D. The army-worm question. Mr. Walsh's reply. < Field Notes, 14 December, 1861.
 - Controversial answer to the assertion of J. H. Klippart (Field Notes, 30 November, 1861), that the larvæ of Leucania unipuncta are viviparous.
- 10. Walsh, B. D. The army-worm and its enemies. <Prairie Farmer, 1861, [v. 24], n. s., v. 8, p. 4. Reprint: <Trans. Ill: Nat. Hist. Soc., v. 1. Prairie Farmer, 6 December, 1861, [v. 24], n. s., v. 8, pp. 370-371.
 - Popular account of some of the parasites of Leucania unipuncta.
- 11. Walsh, B. D. The army-worm. <Prairie Farmer, 1861, [v. 24], n. s., v. 8, pp. 257-258. Reprint: <Trans. Ill. State Agric. Soc., 1861, v. 4, pp. 373-375.
 - Hibernation of the egg; spring burning as a means against Lucania unipuncta; description of Hockeria [=Haltichella] perpulchra n. sp.; notes on other parasites.
- 12. Walsh, B. D. The army-worm and its insect foes. < Prairie Farmer, 1861, [v. 24], n. s., v. 8, pp. 322-323; 337-339; 354-355.
 - Habits and description of larva and imago of Leucania unipuncta; hibernation of the egg; figures of larva, pupa and imago; also of Exorista [=Nemoræa] leucaniæ, Pezomachus minimus, Microgaster [=Apanteles] militaris, Hockeria [=Haltichella] perpulchra and Glyphe viridascens.
- 13. Walsh, B.D. Bug preying on honey-bee. <Prairie Farmer, 1861. S.-b., No. 4, p. 7.

Unknown heteropteron, said to prey on Apis mellifica.

- 14. Walsh, B. D. The pea and its insect foe. <Prairie Farmer, 1861. S.-b., No. 4, p. 12.
 - Unknown leaf-miner and leaf-eater on pea; directions for rearing insects.
- 15. Walsh, B. D. The grain weevil. <Journ. Ill. State Agric. Soc., January, 1862, fig. Reprint: <Trans. Ill. State Agric. Soc. v. 5, pp. 484-485.
 - Sitophilus [= Calandra] remote punctata infests wheat.
- 16. Walsh, B. D. [Ithycerus noveboracensis.] <St. Louis Valley Farmer, March, 1862, v. 14, pp. 82–85, fig.

 Ithycerus noveboracensis attacks twigs of fruit trees in nurseries.
- 17. Walsh, B. D. [Army-worm.] <St. Louis Valley Farmer, 1862, v. 14, p. 161.
 - Correction of mistake of E. S. Washington (St. Louis Valley Farmer, 1862, v. 14, p. 161), who mistook parasitic larvæ for young *Leucania unipuncta*; economy of parasites of army-worm.
- 18. Walsh, B. D. [Brachytarsus variegatus.] <Journ. Ill. State Agric. Soc., March, 1862, pp. 8-12, fig.

 Brachytarsus variegatus parasitic (?) on a large orange-colored cecidomyid (?)

larva in stems of wheat.

- 19. Walsh, B. D. A new insect in wheat, Bruchus. <Journ. Ill. State Agric. Soc., April, 1862. Reprint: <Trans. Ill. State Agric. Soc., v. 5, pp. 485-490.
- 20. Walsh, B. D. [Two apple-tree borers.] < Journ. Ill. State Agric. Soc., June, 1862, pp. 21-23.

 Treats of Chrysobothris femorata and Saperda bivittata [= candida].
- 21. Walsh, B. D. [Colorado potato beetle.] <St. Louis Valley Farmer, July, 1862, pp. 209-210. Reprint (?): <Prairie Farmer, 6 June, 1863, v. —, p. 356, fig.
 - Doryphora 10-lineata infests egg plants, potato and tomato vines, etc., in Kansas and Iowa.
- 22. Walsh, B. D. Fire-blight. Two new foes of the apple and pear. <Prairie Farmer, 6 September, 1862, [v. 26], n. s., v. 10, pp. 147-149, fig. Separate: pp. 4.
 - Eire-blight defined; probably caused by attacks of Chloroneura malefica [=Empoasea viridescens] and C. maligna [=E. obtusa]; characters, habits, and ravages of the same; synoptic separation of several genera of Typhlocybini; two (2) new genera and thirteen (13) new species are described. For a list of the same see the Systematic Index. The figures show C. malefica and C. maligna and the venation of allied forms.
 - (Note.—The descriptive portion with figures is reprinted in Proc. Bost. Soc. Nat. Hist., February, 1864, v. 9, pp. 314-318.)
- 23. Walsh, B. D. Plant lice, the corn-root louse, a new enemy to the corn. <Journ. Ill. State Agric. Soc., September, 1862, pp. 8-13, fig. Reprint: <Trans. Ill. State Agric. Soc., v. 5, pp. 491-497, fig.
 - Aphis maidis (?) infests the roots of young Indian corn as well as the stems of the roasting ears.

- 24. Walsh, B. D. List of the Pseudoneuroptera of Illinois contained in the cabinet of the writer, with descriptions of over forty new species, and notes on their structural affinities. <Proc. Acad Nat. Sci. Phil., September, 1862, pp. 361-402.
 - One hundred and nine (109) species are enumerated; two (2) new genera and forty (40) new species are described. For a list of the same see the Systematic Index.
- 25. Walsh, B. D. [Erythroneura tricincta Fitch.] < St. Louis Valley Farmer, October, 1862, pp. 305-306, fig.

Erythroneura [=Typhlocyba] tricincta on grape-vines.

- 26. Walsh, B. D. Grasshoppers and locusts. < Journ. Ill. State Agric. Soc., November, 1862, pp. 1–3. Reprint: < Trans. Ill. State Agric. Soc., v. 5, pp. 497–499.
 - Injuries of locusts in the Mississippi Valley; differences between locusts and Cicadas.
- 27. Walsh, B. D. On the genera of Aphida found in the United States. < Proc. Ent. Soc. Phil., December, 1862, v. 1, pp. 294–311, figs. 1–8.
 - Synoptic separation of the genera; list of seventy (70) species found in the United States, with food habitat; Aphis [=Nectarophora] rudbeckiw Fitch, A. vitis? Scop., A. maidis? Fitch, $Lachnus\ caryw$ Harris, $Thelaxes\ [=Colopha]$ ulmicola Fitch are described; one (1) new genus and eleven (11) new species are described; see the $Systematic\ Index$ for a list of the same; the figures show venation and other details. Two (2) new ants, $Formica\ aphidicola\ and\ F.\ [=Lasius]\ latipes$, are described.
- 28. Walsh, B. D. [Observations on *Papilio glaucus* and *P. turnus*.] < Proc. Ent. Soc. Phil., February, 1863, v. 1, pp. 349–352.
 - Reasons for considering turnus and glaucus identical; the latter a dimorphic Q of the former; distribution of the two forms; citation of similar case among Pieridw and of analogous one among Dytiscidw; description of larva from which a black Q was raised.
- 29. Walsh, B. D. [Katydid eggs.] < Prairie Farmer, 28 February, 1863, [v. 27], n. s., v. 11, p. 132, fig.

Oviposition of katydid eggs on apple-twigs.

- 30. Walsh, B. D. Fire-blight. < Prairie Farmer, 4 April, 1863, [v. 27], n. s., v. 11, p. 212, fig. S.-b., No. 1, p. 10.
 - Description and figures of eggs of Chloroneura malefica [=Empoasca viride-scens] and the slits in which they are deposited in apple-twigs; cause of fire-blight; suggestion of remedies.
- 31. Walsh, B. D. Insects injurious to fruit trees. < Prairie Farmer, 2 May, 1863, [v. 27], n. s., v. 11, p. 276, fig. S.-b., No. 1, p. 9.
 - Fruit and foliage of pear-trees destroyed by Capsus oblineatus [=Lygus pratensis]; description and figure of imago; habits, food-plants, and remedies.
- 32. Walsh, B. D. The ten-striped spearman. < Prairie Farmer, 6 June, 1863, [v. 27], n. s., v. 11, p. 356, fig. S.-b., No. 1, p. 7.
 - History, ravages, and means against *Doryphora* 10-lineata; figure of imago; mentions other injurious *Phytophaga*.

- 33. Walsh, B. D. The plum-gouger; a new foe of the plum. < Prairie Farmer, 13 June, 1863, [v. 27], n. s., v. 11, pp. 372–373, figs. 1–3. S.-b., No. 1, p. 6.
 - Habits, description, and figures of Anthonomus? prunicida n. sp. [= Coccotorus scutellaris], injurious to plums; comparison with Conotrachelus nenuphar; figure of the latter.
- 34. Walsh, B. D. The plum-gouger. < Prairie Farmer, 11 July, 1863, [v. 28], n. s., v. 12, p. 21, 2 figs. S.-b., No. 1, p. 10.
 - Distribution and ravages of Anthonomus prunicida [= Coccotorus scutellaris]; list of insects found on plum-trees; description of Conotrachelus puncticollis n. sp. [= C. geminatus]; usefulness of Reduvius raptatorius [= Sinca diadema]; figures A. prunicida and R. raptatorius.
- 35. Walsh, B. D. A new fruit foe. < Prairie Farmer, 18 July, 1863, [v. 28], n. s., v. 12, p. 37, fig. S.-b., No. 1., p. 11.
 - Description and figure of image of *Epicærus imbricatus*; injuries to fruit trees and gooseberry-bushes; supposed habits of this and allied species; hibernation of the larva of *Conotrachelus*; description of *C. cratægi* n. sp.
- 36. Walsh, B. D. Leaf-hopper. < Prairie Farmer, 25 July, 1863, [v. 28], n. s., v. 12, p. 53.
 - Proconia [= Oncometopia] undata injurious to grape-vines in southern Illinois; its eggs laid in twigs.
- 37. Walsh, B. D. Locust borers. < Prairie Farmer, 15 August, 1863, [v. 28], n. s., v. 12, p. 101. S.-b., No. 1, p. 12.
 - Description and habits of Clytus [= Cyllene] robiniæ; Clytus of Carya does not attack Robinia; Cossus robiniæ mentioned; larvæ and imagos of Nitidulidæ found under decaying bark.
- 38. Walsh, B. D. Insect friends and insect foes. The twice-stabbed lady-bird. < Prairie Farmer, 1863, [v. 28], n. s., v. 12; 22 August, p. 117, figs. 1, 2; 29 August, p. 133, figs. 3, 4. S.-b., No. 1, pp. 11–12.
 - Description and figure of imago of Chilocorus bivulnerus; figure of larva; destruction of injurious insects by predaceous insects; brief description of and means against Lytta cinerea [= Macrobasis unicolor], found on potato, English bean, and apple, and L. [= Epicauta] vittata, on potato; figures Lytta atrata [= Epicauta pennsylvanica], found on aster and potato, and larva-cases of Solenobia sp. from under bark of apple-trees; note on allies of Solenobia.
- 39. Walsh, B. D. Observations on certain N. A. Neuroptera, by H. Hagen, M. D., of Kænigsberg, Prussia; translated from the original French MS., and published by permission of the author, with notes and descriptions of about twenty new N. A. species of Pseudoneuroptera. < Proc. Ent. Soc. Phil., October, 1863, v. 2, pp. 167-272, fig.
 - Dr. Hagen's comments relate to the species described in No. 24; the author's notes supplement these and describe three (3) new genera and twenty-three (23) new species; for a list of these, see the Systematic Index; descriptions of previously described species, with notes on their habits, distribution, etc., are also given; Batis interlineata [=Siphlurus femoratus] is proposed

- 39. Walsh, B. D.—Continued.
 - (p. 190) in case B. [= S.] femorata Walsh is distinct from B. [= S.] femorata Say; in a note (p. 255) Ophiogomphus mainensis n. sp. is described from Paekard's manuscript; the venation of the Odonata and the terminal abdominal characters of Sialis infurnata are figured.
- 40. Walsh, B. D. On certain remarkable or exceptional larvæ, coleopterous, lepidopterous, and dipterous, with descriptions of several new genera and species, and of several species injurious to vegetation, which have been already published in agricultural journals. < Proc. Bost. Soc. Nat. Hist., February, 1864, v. 9, pp. 286-318.
 - Habits of the larva of Cotalpa lanigera, Pelidnota punctata, Cratonychus [= Melanotus] incertus, and Xyloryctes satyrus; comparative characters of the larva and imago of Halisidota antiphola n. sp. [= tessellaris]; description of larva, pupa, and imago of Sphingicampa (n. g.) distigma n. sp. [= bicolor]; characters and affinities of Dryocampa [= Sphingicampa] bicolor?; description of imagos of Limacodes scapha and L.? [= Phobetron] hyalinus n. sp., and of larva of the last and L.? [= Phobetron?] tetradactylus n. sp.; description of the larva, pupa, and imago of Hipparchiscus n. g. venustus n. sp. [= Aplodes mimosaria]; habits and description of the larva of an undetermined Tabanus, description of the pupa; description of the larva, pupa, and imago of Midas fulvipes n. sp., habits of its larva. On pp. 309-318 descriptions of several new species, with brief notes of food-habits, etc., are reprinted from various agricultural journals.
- 41. Walsh, B. D. On dimorphism in the hymenopterous genus *Cynips*; with an appendix containing hints for a new classification of *Cynipidw* and a list of *Cynipidw*, including descriptions of several new species, inhabiting the oak-galls of Illinois. < Proc. Ent. Soc. Phil., March, 1864, v. 2, pp. 443-500, fig.
 - Detailed observations proving the frequent occurrence of dimorphism in the Cynipidæ; Cynips [= Amphibolips] aciculata, a dimorphie form of C. q. spongifica; concludes from analogy "that aciculata Q generates galls which produce by parthenogenesis \mathcal{F} spongifica exclusively, and that \mathcal{Q} \mathcal{Q} spongifica coupling in June with these & & oviposit in the same month in the young buds of the oak, the eggs lying dormant till the following spring, when some of the eggs produce Q spongifica in June, and some Q aciculata in the autumn or early in the following spring, which last in their turn, as before mentioned, generate & spongifica to appear in the following June;" interesting details concerning the history of the group are given; relations between the true gall-flies (Psenides) and parasitic Cynipide (Inquiline); classification and characters of the same; anatomical structure and homologies of the family; the list includes fifteen (15) species inhabiting the various oaks, with descriptions of their galls and of several new species; seven (7) species of Inquilinæ are described; for the new species, see the Systematic Index; the figures illustrate the anatomy of the abdomen and ovipositor.
- 42. Walsh, B. D. The four-humped Curculio. A new foe of the apple. <Valley Farmer. Reprint: <Prairie Farmer, 27 August, 1864, [v. 30], n. s., v. 14, p. 131, 2 figs. S.-b., No. 1, p. 28.

Description and figures of Anthonomus quadrigibbus, a foe of the apple; comparison with Conotrachelus nenuphar and Anthonomus prunicida [= Coccotorus scutellaris].

43. Walsh, B. D. On the pupa of the ephemerinous genus Batisca Walsh. <Proc. Ent. Soc. Phil., August, 1864, v. 3, pp. 200-206, fig.

Description and figures of the pupa of Batisca obesa Say.

44. Walsh, B. D. On certain entomological speculations of the New England school of naturalists. < Proc. Ent. Soc. Phil., August-September, 1864, v. 3, pp. 207-249.

Discussion of statements, chiefly by Prof. L. Agassiz, upon questions in general entomology. The memoir is divided into sections; in the first, the distribution of insects in North America is discussed in opposition to the views of Agassiz; in the second section the Darwinian theory of the origin of species is discussed and some remarks on the variation of several species are added; in the third and fourth sections the statements of Prof. Agassiz regarding the worm-like larva stage of insects and the resemblances between the pupe of insects and the Crustacea are discussed and criticised; in the fifth section the author argues that the relative rank of the different orders of insects must be determined from a consideration of all the characters of each order and not from the significance of any one character; the sixth section is a review of Dana's classification of insects on the principle of cephalization; and in the seventh section several erroneous statements and generalizations in the same author's paper on classification are pointed out.

45. Walsh, B. D. On phytophagic varieties and phytophagic species. < Proc. Ent. Soc. Phil., November, 1864, v. 3, pp. 403-430.

Many species feed exclusively upon a single food-plant, while other species feed upon many kinds; correlated with this are certain larval or imaginal differences; when these different forms interbreed they are considered the same species and are termed phytophagic varieties; these when separated with a single kind of food-plant form phytophagic races, and these again by a continuation of their isolation form phytophagic species. Phytophagous forms are most abundant where the imago is wingless. Diapheromera velii, Tingis [=Gargaphia] tilia, and T. [=G.] amorpha are described as new; the following are mentioned in the remarks upon species and varieties: Haltica [=Disonycha] alternata, Chrysomela scalaris, Clytus [=Cyllene] robinia, C. [=C.] pictus, Dryocampa [=Sphingicampa] bicolor, Sphingicampa distigma [=bicolor], Bombyx [= Sericaria] mori, Halisidota tessellaris, H. carya, H. antiphola [=tessellaris], Cynips [=Amphibolips] q. spongifica, C. [=Andricus] q. punctata, C. [=A.] q. podagra, and C. [=Amphibolips] q. inanis.

46. Walsh, B. D. On the insects, coleopterous, hymenopterous, and dipterous, inhabiting the galls of certain species of willow. Part 1st—Diptera. <Proc. Ent. Soc. Phil., December, 1864, v. 3, pp. 543-644.

Structural characters, habits, metamorphoses, etc., of *Cecidomyidæ*; remarks on the unity of habits in genera; synopsis of cecidomyidous galls of *Salix*; descriptions of new species, their galls and inquilines; for a list of the new species, see the *Systematic Index*. See No. 197 for Part 2d.

47. Walsh, B. D. The borer. < Colman's Rural World. Reprint: < Prairie Farmer, 6 May, 1865, v. 15, p. 355.

Notice of soap as a means against borers.

48. Walsh, B.D. The new potato-bug and its natural history. < Pract. Ent., 30 October, 1865, v. 1, pp. 1-4.

Record of the spread eastwards of *Doryphora* 10-lineata; description of the egg; food-plants, enemies, and means against the same; comparative characters of *D. juncta* and *D.* 10-lineata; species of *Meloidæ* injurious to the potato.

49. [Walsh, B. D.] The joint-worm. < Pract. Ent., 27 November, 1865, v. 1, pp. 11, 12.

Reasons for disbelieving that the joint-worm is the larva of one of the *Chal-cidide*.

- 50. Walsh, B. D. On phytophagic varieties and phytophagic species, with remarks on the unity of coloration in insects. <Proc. Ent. Soc. Phil., November-December, 1865, v. 5, pp. 194-216, fig. See: <Amer. Journ. Sci., September, 1865, ser. 2, v. 40, pp. 282-284. <Ann. and Mag. Nat. Hist., November, 1865, ser. 3, v. 16, pp. 383, 384.
 - Argues for the origin of races and species by phytophagic isolation; refers to Datana ministra, D. contracta, Halisidota tessellaris, H. antiphola [= H. tessellaris], H. harrisii [= H. tessellaris], Arhopalus [= Cyllene] pictus, A. [= C.] robiniæ, A. infanstus [= C. decorus], Callidium antennatum, C. janthinum, Conotrachelus nenuphar, Doryphora 10-lineata, and D. juncta; unity of coloration explained by assumption of a genetic connection between various species; figure showing bullæ on wing of 1chneumon.
- 51. W[ALSH], B. D. [Cicada years.] < Pract. Ent., 25 December, 1865, v. 1, pp. 18–19.

Answer to inquiry of M. S. Hill; Cicada districts of the United States, as given by Fitch in N. Y. Rept. I, p. 39; habits of C. [= Tibicen] septendecim.

52. Walsh, B. D. Insects injurious to vegetation in Illinois. <Trans. Ill. State Agric. Soc., 1865, v. 5, pp. 469-483, figs.

Notes on injurious insects in Illinois in 1861; life-history and ravages of Leucania unipuncta.

- 53. Walsh, B. D. The "Thrips" of the vine-growers. What is it? <Pract. Ent., 25 December, 1865, v. 1, pp. 20-21.
 - The *Thrips* of fruit-growers probably not a true *Thrips*; food-habits of *Thrips*.
- 54. Walsh, B. D. The grub-worm. < Colman's Rural World, December, 1865. S.-b., No. 1, p. 51.
 - Habits, ravages, and means against Lachnosterna quercina [= fusca]; its larva compared with that of Ligyrus relictus; vernacular names of the Lachnosterna and of Allorhina nitida.
- 55. Walsh, B. D. Borers. < Pract. Ent., 29 January, 1866, v. 1, pp. 25-31, figs. 1-8.
 - Remarks on the name "borer;" figures, natural history, and means against Saperda bivittata [= candida], Chrysobothris femorata, Bostrichus [= Amphicerus] bicaudatus, Egeria [= Sannina] exitiosa, A. tipuliformis, Clytus [= Cyllene] robinia, C. [= C.] pictus, and Cerasphorus [= Chion] cinctus; need of State aid against injurious insects.

- 56. [Walsh, B. D.] [Attagenus? injuring feathers.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of R. Parnell; ravages of and means against Dermestidæ.
- 57. [Walsh, B. D.] [Codling moth.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of Isaac Hicks; natural history and means against Carpocapsa pomonella.
- 58. [Walsh, B. D.] [Aphididæ.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of J. H. Foster, jr.; comparative characters of *Eriosoma* [= Schizoneura] lanigera and Pemphigus pyri [= Schizoneura lanigera].
- 59. [Walsh, B. D.] [Mites mistaken for plant-lice.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of W. M. Smith; distinctive characters of Insecta and Arachnida; description and habitat of the eggs of Aphis mali.
- 60. [Walsh, B. D.] [Saddle-back caterpillar.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of S. P. M.; description of the larva and image of *Empretia stimulea*; urticating properties of the larva.
- 61. W[ALSH], B. D. [Red-humped prominent.] < Pract. Ent., 26 February, 1866, v. 1, p. 35.
 - Answer to inquiry of E. D. Wright; life-habits of Notodonta [= Œdemasia] concinna.
- 62. Walsh, B. D. The joint-worm. < Pract. Ent., 26 February, 1866, v. 1, pp. 37-38.
 - Habitats of the larva of Cecidomyia destructor and of Eurytoma sp.; criticism of A. Fitch's report upon the latter.
- 63. W[ALSH, B. D.]. Entomology all a humbug. <Pract. Ent., 26 February, 1866, v. 1, p. 39.
 - Fewness of entomologists and the multiplicity of their duties the reason why so little has been effected against noxious insects.
- 64. W[ALSH], B. D. [Cocoons of Ichneumon-flies.] < Pract. Ent., 26 March, 1866, v. 1, p. 46.
 - Answer to inquiry of M. S. Hill; life-habits of the Microgasters.
- 65. W[ALSH], B. D. [Mass of eggs.] < Pract. Ent., 26 March, 1866, v. 1, p. 46.
 - Answer to inquiry of T. R. Payne; characters and life-habits of Orgyia leucostigma.
- 66. [Walsh, B.D.] [Apple-tree caterpillar; parasites.] < Pract. Ent., 26 March, 1866, v. 1, p. 46.
 - Answer to inquiries of G. E. Brackett; food-plants of Clisiocampa americana; characters and habits of Tachinida; habits of secondary parasites.
- 67. W[ALSH], B. D. Apple-tree borers. < Pract. Ent., 26 March, 1866, v. 1, p. 47.
 - Reprint, with remarks, of "Apple-tree borers" (Journ. of Commerce); extract from A. Fitch, of larval habits of Saperda bivittata [= candida].

68. Walsh, B. D. The white grub. <Pract. Ent., 30 April, 1866, v. 1, pp. 60-62.

Descriptions and habits of the larval states of Lachnosterna quercina [=fusca] and Ligyrus relictus; ravages and food-habits of the former; the hog as a white-grub destroyer.

- 69. Walsh, B. D. [Apple-tree scales.] < Pract. Ent., 30 April, 1866, v. 1, p. 64.
 - Answer to inquiry of L. S. Pennington; descriptions and ravages of *Mytilaspis* pomorum and *Chionaspis furfurus*; Coccinellidæ as a means against the former; petroleum as a remedy against bark-lice.
- 70. Walsh, B. D. [Clothes-moths.] < Pract. Ent., 30 April 1866, v. 1, p. 64.

Answer to inquiry of S. Canby; habits and means against Tineidw.

71. Walsh, B. D. [Poplar borer.] < Pract. Ent., 30 April, 1866, v. 1, p. 64.

Answer to inquiry of P. C. Truman; larva and imago of Saperda calcarata infesting cottonwood.

72. WALSH, B. D. [Worms in flour and rye.] < Pract. Ent., 30 April, 1866, v. 1, p. 64.

Answer to inquiry of L. D. Hunt; descriptions and habits of *Tenebrio molitor* and *Dermestes lardarius*.

73. Walsh, B. D. Popular remedies for noxious insects. < Pract. Ent., 28 May, 1866, v. 1, pp. 71-74.

Criticism of various worthless remedies against injurious insects.

74. Walsh, B. D. [Serica iricolor Say.] < Pract. Ent., 28 May, 1866, v. 1, p. 77.

Answer to inquiry of B. F. Seibert; description, ravages, and habitat of Serica iricolor.

- 75. Walsh, B. D. [Measuring-worms.] < Pract. Ent., 28 May, 1866, v. 1, p. 77.
 - Answer to inquiry of T. Siveter; habits of Ennomos magnaria [=alniaria]; food-plants, habits, and means against E. [=Eudalimia] subsignaria and Anisopteryx [=Paleacrita] vernata.
- 76. Walsh, B. D. [Lygus pratensis.] < Pract. Ent., 25 May, 1866, v. 1, pp. 77-78.

Answer to inquiry of H. B. Howarth; habits and odors of Lygus pratensis; changes insects pass through.

77. WALSH, B. D. [Apple-tree insects.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.

Answer to inquiry of C. Cooke; an unnamed tineid moth raised from cocoons off twigs of apple-tree; *Hemiteles* sp. and *Microgaster* sp. bred from cocoons attached to eggs of *Clisiocampa americana*.

78. Walsh, B. D. [Destructive current-worm.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.

Answer to inquiry of W. M. Smith; cocoon and sexual characters of Nematus ribis [= ribesii].

79. Walsh, B. D. [Egg slits made by Homoptera.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.

Answer to inquiry of C. Dadant; slits in pear-twigs caused by Chloroneura malefica [= Empoasca viridescens]; Proconia [= Oncometopia] undata oviposits in grape-vines.

80. Walsh, B. D. [Plant-lice.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.

Answer to inquiry of J. Flournoy; characters, habits, encmies, and means against Aphididæ.

- 81. Walsh, B. D. [Insects injuring plum-trees.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.
 - Answer to inquiry of A. C. Hammond; *Ægeria* [=Sannina] exitiosa and an elaterid larva attacking plum-trees; the latter probably attracted by decayed matter, the work of the peach borer.
- 82. Walsh, B. D. Prof. Dana and his entomological speculations. Proc. Ent. Soc. Phil., May-June, 1866, v. 6, pp. 116-121.

Rejoinder to J. D. Dana's reply to criticism upon the classification of insects.

83. Walsh, B. D. Clover-worms. < Pract. Ent., 25 June, 1866, v. 1, pp. 82-83.

Summary of published observations concerning Asopia costalis; points which need explanation; probable remedy.

- 84. Walsh, B. D. Click-beetles. < Pract. Ent., 25 June, 1866, v. 1, p. 83.
 - Note to W.'s "Click-beetles;" snapping of Elateridæ; Jaeger's Life of N. Am. Insects and Emmons' N. Y. Report unreliable.
- 85. Walsh, B. D. Cut-worms. < Pract. Ent., 25 June, 1866, v. 1, pp. 85-86.

Definition, food-habits, and means against cut-worms; climbing habits; cut-worms in California. See Nos. 229, 281.

- 86. Walsh, B.D. The canker-worm. Finding a mare's nest. < Pract. Ent., 25 June, 1866, v. 1, p. 87.
 - Criticism of communication in Western Rural; improbability of the report that Anisopteryx [= Paleacrita] vernata deposits her eggs on the ground.
- 87. Walsh, B. D. Driving nails into fruit trees. < Pract. Ent., 25 June, 1866, v. 1, pp. 87-88.

Uselessness of nails as a means against borers; extracts from various sources, with comments.

- 88. Walsh, B. D. The new potato-bug. <Pract. Ent., 25 June, 1866, v. 1, pp. 88-89.
 - Records Doryphora 10-lineata at Bloomington and Athens, Ill.; rate of its eastern progress; need of legislative action against injurious insects.
- 89. Walsh, B. D. [The painted borer.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.

Answer to inquiry of T. C. Wright; sexual difference in the antennæ of Clyvius [= Cyllene] pictus bred from hickory,

90. Walsh, B. D. [Bug allied to the chinch-bug.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.

Answer to inquiry of H. B. Howarth; comparative characters of *Blissus leu-copterus* and an unnamed lygæid of similar habits.

91. Walsh, B. D. [Canker-worms.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.

Answer to inquiry of C. H. W. Wood; means against Anisopteryx.

92. Walsh, B. D. [Apple-tree tineid.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.

Answer to inquiry of C. Cooke; food-plant of Bucculatrix pomifoliella.

93. Walsh, B. D. [Spittle insect.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.

Answer to inquiry of M. V. B. Hathaway; life-habits of species of Aphrophora.

94. Walsh, B. D. [Insects named.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.

Answer to inquiry of E. Hall; occurrence of Doryphora 10-lineata at Athens, Ill.; Hippodamia [= Megilla] maculata devours the eggs of the same; two species of Cassida affect the sweet-potato.

95. Walsh, B. D. [Longicorn borers.] < Pract. Ent., 25 June, 1866, v. 1, p. 90.

Answer to inquiry of Dr. Trimble; food-habits of Orthosoma cylindricum, [= brunneum], Prionus laticollis, and P. imbricornis.

96. Walsh, B. D. [Canker-worms.] < Pract. Ent., 25 June, 1866, v. 1, p. 90.

Answer to inquiry of F. K. Phænix; season of oviposition of Anisopteryx.

97. Walsh, B. D. [White-pine scale.] < Pract. Ent., 25 June, 1866, v. 1, p. 90.

Answer to inquiry of H. Shimer; Aspidiotus [= Chionaspis] pinifolii infesting Pinus strobus.

98. Walsh, B. D. Doctors differ. < Pract. Ent., 30 July, 1866, v. 1, p. 96.

Criticism of communications in agricultural papers; tanners' oil injurious to trees; sulphur plugged in trees does not protect the leaves; sheep in an orchard do not keep away Conotrachelus nenuphar.

99. Walsh, B. D. The grain plant-louse. < Pract. Ent., 30 July, 1866, v. 1, pp. 96-97.

The insect attacking small grains in Georgia is probably a plant-louse.

100. Walsh, B. D. Popular names for insects. < Pract. Ent., 30 July, 1866, v. 1, p. 97.

The use of the same vernacular name for different insects causes uncertainty in identifying and in suggesting means against injurious insects.

101. Walsh, B. D. Scientific names. < Pract. Ent., 30 July, 1866, v. 1, pp. 97-99.

Explains the necessity of using Latin for scientific names.

102. Walsh, B. D. [Grape-vine insects.] < Pract. Ent., 30 July, 1866, v. 1, pp. 99–100.

Answer to inquiry of C. S. Jackson; habits of an undescribed Fidia; injuries to vines by plant-louse (Aphis vitis?); enemies of Aphididæ.

103. Walsh, B. D. [Wire-worms.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.

Answer to inquiry of A. D. Chesebro; summer fallowing as a means against wire-worms.

104. Walsh, B. D. [Bark-lice.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.

Answer to inquiry of L. É. Harmon; life-history and means against Coccide.

105. Walsh, B. D. [Tortoise beetles.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.

Answer to inquiry of C. K. Yant; food-plants and larval habits of Cassidida.

106. Walsh, B. D. [Grape bark-louse.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.

Answer to inquiry of J. Bird; description and ravages of Lecanium [= Pulvinaria] vitis.

107. Walsh, B. D. [Plum bark-louse.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.

Answer to inquiry of O. B. Douglas.; description of Lecanium sp.

108. Walsh, B. D. [Saw-fly eggs.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.

Answer to inquiry of L. D. Morse; position of saw-fly eggs on oak-leaf; characters of the larvæ of *Tenthredinidæ*.

109. Walsh, B. D. [Wheat midge.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.

Answer to inquiry of R. F.; ravages, description, habits, and means against Cecidomyia [=Diplosis] tritici.

110. Walsh, B. D. [Wheat midge.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.

Answer to inquiry of M. H. Boye; Cecidomyia [=Diplosis] tritici not infrequently passes from larval to imaginal state in the same season; means against the same.

111. Walsh, B. D. [Grape galls.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.

Answer to inquiry of J. H. Foster, jr.; description of the galls and larvæ of 'Cecidomyidæ.

112. Walsh. B. D. [Apple-tree web-caterpillar.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.

Answer to inquiry of Marion Hobart; description of the eggs of Clisiocampa americana; food-plants of the same and of Hyphantria textor [= cunea].

- 113. Walsh, B. D. [Grape-vine beetles.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.
 - Answer to inquiry of M. S. Hill; food-habits of the larva and image of *Pelidnota punctata* and of *Anomala lucicola*.
- 114. Walsh, B. D. [Bot-flies.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of J. B. Fisher; habits and hosts of several species of *Estrida*; parasite infesting *Triplax* [= *Tritoma*] thoracica. See No. 153.
- 115. Walsh, B. D. [Spruce-tree saw-flies.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of J. Barratt; description of cocoon and image of Lophyrus abietis; means against the same. See Pract. Ent., v. 1, p. 114.
- 116. Walsh, B. D. [Stings of insects.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of A. O. Brickman; effects of the stings of bees and other insects.
- 117. Walsh, B. D. [Blackberry scale.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of G. E. Brackett; characters of Lecanium sp. infesting blackberry.
- 118. Walsh, B. D. [Eggs in sumach.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of W. M. Smith; eggs of Orchelimum sp.? or Xiphidium sp.? in pith of sumach.
- 119. W[ALSH], B. D. The "new potato-bug" in Maine. < Pract. Ent., 27 August, 1866, v. 1, p. 105.
 - Extract from Maine Farmer, 26 July, 1866, with comments; the larva of Lema trilineata mistaken for that of Doryphora 10-lineata.
- 120. Walsh, B. D. [The army-worm.] < Pract. Ent., 27 August, 1866, v. 1, p. 107.
 - Extract from Western Rural, 21 July, 1866, with comments; army-worm mistaken for canker-worm; need of precision in using popular names.
- 121. W[ALSH], B. D. [Fire-cure for potato-beetles.] < Pract. Ent., 27 August, 1866, v. 1, pp. 107–108.
 - Extract from letter from M. S. Hill, with comments; burning straw effective as a means against *Meloidw* infesting potato-vines; its use against *Doryphora* 10-lineata; hot water as a means against the onion-magget and peach-borer.
- 122. Walsh, B. D. A new humbug. < Pract. Ent., 27 August, 1866, v. 1, pp. 108-110.
 - Exposure of proposed remedy against Cecidomyia destructor; life-habits of the same.
- 123. W[ALSH], B. D. The striped bug. < Pract. Ent., 27 August, 1866, v. 1, p. 110, figs. 1, 2.
 - Extract from Western Rural, 21 July, 1866, with comments; characters, habits, means against, and figures of *Diabrotica vittata* and *D.* 12-punctata. See No. 148.

- 124. Walsh, B. D. [Grape-vine caterpillars.] < Pract. Ent., 27 August, 1866, v. 1, p. 111.
 - Answer to inquiry of B. Borden; description of the early stages of *Procris* [= Acoloithus] falsarius; characters and habits of the species of *Procris*.
- 125. Walsh, B. D. [Squash-vine insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 111.
 - Answer to inquiry of J. Cope; description and habits of *Epilachna borealis*; habits of *Coccinellidæ*; description and means against the larva of *Trochilium cucurbitæ* [= *Melittia ceto*].
- 126. Walsh, B. D. [Injurious insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 111.
 - Answer to inquiry of T. Conard; larval habits and means against Lema trilineata; work of the grain plant-louse (Nectarophora granaria) and of Trochilium cucurbitæ [= Melittia ceto]; mention of some of the enemies of bees.
- 127. Walsh, B. D. [Grape-leaf galls.] < Pract. Ent., 27 August, 1866, v. 1, pp. 111, 112.
 - Answer to inquiry of E. Daggy; description of the leaf-gall of *Phylloxera* vitifoliw [=vastatrix]; reasons for including it in the *Coccidæ*; dipterous enemy of the same. See No. 154.
- 128. Walsh, B. D. [Grape-vine insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 112.
 - Answer to inquiry of W. H. S.; description and means against an undescribed gall on the tendrils and leaf-stalks of a grape-vine; life-habits of parasitic *Chalcidide*. See No. 160.
- 129. Walsh, B. D. [Grape-vine beetles.] < Pract. Ent., 27 August, 1866, v. 1, pp. 112, 113.
 - Answer to inquiry of W. J. Lawrence; food-habits of *Pelidnota punctata* and of *Colaspis flavida*.
- 130. Walsh, B. D. [Corydalis cornutus.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of J. S. Lewis; natural history and habits of Corydalis cornutus.
- 131. Walsh, B. D. [Tobacco-moth.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of F. W. Noble; transformations and food-plants of Sphinx [=Protoparce] carolina; food-plants of S. 5-maculata [=P. celeus].
- 132. Walsh, B. D. [Datana ministra.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of S. S. Lacy; food-plants and transformations of *Datana* ministra.
- 133. Walsh, B. D. [Grape-vine and fuschia beetles.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of F. K. Phænix; Colaspis flavida injurious to grape-vine shoots; Haltica exapta [=carinata] destructive to fuschia; means against flea-beetles.

134. Walsh, B. D. [Blister-beetles.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.

Answer to inquiry of J. Barber; food-plants of Lytta marginata [=Epicauta cinerea], L. [=E.] vittata, L. atrata [=E. pennsylvanica], and L. cinerea [=Macrobasis unicolor].

135. Walsh, B. D. [Colaspis flavida; Lema trilineata.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.

Answer to inquiry of J. F. Benner; Colaspis flavida destructive to grape-vines; distribution and method of breeding Lema trilineata.

136. Walsh, B. D. [Green worms on gooseberries and currants; *Lema trilineata.*] < Pract. Ent., 27 August, 1866, v. 1, p. 114.

Answer to inquiry of H. Burt; gooseberries and currants injured by larva of a saw-fly?; effects of excessive rain, heat, and drought; food-plants of Lema trilineata.

137. Walsh, B. D. [Galls.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.

Answer to inquiry of H. Shimer; note of galls made by Rhodites radicum, Trypeta solidaginis, and Thelaxes [= Colopha] nlmicola.

138. Walsh, B. D. [Potato and corn insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.

Answer to inquiry of T. H. Parsons; searcity of Lema trilineata in Illinois; Doryphora 10-lineata destroyed by turkeys, but not by fowls; Gortyna sp.? injurious to young Indian corn; rules for breeding insects.

139. Walsh, B. D. [Cocoons on apple-trees.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.

Answer to inquiry of M. S. Hill; description of cocoon and image of Attacus cecropia; the interaction of organisms.

140. Walsh, B. D. Imported insects; the gooseberry saw-fly. <Pract. Ent., 29 September, 1866, v. 1, pp. 117-125, fig.</pre>

List of some injurious European insects imported into America and of American insects imported into Europe; reasons for the increase and abundance of noxious European insects in America, and why American species do not flourish in Europe; natural history, description, and means against Nematus ventricosus [=ribesii]; figure of currant-leaf, showing method of deposition of eggs and work of larva; natural history and description of larva and imago of Pristiphora grossulariæ n. sp.; number of saw-flies infesting the gooseberry and currant in Europe.

141. Walsh, B. D. [Borers.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.

Answer to inquiry of W. H. R. Lykins; western distribution of *Clytus* [= *Cyllene*] robinia.

142. Walsh, B. D. [Wheat midge.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.

Answer to inquiry of C. P. Wickersham; destructiveness of Cecidomyia [=Diplosis] tritici; nature of its cocoon.

143. Walsh, B. D. [Crickets.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.

Answer to inquiry of Marion Hobart; characters and food-habits of Acheta [=Gryllus] abbreviatus; katydids not exclusively vegetarians; Œcanthus niveus found feeding on plant-lice.

- 144. Walsh, B. D. [Walking-stick.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of L. W. Taylor; characters and habits of Spectrum [=Diaphcromera] femorata.
- 145. Walsh, B. D. [Apple-tree insects.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of H. B. Beegle; characters and food-plants of *Limenitis* ursula; description and food-plants of the larva of *Lagoa opercularis*; eharacters of the imago.
- 146. Walsh, B. D. [Mites on grasshoppers.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of W. Riddell; Astoma [= Trombidium] locustarum infesting Caloptenus femur-rubrum.
- 147. W[ALSH], B. D. Grasshoppers and locusts. < Pract. Ent., October, 1866, v. 2, pp. 1-5, 22.
 - Description, migration, and ravages of *Caloptenus spretus*; limits of its range and means of preventing its increase; eonfusion caused by the use of popular names of insects. On p. 22 a "note," accidentally omitted, gives the comparative length of the wing-covers and antennæ of *Caloptenus spretus* and *C. femur-rubrum*.
- 148. W[ALSH], B. D. The striped cucumber-bug. <Pract. Ent., October, 1866, v. 2, p. 5.

Diabrotica vittata attacks German asters. See No. 123.

- 149. W[ALSH], B. D. Fire-blight. < Pract. Ent., October, 1866, v. 2, p. 7.
 - Extract from Horticulturist, with comments; cause of fire-blight unknown; not produced by Scolytus [= Xyleborus] pyri.
- 150. W[ALSH], B. D. The Hessian fly. < Pract. Ent., October, 1866, v. 2, p. 7.
 - Extract from Colman's Rural World, with comments; late sowing as a means against Cecidomyia destructor.
- 151. [Walsh, B. D.] [Injurious caterpillars; plant-lice enemies.] < Pract. Ent., October, 1866, v. 2, pp. 7-8.
 - Answer to inquiry of T. McGraw; habits, descriptions, and food-plants of larva of Dryocampa senatoria and of Datana ministra; food-habits of larva allied to D. ministra; description and habits of larva of Scymnus sp.; S. hamorrhous? bred from gall of Thelaxes [=Colopha] ulmicola.
- 152. [Walsh, B. D.] [Tomato-worm.] < Pract. Ent., October, 1866, v. 2, p. 8.
 - Answer to inquiry of A. A. Jackson; description of the larva and pupa of *Protoparce celeus*; food-plants and harmlessness of the larva.

153. [Walsh, B. D.] [Dipterous larva on swallows.] < Pract. Ent., October, 1866, v. 2, p. 8.

Answer to inquiry of J. B. Fisher; larvæ found on swallows (see No. 114) belong to the Muscidæ and not to the Estridæ; extract from letter of Osten Sacken on larval habits of Muscidæ found with birds.

154. [Walsh, B. D.] [Bark-louse enemy.] < Pract. Ent., October, 1866, v. 2, p. 8.

Answer to inquiry of E. Daggy; see No. 127; according to Osten Sacken the bark-louse enemy belongs to *Leucopis*; systematic position of the genus.

155. [Walsh, B. D.] [Parasitized tomato-worm.] < Pract. Ent., October, 1866, v. 2, pp. 8-9.

Answer to inquiry of T. C. Wright; habits of *Microgaster*, parasitic on *Protoparce celeus*; habits of ichneumonized larvæ.

156. [Walsh, B. D.] [Beetles on buttercup and azalea.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of C. H. Peck; identification of Galeruca [=Adimonia] rufosanguinea, found on Ranunculus acris, and of Prasocuris varipes on Azalea. nudiflora; synoptic table, by J. L. Leconte, of the species of Prasocuris.

157. [WALSH, B. D.] [Flying-bug.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of E. E. Sheldon; Aphodius mistaken for Cecidomyia destructor.

158. [WALSH, B. D.] [Parasitic cocoon.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of H. W. Howarth; description of cocoons made by Microgaster and Pezomachus; habits of Ichneumonidæ.

159. [Walsh, B. D.] [Oak-bark louse.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of T. Meehan; undetermined Coccus on red-oak.

160. [Walsh, B. D.] [Grape-galls.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of W. H. S.; see No. 128; larva of *Leucopis* sp., an enemy to grape-vine gall-louse; probable identity of the maker of the tendril and leaf-galls; varieties of grape infested by *Phylloxera vastatrix*.

161. [WALSH, B. D.] ["Saddle-back."] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of T. M. Harvey; food-plants, description, and urticating properties of *Empretia stimulea*.

162. [Walsh, B. D.] [Corn-worm.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of J. B. Ellis; method of work of two lepidopterous larvæ.

163. [Walsh, B. D.] [Insects named.] < Pract. Ent., October, 1866, v. 2, p. 10.

Answe to inquiry of W. C. Fish; comparative characters of *Entilia sinuata* and E, $\Gamma = Publilia$ concava.

- 164. [Walsh, B. D.] [Grape insects.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of J. H. Garman; description and habits of larva of Procestina [= Harrisina] americana; Procestia [= Oncometopia] undata and several species of Erythroneura [= Typhlocyba] as grape enemies.
- 165. [WALSH, B. D.] [Datana ministra.] < Pract. Ent., October, 1866, v. 2, p. 10. -
 - Answer to inquiry of J. B.; increase of injuries caused by *Datana ministra*; distinctive characters of the larvæ of *Geometrida*.
- 166. [Walsh, B. D.] [Iulus marginatus.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of T. Wiggins; characters and food of *Iulus* [= Spirobolus] marginatus.
- 167. [Walsh, B. D.] [Midge.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of W. G. Morris; characters and habits of the larva and imago of *Chironomus* sp.
- 168. [Walsh, B. D.] [Rat-tailed larva.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of J. A. Lapham; characters and habitats of the larvæ of rat-tailed Syrphide.
- 169. [Walsh, B. D.] [Beetle on thistle.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of W. S. Robertson; characters and food-plants of *Phyllobrotica* [= Diabrotica] longicornis.
- 170. [Walsh, B. D.] [Grape-vine *Aphis.*] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of H.; grape-vine injured by Aphis vitis.
- 171. W[ALSH], B. D. The new potato-bug. <Pract. Ent., November, 1866, v. 2, pp. 13-16, fig.
 - Eastern progress of *Doryphora* 10-lineata; rate of travel; hand-picking the only effective remedy; description of a horse machine for their destruction; pecuniary loss inflicted by the species.
- 172. W[ALSH], B. D. The canker-worm < Pract. Eut., November, 1866, v. 2, p. 16.
 - Tarring trees effective against Paleacrita vernata.
- 173. W[ALSH], B. D. The canker-worm again. < Pract. Ent., November, 1866, v. 2, pp. 16-17.
 - Critical review of article in New England Farmer; value of tarring as a means against Paleacrita vernata.
- 174. Walsh, B. D. Notes by Benj. D. Walsh. < Pract. Ent., November, 1866, v. 2, pp. 19-20.
 - Remarks on statement of H. Shimer in his paper, "The grape leaf gall-coccus" (Pract. Ent., pp. 17-19); number of eggs to each \mathfrak{D} ; tendril and leafgalls probably produced by the same insect; old leaves entirely free from bark-lice by the end of summer; characters of the pupa of Leucopis; systematic position and food-habits of Thysanoptera; criticism of paper by H. Shimer in Prairie Farmer, 3 November, 1861.

175. W[ALSH], B. D. The striped bug. <Pract. Ent., November, 1866, v. 2, p. 20.

Remarks on note by A.; effects of plaster as a means against *Diabrotica* vittata.

- 176. [Walsh, B. D.] [Currant insects; *Psocus venosus.*] < Pract. Ent., November, 1866, v. 2, p. 20.
 - Answer to inquiry of I. Hicks; occurrence of *Pristiphora grossulariæ* in the Eastern States; *Ellopia* [= *Eufitchia*] *ribearia* injurious to currant; date of introduction of "sulphur-cure" on peach-trees; characters and habits of *Psocus venosus*.
- 177. [Walsh, B. D.] [Worm in apple.] < Pract. Ent., November, 1866, v. 2, pp. 20-21.

Answer to inquiry of C. Ward; description and ravages of larva of Trypeta pomonella?

- 178. [Walsh, B. D.] [Insects attracted to light.] < Pract. Ent., November, 1866, v. 2, p. 21.
 - Answer to inquiry of Practical Pomologist; conditions favorable for attracting insects to light; kinds most frequently attracted; the proportion of injurious species attracted.
- 179. [Walsh, B. D.] [Zebra caterpillar.] <Pract. Ent., November, 1866, v. 2, p. 21.

Answer to inquiry of J. H. Parsons; description of the larva of *Mamestra* picta; times of transformation.

- 180. [Walsh, B. D.] [Inquiries answered.] < Pract. Ent., November, 1866, v. 2, p. 21.
 - Answer to inquiry of J. Pettit; identification of several beetles; galls on golden-rod made by Euryptychia saligneana and Trypeta solidaginis; characters of short-winged form of Micropus [=Blissus] leucopterus; its occurrence in Canada.
- 181. [Walsh, B. D.] [Worm in corn.] < Pract. Ent., November, 1866, v. 2, pp. 21-22.

Answer to inquiry of J. B. Ellis; description of larva, habits, transformations, and means against an undetermined noctuid.

- 182. [Walsh, B. D.] [Bag-worms.] < Pract. Ent., November, 1866, v. 2, p. 22.
 - Answer to inquiry of C. P. Wickersham; characters and synonymy of *Thyridopteryx ephemeræformis*.
- 183. [WALSH, B. D.] [Ichneumon cocoons.] <Pract. Ent., November, 1866, v. 2, p. 22.

Answer to inquiry of S. A. N.; characters of cocoons of Microgaster sp.?

184. Walsh, B. D. Note. < Proc. Ent. Soc. Phil., December, 1866, v. 5, p. 260.

Withdraws assertion attributed to C. R. Osten Sacken.

185. W[ALSH], B. D. The old-fashioned potato-bugs. <Pract. Ent., December, 1866, v. 2, pp. 25-27, figs.

Natural history and means against Lema trilineata, Lytta [= Epicauta] vittata, L. atrata [= E. pennsylvanica], L. marginata [= E. cinerea], and L. cinerea [=Macrobasis unicolor]; figures L. trilineata and E. vittata.

- 186. W[ALSH], B. D. Klippart's wheat plant. < Pract. Ent., December, 1866, v. 2, pp. 27-29.
 - Adverse criticism of the entomological portion of the above-named book.
- 187. W[ALSH], B. D. Trimble's insect enemies of fruit and fruit trees. <Pract. Ent., December, 1866, v. 2, pp. 29-30.

 Favorable review of the above-named book.
- 188. W[ALSH], B. D. Answer to the above, by B. D. W. < Pract. Ent., December, 1866, v. 2, pp. 31-33, figs.
 - Answers to questions asked in J. S. Houghton's "Insects in the Orchard" (Pract. Ent., pp. 30-31); food-habits and hibernation of Conotrachelus nenuphar; comparative characters and figures of Aspidiotus conchiformis [=Mytilaspis pomorum] and Coccus harrisii [=Chionaspis furfurus]; natural history of C. harrisii; means against and enemies of the two species; figures Chilocorus bivulnerus and Hippodamia [=Megilla] maculata; food-habits and synonymy of Lytta [=Pomphopæa] sayi; flowing as a means against root-feeding insects; inefficacy of salt, lime, ashes, etc., against the same; insects attacking evergreens not likely to injure fruit trees; cannibal habits of spiders; habits of bees and wasps.
- 189. [Walsh, B. D.] [Cicada; gooseberry-worm.] < Pract. Ent., December, 1866, v. 2, p. 33.
 - Answer to inquiry of M. S. Hill; Cicada [= Tibicen] septendecim compared with an undescribed species; ravages of Pristiphora grossulariæ? on gooseberry bushes.
- 190. [Walsh, B. D.] [Vespidæ.] < Pract. Ent., December, 1866, v. 2, pp. 33, 34.
 - Answer to inquiry of M. Hobart; life-habits of Vespidæ.
- 191. [Walsh, B. D.] [Meal-worms: Timber-borers.] < Pract. Ent., December, 1866, v. 2, p. 34.
 - Answer to inquiry of M. C. D.; characters, food-habits, and means against *Tenebrio molitor* and *T. obscurus*; characters of *Ptinus brunneus* infesting pine timber; corrosive sublimate as a means against timber insects.
- 192. [Walsh, B. D.] [Corn-worm.] < Pract. Ent., December, 1866, v. 2, p. 34.
 - Answer to J. B. Ellis; number of broods of *Heliothis armigera* in the Northern and Southern States.
- 193. [Walsh, B. D.] [Thousand-legged worms.] < Pract. Ent., December, 1866, v. 2, pp. 34-35, fig.
 - Answer to inquiry of G. W. Robinson; description and figure of *Iulus multi-striatus* n. sp. [= Cambala annulata]; habits and means against the same.
- 194. [Walsh, B. D.] [Cranberry-galls, etc.] <Pract. Ent., December, 1866, v. 2, p. 35.
 - Answer to inquiry of W. C. Fish; occurrence of galls made by Cecidomyia sp. on leaves of cranberry; identification of several insects.
- 195. [Walsh, B. D.] Another humbug. <Pract. Ent., December, 1866, v. 2, p. 35.
 - Extract, with comments, from Cultivator and Country Gentleman, 22 November, 1866, exposing worthlessness of P. B. Sheldon's composition for destroying borers,

196. [Walsh, B. D.] Notices. < Pract. Ent., December, 1866, v. 2, p. 35.

Mention of several agricultural journals.

- 197. Walsh, B. D. On the insects, coleopterous, hymenopterous, and dipterous, inhabiting the galls of certain species of willow. Part 2d and last. <Proc. Ent. Soc. Phil., Décember, 1866–January, 1867, v. 6, pp. 223–288.
 - See No. 46 for Part 1st. Supplementary notes on the Cecidomyidæ (gall-makers and guest-flies); criticism of statements of Fitch and Harris; discussion of structure and habits of Tenthredinidæ and insects generally; synopsis of tenthredinidous willow-galls; descriptions of the galls, their makers and inquilines; for the new species described, see the Systematic Index; remarks on species and varieties; figures fore-wings of Tenthredo and Ichneumon.
- 198. W[Alsh], B. D. Plant-lice—their friends and enemies. < Pract. Ent., January, 1867, v. 2, pp. 37-44, figs.
 - Method of work and mode of reproduction of Aphidida; their relations to ants; plant-lice enemies; figures Aphis mali and several enemies of plant-lice; means against Aphidida.
- 199. W[ALSH], B. D. Birds *versus* insects. < Pract. Ent., January, 1867, v. 2, pp. 44-47.

Insectivorous birds not necessarily beneficial; a bird must be shown to eat at least thirty (30) times as many injurious insects as it does beneficial ones before it can be considered useful.

200. [Walsh, B. D.] [Insects named.] < Pract. Ent., January, 1867, v. 2, p. 47.

Answer to inquiry of J. Pettit; identification of many species, mostly beetles; characters and food-habits of Clerus nigripes [= 4-guttatus]; habitat of Leptura capitata and Tingis [=Corythuca] ciliata.

201. [Walsh, B. D.] [Apple-tree insects.] < Pract. Ent., January, 1867, v. 2, p. 47.

Answer to inquiry of M. Hobart; increasing distribution of Aspidiotus conchiformis [= Mytilaspis pomorum]; enemies of the same; characters of the eggs of Aphis mali.

202. [Walsh, B. D.] [Work in entomology.] < Pract. Ent., January, 1867, v. 2, p. 47.

Answer to G. Scarborough; advice for the study of entomology by beginners.

- 203. W[ALSH], B. D. The true *Thrips* and the bogus *Thrips*. < Pract. Ent., February, 1867, v. 2, pp. 49–52, figs.
 - Natural history, characters, habits, and figures of *Thrips* sp.; characters of the larva and image of *Haltica chalybea*; figure of image; habits of species of *Erythroneura* [= *Typhlocyba*]; description and figures of *E.* [= *T.*] vitis.
- 204. W[ALSH], B. D. Universal remedies. < Pract. Ent., 1867, v. 2, p. 52.

Exposure of proposed universal remedy against fruit-tree insects.

205. W[ALSH], B. D. Poisoning noxious insects. < Pract. Ent., February, 1867, v. 2, pp. 52-53.

Sugaring with poison as a means against Noctuide.

- 206. W[ALSH], B. D. Remarks by B. D. W. < Pract. Ent., February, 1867, v. 2, p. 54.
 - Comments on S. S. Rathvon's "Bag-worms" (Pract. Ent., pp. 53-54); distribution and food-plants of Thyridopteryx ephemeræformis.
- 207. W[ALSH], B. D. Habits of the tree-cricket (*Ecanthus niveus*). < Pract. Ent., February, 1867, v. 2, p. 54, figs.
 - Characters of eggs and method of deposition; characters, food-habits, and figures of the imago.
- 208. [Walsh, B. D.] Importing European parasites. < Pract. Ent., February, 1867, v. 2, pp. 54-55.
 - Imaginary correspondence between Asa Fitch and John Curtis on the importation of parasitic insects.
- 209. [Walsh, B. D.] [Screw-worm.] < Pract. Ent., February, 1867, v. 2, p. 55.
 - Answer to inquiry of L. D. Morse; nndetermined larva found in osageorange seed; hominivorous habits of Lucilia macellaria.
- 210. [Walsh, B. D.] [Attacus cecropia.] < Pract. Ent., February, 1867, v. 2, p. 55.
 - Answer to inquiry of T. T. Smith; characters of larva and image of Attacus cecropia; food-plants; method of issnance from cocoon; nature of fluid discharged when emerging from cocoon.
- 211. [WALSH, B. D.] [Apple-worm.] < Pract. Ent., February, 1867, v. 2, pp. 55-56.
 - Answer to inquiry of W. C. Fish; characters of the pupa of Sciara mali.
- 212. [Walsh, B. D.] [Oak-gall.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of W. Muir; characters of an undetermined cynipid gall on oak.
- 213. [Walsh, B. D.] [Cicada: Grape-vine Procris.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of M. S. Hill; variations in the image of Cicada [= Tibicen] septendecim and in the song of the same; characters of the larva of Procris [= Harrisina] americana.
- 214. [Walsh, B. D.] [Nitidulidæ.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of W. H. S.; food-habits of Ips fasciatus and Nitidula bipustulata.
- 215. [Walsh, B. D.] [Turnip enemy.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of F. T. Pember; characters and food-habits of *Polydesmus complanatus*.
- 216. [WALSH, B. D.] [Wheat midge.] < Pract. Ent., February, 1867, v. 2, p. 57.
 - Answer to inquiry of C. P. Wickersham; nature of the membrane enveloping the full-grown larva of *Diplosis tritici*,

217. [Walsh, B. D.] [Queries answered.] < Pract. Ent., February, 1867, v. 2, p. 57.

Answer to inquiry of R. Middleton; toads, frogs, and spiders beneficial; effects of the stings of various insects.

- 218. [Walsh, B. D.] [Scales.] < Pract. Ent., February, 1867, v. 2, p. 57.
 - Answer to inquiry of M. M. S.; characters and ravages of scales on oleander and orange; means against *Coccida*.
- 219. [Walsh, B. D.] [Hundred-legged worm.] < Pract. Ent., February, 1867, v. 2, p. 57.
 - Answer to inquiry of S. P. Monks; identification of *Polydesmus virginiensis*; distinctive characters of Myriapoda.
- 220. [Walsh, B. D.] [Bark-borers.] < Pract. Ent., February, 1867, v. 2, pp. 57-58.

Answer to inquiry of C. V. Riley; synoptic table and food-habits of several species of *Scolytus*; descriptions of *S. fagi* n. sp. and *S. caryw* n. sp. [=4-spinosus].

- 221. [Walsh, B. D.] [Aleurodes.] < Pract. Ent., February, 1867, v. 2, p. 58.
 - Answer to inquiry of D. F. C.; characters and food-plants of an undescribed species of *Aleurodes*.
- 222. W[ALSH], B. D. The critic criticized. <Pract. Ent., February, 1867, v. 2, p. 58. Answer [by C. V. Riley]. <Prairie Farmer, 16 March, 1867, [v. 35], n. s., v. 19, p. 169. S.-b., No. 2, p. 58. Criticism of article by C. V. Riley (Prairie Farmer, 19 January, 1867, [v. 35], n. s., v. 19, p. 37).
- 223. [Walsh, B. D.] Obituary. '<Pract. Ent., February, 1867, v. 2, p. 58.

Notice of the late Brackenridge Clemens.

- 224. W[ALSH], B. D. Wire-worms. < Pract. Ent., March, 1867, v. 2, pp. 61-62, figs.
 - Characters of Myriapoda and larvæ of *Elateridæ*, popularly called wire-worms; food-habits of larva of *Ludius attenuatus*; figure of larva and imago of the same; confusion caused by the use of the term "wire-worms" by correspondents of the Country Gentleman.
- 225. W[ALSH], B. D. Imported insects.—The onion-fly. <Pract. Ent., March, 1867, v. 2, p. 64, figs.
 - Figures of larva and imago of Anthomyia ceparum and of Ortalis [= Tritoxa] flexa; comparative characters of imagos; distribution and means against the two species.
- 226. W[ALSH], B. D. [Means against cut-worms.] < Pract. Ent., March, 1867, v. 2, p. 66.
 - Recommends, in reply to J. Townley's inquiry, sugaring with poisoned molasses as a means against cut-worms.

- 227. W[ALSH], B. D. Entomology indeed run mad. <Pract. Ent., March, 1867, v. 2, pp. 66, 67.
 - Reprint of "The bud-worm" (N. C. Rural Journ., September, 1866), with criticism and correction of some of the errors.
- 228. W[ALSH], B. D. The imported gooseberry saw-fly. < Pract. Ent., March, 1867, v. 2, p. 67.
 - Reason for re-appearance in June and for the non-appearance of the second broad of Nematus ventricosus [=ribesii] in New York in 1866.
- 229. W[ALSH], B. D. Tree cut-worms. <Pract. Ent., March, 1867, v. 2, p. 67.
 - Cut-worm described (Pract. Ent., v. 1, pp. 85-86) identified as *Hadena chenopodii* [= Mamestra trifoli]. See Nos. 85, 281.
- 230. W[ALSH], B. D. Doctoring fruit trees again. <Pract. Ent., March, 1867, v. 2, pp. 67, 68.
 - Extract from Industrial Gazette, 15 December, 1866, with criticism; calomel, sulphur, and substances insoluble in water not absorbed into the circulation of trees.
- 231. W[ALSH,] B. D. The grape-vine Colaspis (Colaspis flavida Say). <Pract. Ent., March, 1867, v. 2, pp. 68-69, fig.
 - Ravages of *Colaspis flavida*; comparison with allied species; remarks on the law of priority and on certain insects becoming suddenly injurious; figures *C. flavida*.
- 232. W[ALSH], B. D. Another universal remedy. <Pract. Ent., March, 1867, v. 2, p. 69.
 - Extract from reprint (?) in Cultivator and Country Gentleman, 10 January, 1867, with criticism of proposed remedy against insects injurious to trees.
- 233. W[ALSH], B. D. Borers. The plug-ugly theory. <Pract. Ent., March, 1867, v. 2, p. 69.
 - Criticism of theory that a plug driven into a hole in a tree is death to borers.
- 234. W[ALSH], B.D. A mass of mistakes. < Pract. Ent., March, 1867, v. 2, p. 70.
 - Correction of errors in "Remedy for the borer" (Prairie Farmer, 9 February, 1867).
- 235. W[ALSH], B. D. Hop-growing in the West. < Pract. Ent., March, 1867, v. 2, p. 70.
 - Extract from Country Gentleman, 31 January, 1867, showing ravages of *Phorodon humuli* in Vermont; danger of its introduction into the Western States.
- 236. W[ALSH], B. D. Thousand-legged worms. <Pract. Ent., March, 1867, v. 2, p. 70.
 - Extract from letter of J. H. Parsons, with comments; injuries to onions by Iulus multistriatus [= Cambala annulata]; I. multistriatus undistinguishable from I. cæruleo-cinctus.

237. W[ALSH], B. D. Fighting the Curculio. <Pract. Ent., March, 1867, v. 2, p. 71.

Extract from Genesee Farmer for 1853, p. 125; ridicule of additions to "jarring" as a means against Conotrachelus nenuphar.

238. W[ALSH], B.D. ["Bushels of butterflies."] < Pract. Ent., March, 1867, v. 2, p. 71.

Extract from Cincinnati Gazette, August, 1866, with comments; occurrence of dead imagos of Attacus cecropia? in large numbers ("10 bushels") in Newport, Ohio; food-plants of the larva.

239. [Walsh, B. D.] [Seab on potato.] < Pract. Ent., March, 1867, v. ; 2, pp. 71-72.

Answer to inquiry of T. L. J. Baldwin; account of blister-like scabs on potatoes, probably caused by *Sciara* sp.; means against the same.

240. [Walsh, B. D.] [Mass of eggs.] < Pract. Ent., March, 1867, v. 2, p. 72.

Answer to inquiry of E. Daggy; egg-masses of Clisiocampa americana; food-plants of C. americana and C. sylvatica [=disstria].

241. [Walsh, B. D.] [Food of katydids.] < Pract. Ent., March, 1867, v. 2, p. 72.

Answer to inquiry of C. M.; food of Microcentrum retinervis largely carnivorous.

242. [Walsh, B. D.] [Caterpillar nest on wild cherry.] < Pract. Ent. March, 1867, v. 2, p. 72.

Answer to inquiry of H. Morey; undetermined larva on wild cherry, foodplants of *Hyphantria textor* [=eunea].

243. [Walsh, B. D.] [Insect eggs.] < Pract. Ent., March, 1867, v. 2, p. 72.

Answer to inquiry of P. Ferris; undetermined eggs of Lepidoptera and Hemiptera on apple-tree twigs.

244. [WALSH, B. D.] [Attacus polyphemus, etc.] < Pract. Ent., March, 1867, v. 2, pp. 72-73.

Answer to inquiry of M. M. S.; variations in Attacus [= Telca] poluphemus; habits and times of transformations of Arctia [= Pyrrharctia] isabella; references to history of Dryocampa rubicunda.

245. [Walsh, B. D.] [Cossus robiniæ, etc.] < Pract. Ent., March, 1867, v. 2, p. 73.

Answer to inquiry of J. Townley; food-habits and injuries of Cossus robiniæ; result of freezing insects; characters of some wood-borers.

246. [Walsh, B. D.] [Museum pests.] < Pract. Ent., March, 1867, v. 2, p. 73.

Answer to inquiry of F. L. Van Arsdale; means against Dermestidæ.

247. [Walsh, B. D.] [Katydid eggs.] < Pract. Ent., March, 1867, v. 2, p. 73.

Answer to C. M. B.; characters of eggs of Microcentrum retinervis.

248. [Walsh, B. D.] [Squash-vine borer.] < Pract. Ent., March, 1867, v. 2, p. 73.

Answer to inquiry of E. Nason; ravages and means against Trochilium cucurbita [= Melittia ceto].

- 249. [Walsh, B. D.] [Apple-twig borers, etc.] < Pract. Ent., March, 1867, v. 2, pp. 73–74.
 - Answer to inquiry of A. W. Brumbaugh; work of Bostrichus [= Amphicerus] bicaudatus; characters and habits of Twniopteryx fasciata; ravages, habits, and means against Macrodactylus subspinosus.
- 250. [Walsh, B. D.] [Hessian fly.] < Pract. Ent., March, 1867, v. 2, p. 74.

Answer to inquiry of E. E. Sheldon; hibernation of the larva of Cecidomyia destructor.

251. [Walsh, B. D.] [Tree-cricket.] < Pract. Ent., March, 1867, v. 2, p. 74.

Answer to inquiry of J. M. Cole; food-habits of Œcanthus niveus; characters of its eggs.

252. [Walsh, B. D.] [Tineids in bee-hives.] < Pract. Ent., March, 1867, v. 2, p. 74.

Answer to inquiry of J. H. Hunt; characters of case and larva of undetermined tineid.

253. W[ALSH], B. D. Jumping to conclusions. <Pract. Ent., March, 1867, v. 2, p. 74.

Criticism of article "The potato-bug" (Wisconsin Farmer, 2 March, 1867).

254. W[ALSH], B. D. The common Curculio and its allies. <Pract. Ent., April, 1867, v. 2, pp. 75-81.

Characters of the Rhyncophora; characters, ravages, life-history, food-habits, number of broods, and means against Conotrachelus nenuphar; ravages, habits, characters, and means against Anthonomus prunicida [= Coccotorus scutellaris] and A. 4-gibbus; mention of additional injurious weevils; comparative characters of the three species.

255. W[ALSH], B. D. Remarks by B. D. W. < Pract. Ent., April, 1867, v. 2, p. 82.

Comments on C. V. Riley's article, "The imported apple-tree bark-louse" (Prairie Farmer, 23 March, 1867, [v. 35], n. s., v. 19, p. 184).

256. W[ALSH], B. D. Confessing the corn. < Pract. Ent., April, 1867, v. 2, p. 82.

Acknowledgment of mistake in quoting from Prairie Farmer.

257. [Walsh, B. D.] [Bibio albipennis.] < Pract. Ent., April, 1867, v. 2, p. 83.

Answer to inquiry of H. Burt; habits of larva of Bibio albipennis.

258. [Walsh, B. D.] [Lice on horses.] < Pract. Ent., April, 1867, v. 2, p. 83.

Answer to inquiry of L. West; comparative characters of Pediculina and Mallophaga.

259. [Walsh, B. D.] [Cocoon on wild cherry.] < Pract. Ent., April, 1867, v. 2, p. 83.

Answer to inquiry of A. A. Baker; sexual differences of Attacus promethea; food-plauts of its larva.

260. [Walsh, B. D.] [Lepidopterous case-bearer.] < Pract. Ent., April, 1867, v. 2, p. 83.

Answer to inquiry of J. H. Hunt; larval characters of case-bearing Lepidoptera.

261. [Walsh, B. D.] ["Thousand-legged worms."] < Pract. Ent., April, 1867, v. 2, p. 83.

Answer to inquiry of J. H. Parsons; *Inlus virgatus* eating dead seeds; injurious habits of species of *Polydesmus*.

262. [Walsh, B. D.] [Toads: Glorified squash-bug.] < Pract. Ent., 1867, v. 2, p. 83.

Answer to inquiry of F. C. Hill; food of toads; poisonousness of *Prionotus* novenarius [= Prionidus cristatus].

263. [Walsh, B. D.] [Galls on blackberry.] < Pract. Ent., April, 1867, v. 2, pp. 83-84.

Answer to inquiry of J. H. Tice; identification of the gall of *Diastrophus* nebulosus; genera of *Cynipidw* confined to single plants.

264. [Walsh, B. D.] [Bark-lice.] < Pract. Ent., April, 1867, v. 2, p. 84.

Answer to inquiry of Dr. Houghton; comparative harmlessuess of Coccus harrisii [=Chionaspis furfurus]; means against Coccidæ.

265. [Walsh, B. D.] [Plant-lice.] < Pract. Ent., April, 1867, v. 2, p. 84.

Answer to inquiry of E. Orton; description of eggs of *Lachnus strobi*; means against *Aphidida*.

266. [Walsh, B. D.] [Bag-worms.] < Pract. Ent., April, 1867, v. 2, pp. 84-85.

Answer to inquiry of J. Murphy; food-habits and means of dispersal of Thyridopteryx cphemeræformis.

267. [Walsh, B. D.] [Insects injurious to grape.] < Pract. Ent., April, 1867, v. 2, p. 85.

Answer to inquiry of J. Wood; supposed Curculio punctures iu grapes.

268. [Walsh, B. D.] [Attacus promethea.] < Pract. Ent., April, 1867, v. 2, p. 85.

Answer to inquiry of J. B. Lyon; mention of undetermined parasite of Attacus promethea.

269. [Walsh, B. D.] [Borer in hickory.] < Pract. Ent., April, 1867, v. 2, p. 85.

Answer to inquiry of B. Norris; characters of young and mature larva of Clytus [= Cyllene] pictus.

270. [WALSH, B. D.] [Apple-tree plant-lice.] < Pract. Ent., April, 1867, v. 2, p. 85.

Answer to inquiry of W. W. Linn; identification of eggs of Aphis mali; enemies of the same.

- - Answer to inquiry of I. Hicks; parasites and encmies of undescribed barklouse on tulip-tree; Thyridopteryx ephemeræformis destructive to evergreens.
- 272. W[ALSH], B. D. The grape-vine Fidia. (Fidia viticida, new species.) < Pract. Ent., May, 1867, v. 2, pp. 87–88, fig.
 - Ravages, description, and figure of Fidia viticida n. sp.; modification of the clasping organs of insects.
- 273. W[ALSH], B. D. Enemies of the rice crop. <Pract. Ent., May, 1867, v. 2, p. 89.
 - Injuries caused by and means adopted against the water-weevil [=Lissorhop-trus simplex] and the rice-grub = [Chalepus trachypygus.]
- 274. [Walsh, B. D.] The canker-worm once more. < Pract. Ent., May, 1867, v. 2, p. 89. Abstract: < Cultivator and Country Gentleman, 6 June, 1867, v. 29, p. 370.
 - Extract from and review of W. G. C.'s "My trials with the canker-worms: remedy" (Iowa Homestead, 13 March, 1867); sorghum as a means against Anisopteryx.
- 275. W[ALSH], B. D. None so blind as those who shut their eyes. < Pract. Ent., May, 1867, v. 2, pp. 89–90.
 - Criticism of error by editors of agricultural papers; rate of eastward progress of Doryphora 10-lineata.
- 276. W[ALSH], B. D. Self-taught entomologists. <Pract. Ent., May, 1867, v. 2, pp. 91-92.
 - Criticism of misstatements found in agricultural papers; first occurrence of Doryphora 10-lineata in Illinois in 1864.
- 277. W[ALSH], B. D. Pear-tree and apple-tree insects. < Pract. Ent., May, 1867, v. 2, pp. 92-93.
 - Food-plants of Attacus cecropia, Clisiocampa americana, and Saperda bivittata [= candida].
- 278. W[ALSH], B. D. More universal remedies. <Pract. Ent., May, 1867, v. 2, p. 93. .
 - Reprint and criticism of "Destruction of insects" (Mo. Rept. Bureau Agric., February, 1867, p. 60).
- 279. W[ALSH], B. D. Hop-growing in the West. < Pract. Ent., May, 1867, v. 2, pp. 93-94.
 - Occurrence in Michigan of *Phorodon humuli*; its destructiveness; popular ignorance of entomology.
- 280. W[ALSH], B. D. The wheat midge—jumping to a conclusion. <Pract. Ent., May, 1867, v. 2, p. 94.
 - Extract, with comment, from Maryland Farmer and Mcchanic, August, 1865; Diplosis tritici not infested with parasites in America; food-habits of Thrips.

- 281. W[ALSH], B. D. Tree cut-worms. < Pract. Eut., May, 1867, v. 2, p. 94.
 - "Dark-sided cut-worm" an Agrotis and not Hadena chenopodii [= Mamestra trifolii]. See Nos. 85, 229.
- 282. W[ALSH], B. D. Quacks and physicians. < Pract. Ent., May, 1867, v. 2, p. 95.
 - Criticism of proposed universal remedy against insects injurious to the rose (Western Rural, 30 March, 1867).
- 283. W[ALSH], B. D. The ephemeron or May-fly. < Pract. Ent., May, 1867, v. 2, p. 95.

Length of duration of life of Ephemeridæ.

- 284. W[ALSH], B. D. A groundless fear. < Pract. Ent., May, 1867, v. 2, p. 95.
 - Extract from Mo. Rept. Bureau Agric., February, 1867, p. 62; food-habits of larva and imago of Lytta atrata [= Epicauta pennsylvanica]; places of oviposition.
- 285. W[ALSH], B. D. Fire-blight. < Pract. Ent., May, 1867, v. 2, p. 96.

Probable cause.

- 286. [Walsh, B. D.] [Insect eggs.] < Pract. Ent., May, 1867, v. 2, p. 96.
 - Answer to inquiry of W. Willock; identification of eggs of *Œcanthus niveus* and of one of the *Membracidw* on grape twigs.
- 287. [WALSH, B. D.] [Attacus cecropia.] < Pract. Ent., May, 1867, v. 2, p. 96.

Answer to inquiry of J. B. H.; characters of Attacus cecropia.

- 288. [Walsh, B. D.] [Twig pruners.] < Pract. Ent., May, 1867, v. 2, p. 96.
 - Answer to inquiry of M. W. Philips; twigs of pear and other trees amproated in the same manner as oak twigs by *Elaphidion putator* [=villosum].
- 289. [Walsh, B. D.] [Blister-beetles.] < Pract. Ent., May, 1867, v. 2, p. 97.
 - Answer to inquiry of J. M. Tracy; food-habits, ravages, and means against $Lytta \ [= Pomphop \alpha a] \ anea;$ food-plants of $L. \ [= P.] \ tarsalis;$ tabular separation of three species of $Lytta \ [= Pomphop \alpha a]$.
- 290. [Walsh, B. D.] [Means against Curculio: *Œstridæ*.] < Pract. Ent., May, 1867, v. 2, p. 97.
 - Answer to inquiry of C. Greene; ineffective means against Conotrachelus neurphar; Estridæ infesting the ox and rabbit.
- 291. [WALSH, B. D.] [Insect éggs.] < Pract. Ent., May, 1867, v. 2, p. 97.
 - Answer to inquiry of P. Ferris; identification of eggs of Clisiocampa sylvatica [=disstria] and Aphis mali.

292. W[ALSH], B. D. The wheat midge. <Pract. Ent., June, 1867, v. 2, pp. 99-101.

Extract from letter from S. S. Rathvon, with remarks; natural history of Diplosis tritici.

293. W[ALSH], B. D. The new or Colorado potato-bug. <Pract. Ent., June, 1867, v. 2, pp. 101-102.

Distribution, habits, and extent of injuries of Doryphora 10-lineata.

294. [Walsh, B. D.] [Gall-flies; saw-flies.] < Pract. Ent., June, 1867, v. 2, pp. 102–103.

Answer to inquiry of M. W. Philips; mention of various insects which are gall-makers; their modes of life; characters of the larvæ of Cecidomyidæ and Tenthredinidæ.

- 295. [Walsh, B. D.] [Bark-beetles.] < Pract. Ent., June, 1867, v. 2, p. 103.
 - Answer to inquiry of W. C. Fish; identification of specimens sent; habits of Tomicus pusillus [= Pityophthorus minutissimus] and of T. ramulorum [= P. micrographus].
- 296. [WALSH, B. D.] [Apple-tree insects.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of S. Cassi; characters, food-plants, and ravages of Arctia [= Spilosoma] virginica; supposed work of Epicarus imbricatus.

297. [Walsh, B. D.] [Attacus cecropia.] < Praet. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of T. McGraw; larvæ of Attacus cecropia usually solitary.

298. [Walsh, B. D.] [Grape-vine flea-beetle.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of A. Kelley; hibernation of the image of Haltica chalybea; ravages and means against the same.

299. [WALSH, B. D.] [Sphyracephala brevicornis.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of H. B. Howarth; characters and habitat of Sphyrace-phala brevicornis.

300. [Walsh, B. D.] [May-bug.] <Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of C. H. R.; characters and food-habits of the larva of Lachnosterna quercina [== fusca].

301. [Walsh, B. D.] [Blister-beetles: plant-louse enemy.] < Pract. Ent., June, 1867, v. 2, p. 104.

Answer to inquiry of Dr. Houghton; food-habits of $Lytta = Pomphop \alpha a$ and L = P. sayi; Chilocorus bivulnerus as a means against Coccide.

302. [Walsh, B. D.] [Blackberry borers.] < Pract. Ent., June, 1867, v. 2, p. 104.

Answer to inquiry of A. S. Fuller; characters of undetermined larva (*Æge-ria?*) and of *Oberea perspicillata* [= bimaculata] boring in blackberry stems; food-habits of *Ægeridw*.

- 303. W[ALSH], B. D. Scientific nomenclature. <Pract. Ent., July, 1867, v. 2, pp. 105-106.
 - Criticism of the practice of establishing new genera upon slight characters, as illustrated by the American silk-worms.
- 304. W[ALSH], B. D. Currant plant-lice (Aphis ribis). < Pract. Ent., July, 1867, v. 2, p. 106.
 - Method of work of Aphis [=Myzus] ribis.
- 305. [Walsh, B. D.] [Lucanus elaphus.] < Pract. Ent., July, 1867, v. 2, p. 107.
 - Answer to inquiry of S. Hayeraft; characters and habits of *Lucanus elaphus*; rarity of the Q; use of prolonged jaws in Z insects.
- 306. [Walsh, B. D.] [New York weevil.] < Pract. Ent., July, 1867, v. 2, p. 107.
 - Answer to inquiry of T. Gregg; habits and distribution of Ithycerus noveboracensis.
- 307. [Walsh, B. D.] [Codling-moth.] < Pract. Ent., July, 1867, v. 2, p. 107.
 - Answer to inquiry of H. Morey; number of broods, habits, etc., of Carpocapsa pomonella.
- 308. [Walsh, B. D.] [Hickory borer, etc.] < Pract. Ent., July, 1867, v. 2, p. 107.
 - Answer to inquiry of T. Wiggins; times of cmergenee of Clytus [= Cyllene] pictus and C. [= C.] robiniæ; habits of Corydalus cornutus.
- 309. [Walsh, B. D.] [Wasps.] < Pract. Ent., July, 1867, v. 2, p. 107.

 Answer to inquiry of G. W. Smith; food-habits of digger-wasps.
- 310. [Walsh, B. D.] [Parasitic flies, etc.] < Pract. Ent., July, 1867, v. 2, pp. 107–108.
 - Answer to inquiry of S. P. Monks; method of transformation of parasitie Hymenoptera; use of Latin plurals for English words.
- 311. [Walsh, B. D.] [Workers among Hymenoptera.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of W. V. Andrews; economy of workers among the social Hymenoptera.
- 312. [Walsh, B. D.] [Cut-worms.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of V. Abbey; means against Agrotis sp.?
- 313. [Walsh, B. D.] [Potato-beetle enemies.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of W. Smith; scutellerid enemy of larva of *Doryphora* 10-lineata; katydids as an enemy of the same.
- 314. [Walsh, B. D.] [Black-bug.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of J. R. Tewksbury; characters and habits of *Pirates* [= Melanolestes] picipes.

- 315. [Walsh, B. D.] [Burying-beetles.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of A. D. Strong; characters of Necrophorus marginatus; habits of Silphidæ; mite parasites of Silphidæ and other beetles.
- 316. [Walsh, B. D.] [Insects named.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of M. S. Hill; identification of specimens sent; food-plants of larva of Chrysomela [= Gastroidea] cyanea.
- 317. [Walsh, B. D.] [Spring-tails.] < Pract. Ent., July, 1867, v. 2, pp. 108-109.
 - Answer to inquiry of E. Daggy; characters, habits, and means against Poduridæ.
- 318. [Walsh, B. D.] [Blackberry gall insects.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of Aculeus; number of abdominal segments of *Diastrophus* and *Aulax*; reasons for considering the former the gall-maker and the latter the guest-fly.
- 319. [Walsh, B. D.] [Pine and cedar longicorns.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of J. Barratt; sexual characters and food-habits of Callidium antennatum and C. janthinium.
- 320. [Walsh, B. D.] [White-pine plant-louse, etc.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of C. H. Peck; Lachnus strobi on Pinus strobus; undetermined bark-louse, with enormous lateral thoracic plates, found on Rhus glabra.
- 321. [Walsh, B. D.] [Cut-worms.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of J. Townley; food-habits of climbing cut-worms.
- 322. [Walsh, B. D.] [Currant plant-louse.] < Pract. Ent., July, 1867, v. 2, pp. 109, 110.
 - Answer to inquiry of R. L. Walker; Coccinellidæ as a means against Aphis [= Myzus] ribis.
- 323. [Walsh, B. D.] [Beetles.] < Pract. Ent., July, 1867, v. 2, p. 110.

 Answer to inquiry of I. A. Plucke; characters and food-habits of Tetraopes tornator [= tetraophthalmus] and of T. femoratus; comparative characters of Chrysomela bigsbyana and of C. philadelphica; characters of Chrysomelidæ and Coccinellidæ.
- 324. W[ALSH], B. D. Fertilizing plants. < Pract. Ert., July, 1867, v. 2, p. 110.
 - Part played by insects in fertilizing plants.
- 325. W[ALSH], B. D. The peach-worm. <Pract. Ent., July, 1867, v. 2, p. 110.
 - Larvæ of Ephestia zew [=interpunctella] injurious to dried peaches; foodplants and encmies of the same.

326. W[ALSH], B. D. Valedictory. <Pract. Ent., August-September, 1867, v. 2, p. 111.

Note of thanks for aid received in editing the Practical Entomologist.

- 327. W[ALSH], B. D. The State entomologist of Illinois. < Pract. Ent., August-September, 1867, v. 2, p. 111. Reprint: < Can. Farmer, 1 October, 1867, v. 4. S.-b., No. 2, p. 98.
 - Bill for appointment of State entomologist of Illinois not carried into effect.
- 328. W[ALSH], B. D. The three so-called army-worms. < Pract. Ent., August-September, 1867, v. 2, pp. 111-114.
 - Confusion caused by the use of the same popular name for Anomis [= Aletia] xylina, Leucania unipuncta, and Clisiocampa sylvatica [= disstria]; habits, etc., of the three species; food-habits of their larvæ; characters, parasites, and means against C. sylvatica [= disstria].
- 329. W[ALSH], B. D. The little Turk and its crescent. < Pract. Ent., August-September, 1867, v. 2, pp. 114-115.

 Comments on letter from F. C. Hill; oviposition of Conotrachelus nenuphar.
- 330. [Walsh, B. D.] Apple-worm. (Carpocapsa pomonella.) < Pract. Ent., August-September, 1867, v. 2, p. 115.

Prevalence of C. pomonella in 1867; food-habits of its larva.

331. W[ALSH], B. D. Spindle-worms. < Pract. Ent., August-September, 1867, v. 2, pp. 115-116.

Habits, etc., of Achatodes zew and of Gortyna nitela.

- 332. W[ALSH], B. D. A plant growing out of an insect. < Pract. Ent., August-September, 1867, v. 2, p. 116.
 - Larvæ of Lachnosterna fusca infested with Cordyceps turned up by the plow in large numbers; larvæ supposed to have caten poisonous seed which has germinated after killing the larvæ; sowing the seed as a means against the larvæ.
- 333. [Walsh, B. D.] The imported gooseberry saw-fly. < Pract. Ent., August-September, 1867, v. 2, p. 116.
 - Changes in the larva of Nematus ventricosus [= ribcsii] during the last moult; ravages in Columbia County, N. Y.; extent of spread from Rochester, N. Y.
- 334. W[ALSH], B. D. The Colorado potato-bug. <Pract. Ent., August-September, 1867, v. 2, p. 116.
 - Spread of Doryphora 10-lineata into southern Michigan and western Indiana.
- 335. W[ALSH], B. D. Apple-tree plant-lice. (Aphis mali.) < Pract. Ent., August-September, 1867, v. 2, pp. 116-117.

 Effects of early frosts on Aphis mali.
- 336. W[Alsh], B. D. The tent-caterpillar of the apple-tree. (*Clisio-campa americana*.) < Pract. Ent., August-September, 1867, v. 2, p. 117.

Scarcity of C. americana in 1867 due to the abundance of an egg parasite.

- 337. [Walsh, B. D.] A new foe of the corn. < Pract. Ent., August-September, 1867, v. 2, pp. 117-118.
 - Ravages and description of Sphenophorus zew n. sp. [= sculptilis]; habits of the larva and imago.
- 338. [Walsh, B. D.] [*Rhyssa atrata*.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of H. K. Smith; Rhyssa [= Thalessa] atrata ovipositing in hickory.
- 339. [Walsh, B. D.] [Grape-vine beetles.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of L. D. Morse; ravages and food-plants of *Fidia viticida* and of *F. longipes*.
- 340. [Walsh, B. D.] [Currant insects.] <Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of A. M. Burns; identification of *Egeria tipuliformis*; mention of species called currant-worms; destructiveness of *Nematus ventricosus* [=ribesii].
- 341. [Walsh, B. D.] [Rascal leaf-crumpler.] < Pract. Ent., August–September, 1867, v. 2, p. 118.
 - Answer to inquiry of J. M. K.; times of transformation and means against Phycita nebulo [= Acrobasis indiginella].
- 342. [Walsh, B. D.] [Strawberry enemy.] '<Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of C. Faxon; characters, habits, and means against *Polydesmus serratus*; roots of strawberry plants injured by the same.
- 343. [Walsh, B. D.] [Eight-spotted forester.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of A. Gilbert; characters and food-plants of Alypia octomaculata.
- 344. [WALSH, B. D.] [Maple bark-louse.] < Pract. Eut., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of B. W. McLain; characters of undescribed coccid on maple.
- 345. [Walsh, B. D.] [Apple-tree bark-lice.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of A. Gilbert; identification of Aspidiotus harrisii [= Chionaspis furfurus] and of A. conchiformis [= Mytilaspis pomorum] on appletrees.
- 346. [Walsh, B. D.] [Habits of larvæ and pupæ.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of M. M. S.; insects which transform under ground enter the earth in the larval stage; apparatus used by pupæ in forcing their way to the surface of the ground.
- 347. [Walsh, B. D.] [Potato-beetles: Cicindela sex-guttata.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of P. B. Sibley; mention of Lytta cinerea [=Macrobasis unicolor], and of Doryphora 10-lineata as destructive to potato vines; supposed larval habits of Cicindela sex-guttata.

- 348. [Walsh, B. D.] [Wavy-striped flea-beetle.] < Pract., Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of E. T. Snelling; food-habits of Haltica striolata [= Phyllotreta vittata].
- 349. [Walsh, B. D.] [Parasites of cut-worms.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of J. Edgerton; metamorphoses of tachinid parasites of Noctuida.
- 350. [Walsh, B. D.] [Eggs on sugar-maple.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of W. Priehard; egg-rings of Clisiocampa americana? on sugar-maple.
- 351. [Walsh, B. D.] [Click-beetles.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of J. H. Foster, jr.; food-habits of Melanotus communis and of Monocrepidius vespertinus.
- 352. [WALSH, B. D.] [Chrysobothris femorata.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of J. J. Kelly; food-habits of Chrysobothris femorata.
- 353. [Walsh, B. D.] [Gall on chestnut.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of W. Kite; mention of an undetermined *Phylloxera* gall on the flower-catkin of the chestnut.
- 354. [Walsh, B. D.] [Grape-vine beetle: Bugs.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of B. F. Lazear; habits of *Pelidnota punctata*; swarming of *Corimelana pulicaria*; defensive odors of Heteroptera.
- 355. [Walsh, B. D.] [Squash-vine insects.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of J. H. Parsons; larval habits of *Diabrotica vittata*; confusion eaused by popular names.
- 356. [Walsh, B. D.] [Mites.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of C. Carpenter; characters and life-habits of Acarina.
- 357. [Walsh, B. D.] [Haltica cucumeris.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of G. S.; food-habits of Haltica [= Crepidodera] cucumeris.
- 358. [Walsh, B. D.] [Insects named.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of M. H. Boye; food-habits of Conotrachelus cratægi, Melanotus communis, Pelidnota punctata, Desmocerus palliatus, Lucidota atra, Podabrus rugulosus, and P. modestus; variation of Haltica [== Crepidodera] helxines; characters of Dolerus arvensis.

- 359. [Walsh, B. D.] [Insects named.] <Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of H. Burt; identification of several insects; climbing habits of cut-worms.
- 360. [Walsh, B. D.] [Hickory-galls.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of W. C. Fish; characters of galls made by *Phylloxera* caryæglobuli and *P. caryæcaulis*; identification of several beetles.
- 361. [Walsh, B. D.] [Yellow swallow-tail; Rose-beetle.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of J. Barratt; identification of Papilio turnus; transformation and ravages of Macrodactylus subspinosus.
- 362. [Walsh, B. D.] [Blister-beetles.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of D. W. Kauffman; food-plants and characters of *Macrobasis murina* [= unicolor] and of *Epicauta atrata* [= pennsylvanica].
- 363. [Walsh, B. D.] [Tent-caterpillars.] < Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Answer to inquiry of S. R. Williams; occurrence of Clisiocampa americana on pear-trees and of C. sylvatica [== distria] on black-walnut.
- 364. [Walsh, B. D.] [Currant-worms.] < Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Answer to inquiry of A. H. Mills; characters of the larva of Nematus ventricosus [=ribesii], of Ellopia [= Eufitchia] ribearia, and of Pristiphora grossularia.
- 365. [Walsh, B. D.] [Potato-beetle enemies.] < Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Answer to inquiry of W. Maus; description and habits of Lebia grandis; hemipterous enemics of Doryphora 10-lineata.
- 366. W[ALSH], B. D. The Colorado potato-bug. <Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Extract from Mo. Rept. U. S. Agric. Bureau, 1866, p. 344; doubts the occurrence of *Doryphora* 10-lineata in Pennsylvania.
- 367. [Walsh, B. D.] Entomology. < Prairie Farmer, December, 1867, [v. 36], n. s., v. 20, pp. 358-359.
 - Address, with discussion by others, at the third annual meeting of the Southern Illinois Fruit Growers and Shippers' Association; ravages and means against Pemphigus pyri [= Schizoncura lanigera], Carpocapsa pomonella, Trypeta pomonella, Semasia [= Grapholitha] pruinivora, Caliodes [= Craponius] inaqualis, Conotrachelus nenuphar.
- **368.** Walsh, B. D. The grasshoppers. <Iowa Homestead, 15 January, 1868 (v. 13?), p. 9. S.-b., No. 2, p. 119.
 - Range of swarms of Caloptonus spretus in Iowa in 1867.
- 369. Walsh, B. D. Birds vs. insects. < Prairie Farmer, 1868, [v. 37], n. s., v. 21; 30 May, pp. 346-347; 6 June, pp. 362-363; 13 June, pp. 378, 379.

- 370. Walsh, B. D. The seventeen-year locust. <Dixie Farmer, 11 June, 1868. S.-b., No. 3, p. 23.
 - Periodicity and local distribution of the various broods of Cicada [= Tibicen] septendecim.
- 371. Walsh, B. D. Entomological. Do locusts sting? < Chicago Republican, 1868. S.-b., No. 3, p. 25.
 - Sting of Cicada not poisonous; sting of Stizus grandis [== Sphecius speciosus] mistaken for it; habits of the Stizus.
- 372. Walsh, B. D. The bug-hunter in Egypt. A journal of an entomological tour into south Illinois by the senior editor. < Amer. Ent., 1868, v. 1; September, pp. 6-14, figs. 1-7; October, pp. 29-36, figs. 23-29.
 - Habits and figures of Macronema zebratum, Palingenia [= Hexagenia] bilineata, Cicada sp., Stizus grandis [= Sphecius speciosus], Xylocopa carolina [= virginica, and Anthophora sponsa [=abrupta]; sting of Cicada not poisonous, that of the Stizus mistaken for it; habits of Conotrachelus nanuphar; description of Hull's Curculio-catcher; Conotrachelus anaglypticus and Colaspis sp. injurious to plum-trees; figures of Rhopalus sp. and Euryomia [= Euphoria | melancholica injurious to pear-trees; figure and habits of Metapodius nasalus [= fcmoratus]; predaceous habits of Arma [= Podisus] spinosus, Evagoras viridis [=Diylodus luridus], and Stiretrus fimbriatus [=anchorago]; work of Aspidiotus harrisii [= Chionaspis furfurus] upon pear, and of Cecidomyia [= Diplosis] tritici in wheat; occurrence of Endrosa [= Lachnosterna] quercus, Doryphorajuneta, and Trichius delta in Illinois; mite enemy of Aspidiotus conchiformis [= Mytilaspis pomorum]; injury to fruit by Apis mellifica; hot water as a means against Schizoneura lanigera; habits and figures of Chrysopa sp.; figures of Aspidoglossa subangulata, Harpalus pennsylvanicus, and Evarthrus orbatus; figures and descriptions of larva of a carabid and of Chauliognathus pennsylvanicus; Curculio-feeding habits of Carabidæ.
- 373. Walsh, B. D. First annual report on the noxious insects of the State of Illinois. <Trans. Ill. State Hortic. Soc. for 1867, n. s., v. 1, Appendix. Separate: <Chicago, 1868, 103 pp., figs. 1-3.

TABLE OF CONTENTS.

INTRODUCTORY	3
CHAPTER 1. The grape Curculio. Caliodes [= Craponius] inaqualis Say.	
Its history—Guest insects, parasites, and cannibals—Remedy	13
CHAPTER 2. The grape-leaf gall-louse. Dactylosphara vitifolia, Fitch	
$[=Phylloxera\ vastatrix].$	
Peculiar to the Clinton and other cultivated varieties of Frost	
grape—Practical inference therefrom	21
Chapter 3. The rose-bug. Macrodactylus subspinosus Linn.	
Device for destroying it on grape-vines	24
CHAPTER 4. The grape-root borer. Egeria [= Sciapteron] polistiformis	
Harris.	
Its history—Its scientific nomenclature—Remedies	24
CHAPTER 5. The apple-worm or codling-moth. Carpocapsa pomonella	
Linn.	
Double-brooded—Practical inference therefrom—Remedies	27
Chapter 6. The apple-magget fly. Trypeta pomonella Walsh.	
Occurs at present only in certain Eastern States on the apple,	

though it was found long ago in Illinois on the crab-Its history.

29

079	WATER P. D. Continued	
010.	WALSH, B. D.—Continued. CHAPTER 7. The rascal leaf-crumpler. Phycita nebulo Walsh [= Acro-	
	basis indiginella].	
	A peculiarly Northwestern species, not found either East or South—	
	Easily destroyed	34
	Chapter 8. The oyster-shell bark-louse. Aspidiotus conchiformis Gmel. $[=Mytilaspis\ pomorum].$	
	Its history—Loses almost all its organs when only a few days old,	
	and becomes thenceforth as stationary as a cabbage—How it spreads from tree to tree—Mites and their natural history—Plant-	
	feeding mites, parasitic mites, and cannibal mites—The bark-	
	louse largely preyed on by a minute cannibal mite—Uscless and	
	useful remedies	34
hape	Chapter 9. Harris' bark-louse. Aspidiotus harrisii Walsh [=Chionaspis furfurus].	
	How it differs from the preceding—The geographical distribution.	53
	CHAPTER 10. The apple-root plant-louse. Pemphigus pyri Fitch [= Schizoneura lauigera].	*
	Often confounded in Illinois with the true "Woolly plaut-louse"	
	The differences—Its history—Causes a form of "rotten-root"—	
	Its supposed cannibal foes—Remedies.	55
	CHAPTER 11. The plum Curculio. Conotrachelus nenuphar Herbst. Its history—Its peculiar crescent cut explained—Double-brooded—	
	Its supposed enemies, the so-called "Curculio parasite" and the	
	Baltimore oriole—Remedies	64
	Chapter 12. The plum-gouger. Anthonomus prunicida Walsh [= Coccotorus scutellaris].	
	Bores a round hole in the plum, instead of a crescent cut—How	
	and why it does this-Differs in many other respects from the	
	Curculio	72
	CHAPTER 13. The plum-moth. Semasia [=Grapholitha] prunivora Walsh.	
	Its history—Probably a guest-moth, and therefore not injurious	78
	CHAPTER 14. The hateful grasshopper. Caloptenus spretus Walsh.	
	The Rocky Mountains its natural home—Invades in certain years certain neighboring districts, such as Texas, Missouri, Kansas,	۰
	Nebraska, and Iowa—Lays there millions of eggs, which develop	
	into barren grasshoppers only—Invasion of 1866—Damage done	
	by the young grasshoppers in the spring of 1867—In Kausas	
	about one-eighth of the field crops and seven-eighths of the gar- den crops destroyed by them—Grashopper invasious of A. D.	
	1820, 1856, 1857, 1864, and 1867—Probable results in 1868 of	
	the invasion of 1867—Has never yet come within 115 miles of	
	Illinois—Can not spread into Illinois as the Colorado potato-bug	
	has done—Reasons why, in all human probability, it can never	20
	reach Illinois at all	82

374. Walsh, B. D. A friend unmasked. < Amer. Ent., November, 1868, v. 1, pp. 51-52, fig. 55.

Description and figures of larva and adult of Chauliognathus pennsylvanicus; characters and habits of the Telephorida.

- 375. Walsh, B. D. Appendix to the article on "Wasps and their habits," in No. 7. < Amer. Ent., April, 1869, v. 1, pp. 162-164.
 - Habits of Chrysididæ; descriptions of Stizus [Megastizus] brevipennis n. sp., Agenia subcorticalis n. sp., A. architectus Say, A. cupida Cress., A. bombycina Cress., Ceropales rufiventris n. sp., and Ammophila pictipennis n. sp.; tabular separation of the genera of Sphecidæ. See No. 543.
- 376. Walsh, B. D. Mr. Couper's thorn-leaf gall. <Ca. Ent., 15 April, 1869, v. 1, pp. 79-80.
 - Description of cecidomyidous leaf-gall on *Cratægus*; mentions of three additional Cecidomyidous, and an Acaridous gall on the same; *Anthonomus cratægi* inquilinous in gall on *Cratægus*.
- 377. Walsh, B. D. The six worst insect enemies of the fruit growers in northern Illinois. <Trans. North. Ill. Hortic. Soc. for 1867-768, pp. 91-96.
 - Treats of Carpocapsa pomonella, Aspidiotus conchiformis [=Mytilaspis pomorum], Anisopteryx [= Paleacrita] vernata, Saperda bivittata [= candida], Chrysobothris femorata, Conotrachelus nenuphar, and Anthonomus prunicida [= Coccotorus scutellaris].
- 378. Walsh, B. D. Squash borer. < Cultivator and Country Gentleman, 30 September, 1869, v. 34, p. 256. Extract: < Prairie Farmer, 30 October, 1869.
 - Characters of larva and image of *Egeria cucurbitæ* [= *Melittia ceto*]; natural history, means against, and distribution of the same.
- 379. Walsh, B. D. Apple-tree borers. <Journ. Ill. State Agric. Soc. Reprint: <Trans. Ill. State Agric. Soc., v. 5, pp. 499-501.
 - Comparative differences between Saperda and Chrysobothris; soap as a means against the same.
- 380. Walsh, B. D. On a species of *Hemiteles* (*Ichneumonidæ*), ascertained by the editor to be parasitic in Canada on the imported current-worm fly (*Nematus ventricosus* Klug). <Can. Ent., 1 October, 1869, v. 2, pp. 9-12.
 - Division of *Hemiteles* into two groups; description of *H. nemativorus* n. sp.; larval and pupal history of the same by C. J. S. Bethune.
- 381. Walsh, B. D. The imported currant-worm fly (Nematus ventricosus Klug) and its parasite (Hemiteles nemativorus Walsh) <Ca. Ent., 15 November, 1869, v. 2, pp. 31–33.
 - Correction of errors in No. 380; sexual characters of *Hemiteles*; variation in the time at which the image state is assumed by hibernating insects; heteropterous enemies of *Nematus ventricosus* [= ribesii].
- 382. Walsh, B. D. Larvæ in the human bowels. One of Mr. Walsh's posthumous articles. <Amer. Ent., March, 1870, v. 2, pp. 137–141, fig. 93.
 - Characters and habits of larval *Homalomyia*; description of the larva of *H. wilsoni*, *H. leidyi*, and *H. prunivora* n. spp.; mention of recorded cases of diptera in the human bowels; figures larva of *H. wilsoni*.

383. Walsh, B. D. One day's journal of a State entomologist. < Amer. Ent. and Bot., May, 1870, v. 2, pp. 197-199.

Sketch of an average day's work of a State entomologist.


384. Walsh, B. D. On the group Eurytomides of the hymenopterous family Chalcididæ; with remarks on the theory of species and a description of Antigaster, a new and very anomalous genus of Chalcididæ. <Amer. Ent. and Bot., 1870, v. 2, September, pp. 297-301, fig. 1, 2; October, pp. 329-335, fig. 3-6; December, pp. 367-370, fig. 7-10.

Descriptions, habits, and figures of many Eurytomides; synoptic tables of the species of Eurytoma and Decatoma. For a list of the new species see the Systematic Index.

385. Walsh, B. D. Descriptions of North American Hymenoptera. Trans.Acad.Sci.St.Louis, 7 May—9 June, 1873, v. 3, pp. 65–166, fig.

Printed posthumously, with notes by E. T. Cresson; description of many previously described *Tenthredinidæ* and *Ichneumonidæ*; five (5) new genera and forty-five (45) new species of *Ichneumonidæ* are described, for a list of which see the *Systematic Index*; figures venation of front wing of *Ichneumona*.

4 ENT



U.S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

 \mathbf{OF}

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

BY

SAMUEL HENSHAW.


PART II.

THE MORE IMPORTANT JOINT WRITINGS

OF

B. D. WALSH and C. V. RILEY.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1889.



BIBLIOGRAPHY OF THE MORE IMPORTANT CONTRIBUTIONS TO AMERICAN ECONOMIC ENTOMOLOGY.

PART II.

- 386. [Walsh, B. D., and C. V. Riley.] Salutatory. To the agriculturists and horticulturists of the United States. <Amer. Ent., September, 1868, v. 1, pp. 1-3.
 - Importance and extent of the ravages of insects; practicability of the diminution of these ravages; value of entomological work; aims of the American Entomologist.
- 387. [Walsh, B. D., and C. V. Riley.] Hogs vs. bugs. < Amer. Ent., September, 1868, v. 1, pp. 3-6.
 - Effectiveness of swine as a means against Conotrachelus nenuphar and Carpocapsa pomonella; quotations showing good results from pasturing swine in fruit orchards.
- 388. [Walsh, B. D., and C. V. Riley.] Insect changes. <Amer. Ent., September, 1868, v. 1, p. 6.

 Definition of the different stages of insect transformation.
- 389. [WALSH, B. D., and C. V. RILEY.] A new bark-louse on the osage orange. <Amer. Ent., September, 1868, v. 1, p. 14, fig. 8. Seasons, food-plants, characters, and figures of Lecanium macluræ n. sp., [=Pulvinaria innumerabilis] and of L. acericola n. sp. [= P. innumerabilis]; characters and economic importance of the genus Aspidiotus.
- 390. [Walsh, B. D., and C. V. Riley.] Entomological ignorance in the South. <Amer. Ent., September, 1868, v. 1, pp. 14-16.

 Reprint and criticism of article, "How to destroy the cotton-worm—a suggestion."
- 391. [Walsh, B. D., and C. V. Riley.] Grasshoppers. <Amer. Ent., September, 1868, v. 1, p. 16.
 - Ravages of Acrididæ in Illinois, Iowa, and Missouri; mention of the species committing these ravages; means against the same.
- 392. [Walsh, B. D., and C. V. Riley.] The old and the new philosophy. <Amer. Ent., September, 1868, v. 1, p. 17.

 Defense of the practical application of science.
- 393. [Walsh, B. D., and C. V. Riley.] The animal kingdom. < Amer. Ent., September, 1868, v. 1, pp. 17-18.
 - Definition of the "four grand branches of the animal kingdom;" mention of their respective classes.

- 394. [Walsh, B. D., and C. V. Riley.] Ad interim committees. <Amer. Ent., September, 1868, v. 1, p. 18.
 - Duties of ad interim committees of horticultural societies; the American Entomologist ready to publish any scientific discoveries or observations with regard to rots or diseases of a vegetal nature.
- 395. [Walsh, B. D., and C. V. Riley.] On our table. < Amer. Ent., September, 1868, v. 1, p. 18.
 - Notice of W. H. Edwards's "The butterflies of North America," pt. 1; of A. S. Packard, jr.'s "Guide to the study of insects," pt. 2; and of the "Canadian Entomologist," v. 1, No. 1.
- 396. [Walsh, B. D., and C. V. Riley.] "Fire-flies." < Amer. Ent., September, 1868, v. 1, p. 19, fig. 9.
 - Answer to inquiry of W. McC.; characters and transformations of *Photinus* pyralis; figures larva, with details of structure, pupa, and imago.
- 397. [Walsh, B. D., and C. V. Riley.] A new grape root-borer. <Amer. Ent., September, 1868, v. 1, p. 19.
 - Answer to inquiry of W. D. F. Lummis; characters and ravages of an undetermined longicorn larva; food-habits of Orthosoma cylindricum [== brunnenm].
- 398. [Walsh, B.D., and C.V. Riley.] Bugs on grape-vines mistaken for chinch-bugs. <Amer. Ent., September, 1868, v. 1, p. 19.

 Answer to inquiry of F. Hecker; characters, ravages, and food-plants of Blissus lencopterus and of Piesma cinerea.
- 399. [Walsh, B. D., and C. V. Riley.] Leaf-hoppers of the grape. <Amer. Ent., September, 1868, v. 1, p. 19.

Answer to inquiry of F. Hecker; characters and ravages of *Proconia* [= Oncometopia] undata.

- 400. [Walsh, B. D., and C. V. Riley.] Leaf-bugs. < Amer. Ent., September, 1868, v. 1, p. 19.
 - Answer to inquiry of C. H. Pcck; food-habits of Tingis = Gargaphia tilia and of T. = Corythuca ciliata; punctures on leaves caused by Hemiptera.
- 401. [Walsh, B. D., and C. V. Riley.] Potato bugs. <Amer. Ent., 1868, v. 1, October, pp. 21–27, fig. 10–19; November, pp. 41–49, fig. 33–48.
 - Failure of popular authors to distinguish between the different insects infesting the potato plant; brief accounts and figures of one or more stages of the following species: Gortyna nitela, Baridius [= Trichobaris] trinotata, Sphinx 5-maculata [= Protoparce celeus], Lytta [= Epicauta] vittata, L. atrata [= E. pennsylvanica], L. marginata [= E. cinerea], L. cinerea [= Macrobasis unicolor], L. murina [= M. unicolor], Lema trilineata, and Haltica [= Crepidodera] cucumeris; migrations, habits, enemies, and means against Doryphora 10-lineata; figures of all stages of D. 10-lineata and D. juncta and of numerous enemics of the former.
- 402. [Walsh, B. D., and C. V. Riley.] "Grasshoppers." Their devastations in western Iowa and the good that has resulted from them. <Amer. Ent., October, 1868, v. 1, pp. 27–28.
 - Letter of M. C. Nickerson, with comments; seeds of Vilfa vaginæflora carried by "grasshoppers."

- 403. [Walsh, B. D., and C. V. Riley.] An apple growing on a grapevine. <Amer. Ent., October, 1868, v. 1, p. 28.
 - Extract from Richmond (Va.) Whig, with criticism; a gall of Cecidomyia vitispomum mistaken for an apple growing on a grape-vine. See Nos. 332, 436, 478.
- 404. [Walsh, B. D., and C. V. Riley.] Ticks and Texas fever. <Amer. Ent., October, 1868, v. 1, p. 28.

 Improbability that *Ixodes bovis* is the cause of the Texas fever of cattle.
- 405. [Walsh, B. D., and C. V. Riley.] Scientific symbols. <Amer. Ent., October, 1868, v. 1, p. 28. Reprint: <Op. cit., November, 1869, v. 2, p. 50.

Explanation of the use of the signs designating the sexes.

- 406. [WALSH, B. D., and C. V. RILEY.] A swarm of butterflies. Amer. Ent., October, 1868, v. 1, pp. 28–29, figs. 20–22.

 Record of swarms of Danais archippus; food-plant of larva; figures larva, chrysalis, and imago.
- 407. [WALSH, B. D., and C. V. RILEY.] The sting of the 17-year Cicada. <Amer. Ent., October, 1868, v. 1, pp. 36-37.

 Communications from F. W. Collins, R. Richardson, and B. Borden, on the reputed sting of Cicada [= Tibicen] septendecim and on the habits of Stizus grandis [= Sphecius speciosus].
- 408. [WALSH, B. D., and C. V. RILEY.] On our table. <Amer. Ent., October, 1868, v. 1, p. 37.

 Notice of "The insect world," by L. Figuier.
- 409. [WALSH, B. D., and C. V. RILEY.] Squash-bug; its change of color. <Amer. Ent., October, 1868, v. 1, p. 37.
 Answer to inquiry of J. Periam; changes of color of Covens [= Anasa] tristis during its metamorphoses.
- 410. [Walsh, B. D., and C. V. Riley.] White grub; immunity from it next year in Clinton County, Mo. <Amer. Ent., October, 1868, v. 1, p. 37.

Answer to inquiry of J. P. McCartney; life-habits of Lachnosterna fusca.

- 411. [Walsh, B. D., and C. V. Riley.] Insect enemies of the Colorado potato beetle. <Amer. Ent., October, 1868, v. 1, p. 37.

 Answer to inquiry of S. H. Kriedelbaugh; identification of three insect enemies of Doryphora 10-lineata.
- 412. [WALSH, B. D., and C. V. RILEY.] "Harvest bugs" in America, misnamed "jiggers." <Amer. Ent., October, 1868, v. 1, p. 38.

 Answer to inquiry of M. McKenzie; habits of parasitic Acarina; distribution of and injuries caused by Sarcopsylla penetrans.
- 413. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., October, 1868, v. 1, p. 38.
 - Answer to inquiry of J. T. Smith; food-plants of Aphis mali and of Notodonta [= Œdcmasia] concinna; Campoplex [= Limneria] fugitiva parasitic on larva of Euchwtes eglc.
- 414. [Walsh, B. D., and C. V. Riley.] Grapes cut off by the tree cricket. <Amer. Ent., October, 1868, v. 1, p. 38, fig. 30-31.

 Answer to inquiry of J. H. Tice; food-habits of *Ecanthus niveus*; deposition

of eggs; figures both sexes.

- 415. [Walsh, B. D., and C. V. Riley.] The Buffalo tree-hopper. <Amer. Ent., October, 1868, v. 1, p. 38.
 - Answer to inquiry of P. Rickert; characters, habits, and means against Ceresa bubalus.
- 416. [Walsh, B. D., and C. V. Riley.] A scavenger mistaken for a foe. <Amer. Ent., October, 1868, v. 1, p. 38.
 - Answer to inquiry of Simmons and Tillson; characters and habits of the larva of Copris carolina.
- 417. [Walsh, B. D., and C. V. Riley.] Insect foes of the bark-louse. <Amer. Ent., October, 1868, v. 1, p. 39, fig. 32.
 - Answer to inquiry of J. Weed; habits and figures of larva and imago of Chilocorus bivulnerus; food-habits and figure of Epilachna borealis.
- 418. [Walsh, B. D., and C. V. Riley.] Best works on entomology. <Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of B. M. Reynolds; mention of several important works on entomology.
- 419. [Walsh, B. D., and C. V. Riley.] Larvæ in plum-gum. < Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of P. Rickert; unknown larva (probably dipterous) found in gum of plum tree; *Mycetophila* sp., bred from gum of peach-tree.
- 420. [Walsh, B. D., and C. V. Riley.] Insect foes of the hop vine. <Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of H. C. Freeman; characters and habits of larva of Saturnia [=Hyperchiria] io and of Vanessa [=Grapta] interrogationis.
- 421. [Walsh, B. D., and C. V. Riley.] Insect foe of the apple-tree borer. <Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of J. E. Walker; undetermined carabid larva preying upon larva of Saperda candida.
- 422. [Walsh, B. D., and C. V. Riley.] Oak tree caterpillars. < Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of W. W. Daniells; characters and habits of larva of *Edema* albifrons.
- 423. [Walsh, B. D., and C. V. Riley.] Bark-lice on apple trees. <Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of J. G. Fleck; Mytilaspis pomorum destroyed by a minute mite.
- 424. [Walsh, B. D., and C. V. Riley.] The "saddle-back" caterpillar. <Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiries of E. Baxter and O. A. Kenyon; characters, habits, and urticating properties of the larva of *Empretia stimulea*.
- 425. [Walsh, B. D., and C. V. Riley.] The regal walnut caterpillar. <a href="#
 - Answer to inquiry of M. Copley; characters and food-plants of larva of Citheronia regalis; times of metamorphosis; characters of the imago.
- 426. [Walsh, B. D., and C. V. Riley.] The Hessian fly. < Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of F. D. Carson; ravages of Cecidomyia destructor; late sowing as a means of avoiding the same.

- 427. [Walsh, B. D., and C. V. Riley.] Red cedar caterpillar. < Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of R. Peter; habits, characters, and means against Thyridopteryx ephemeræformis.
- 428. [Walsh, B. D., and C. V. Riley.] Wheat-midge winter killed. < Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of J. P. Alexander; cause of the scarcity of Diplosis tritici.
- 429. [Walsh, B. D., and C. V. Riley.] Grape vine borer. < Amer. Ent., October, 1868, v. 1, p. 40. Answer to inquiries of A. Barter and J. H. Hogan; undetermined cerambycid

larva injurious to grape-vines.

- 430. [Walsh, B. D., and C. V. Riley.] Popular names and scientific names. < Amer. Ent., November, 1868, v. 1, p. 49, figs. 49-51. Indefiniteness of popular names; figures Julus sp., and the larva and imago of one of the Elaterida.
- 431. [WALSH, B. D., and C. V. RILEY.] Entomological ignorance in the North. < Amer. Ent., November, 1868, v. 1, pp. 50-51, figs. 52-54.
 - Confusion existing in regard to the meaning of the word locust; habits of Acridida and Cicadida; figures types of the two families and of twig with eggs of Cicada sp.
- 432. [WALSH, B. D., and C. V. RILEY.] Tit for tat. < Amer. Ent., November, 1868, v. 1, p. 52. Ridicule of an absurd entomological item.
- 433. [Walsh, B. D., and C. V. Riley.] Grasshoppers. < Amer. Ent., November, 1868, v. 1, p. 53. Abundance and ravages of several species of Acridide and of Acheta [=Gryllus]

abbreviatus in the northern central United States, and scarcity of the same in New York in 1868.

- 434. [Walsh, B. D., and C. V. Riley.] To keep seed peas from bugs. < Amer. Ent., November, 1868, v. 1, p. 53. Habits of and means against Bruchus pisi.
- 435. [Walsh, B. D., and C. V. Riley.] Mind how you pack insects. Amer. Ent., November, 1868, v. 1, p. 54. Care needed to avoid the introduction of noxious insects.
- 436. [Walsh, B. D., and C. V. Riley.] The apple growing on a grape vine. < Amer. Ent., November, 1868, v. 1, p. 54. The "vegetable phenomenon," see Nos. 332, 403, 478, proved to be a gall.
- 437. [WALSH, B. D., and C. V. RILEY.] The late exhibition of useful and destructive insects at Paris. < Amer. Ent., November, 1868, v. 1, p. 55.

Notice of the formation of the Société d'Insectologie Agricole at Paris, and of the second exhibition of the society.

438. [Walsh, B. D., and C. V. Riley.] Destroying black ants in gardens. < Amer. Ent., November, 1868, v. 1, p. 55.

Answer to inquiry of W. S. Patten; means against Formicida in gardens.

- 439. [Walsh, B. D., and C. V. Riley.] Entomological quackery. <Amer. Ent., November, 1868, v. 1, p. 56.
 - Reprint of "The Curculio (Iowa Homestead, 22d July, 1868), with comments; means against Conotrachelus nenuphar.
- 440. [Walsh, B.D., and C.V. Riley.] The cruel bug-hunters. < Amer. Ent., November, 1868, v. 1, p. 56.
 - Inscets are not susceptible of such feelings of pain and pleasure as are felt by higher animals.
- 441. [Walsh, B. D., and C. V. Riley.] Honey bees eating grapes. Amer. Ent., November, 1868, v. 1, p. 56.
 - Reprint of a letter by T. W., from Ohio Farmer, with comments; Apis melli-fica as an enemy of sound fruit.
- 442. [Walsh, B. D., and C. V. Riley.] Twigs amputated by some unknown animal. <Amer. Ent., November, 1868, v. 1, p. 57.

 Answer to inquiry of G. Burnside; method of work of an undetermined fruit-
- tree pruner [= Oncideres cingulata?].

 443. [WALSH, B. D., and C. V. RILEY.] Twigs girdled by some animal. <Amer. Ent., November, 1868, v. 1, p. 57.
 - Answer to inquiries of Judge Brown and P. Earle; method of work of an undetermined fruit-tree pruner [= Oncideres cingulata?]; trees affected.
- 444. [Walsh, B. D., and C. V. Riley.] Hop-vine caterpillar. < Amer. Ent., November, 1868, v. 1, p. 57.
 - Answer to inquiry of H. J. Dunlap; larva of Grapta interrogationis feeding on hop-vines.
- 445. [Walsh, B. D., and C. V. Riley.] "Galls" on leaves of soft maple. Amer. Ent., November, 1868, v. 1, p. 57.
 - Answer to inquiry of A. L. Child; characters of undescribed mite-galls and their architects; mode of formation of leaf-galls by mites.
- 446. [Walsh, B. D., and C. V. Riley.] Unsightly galls on the cottonwood. <Amer. Ent., November, 1868, v. 1, p. 57.
 - Answer to inquiry of A. L. Child; character of galls made by *Pemphigus vagabondus* and *P. populicaulis*; their effects upon the cottonwood and means against them.
- 447. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1868, v. 1, p. 57.
 - Answer to inquiry of E. Baxter; Gordius aquaticus a host in Orchelimum gracile; Cetonia [= Euphoria] inda destructive to the peach.
- 448. [Walsh, B. D., and C. V. Riley.] The stick-bug. Amer. Ent., November, 1868, v. 1, p. 58. Reprint: Can. Farmer, 15 January, 1870.
 - Description, vernacular names, and habits of Spectrum [=Diapheromera] femorata.
- 449. [Walsh, B. D., and C. V. Riley.] Woolly lice on the beech. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of F. H. Guiwits; habits and food-plants of Pemphigus imbricator.

- 450. [Walsh, B. D., and C. V. Riley.] The sheep-bot or head-maggot. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of R. W. Seott; Estrus ovis ordinarily oviparous; sometimes the eggs hatch prematurely inside the body.
- 451. [Walsh, B. D., and C. V. RILEY.] Leaf-miners of the locust. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of R. W. Seott; characters, habits, ravages, and means against *Hispa scutcllaris* [= Odontota dorsalis].
- 452. [Walsh, B. D., and C. V. Riley.] Leaf-hoppers on celery. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of R. Parnell; mention of two undescribed *Tettigonidæ* infesting early eclery.
- 453. [Walsh, B. D., and C. V. Riley.] Parsnip eaterpillars.—Scorpions. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of F. Brewer; food-plants of larva of Papilio asterias; habits of the imago; habits and sting of Buthus carolinianus; its occurrence in Missouri.
- 454. [Walsh, B. D., and C. V. Riley.] Noxious insects named. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of J. H. Parsons; larval characters and food-plants of Hyphantria textor [=cunea], Notodonta [= Edemasia] concinua, Arctia [= Spilosoma] virginica, Carpocapsa pomouella, and of Trypeta pomouella.
- 455. [Walsh, B. D., and C. V. Riley.] Cut-worms destroying recently sown wheat. < Amer. Ent., November, 1868, v. 1, p. 59.

 Answer to inquiry of T. R. Allen; characters, habits, and means against undetermined *Noctuidæ* attacking wheat.
- 456. [Walsh, B. D., and C. V. Riley.] The spined soldier bug. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of I. Hieks; Podisus spinosus as an enemy of Hyphautria textor [= cunea].
- 457. [Walsh, B. D., and C. V. Riley.] The preying Mantis, alias Devil's riding horse, etc. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of A. Pettit; habits of Mantis [=Phasmomantis] carolina; egg-mass of the same.
- 458. [Walsh, B. D., and C. V. Riley.] The pigeon Tremex. < Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of F. Brewer; characters of Tremex columba; food-habits and parasites of its larva.
- 459. [Walsh, B. D., and C. V. Riley.] Flesh-worms. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of W. J. Stuart; larval habits of an undetermined museid.
- 460. [Walsh, B. D., and C. V. Riley.] Fall web-worm on hickory. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of W. W. Daniels; transformation of Hyphantria textor [=cunea]; characters of its larva and imago.

- 461. [Walsh, B. D., and C. V. Riley.] Butterflies named. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of W. W. Butterfield; characters of Colias philodice, Vanessa [= Grapta] comma, and Danais archippus.
- 462. [Walsh, B. D., and C. V. Riley.] Museum pest. < Amer. Ent., November, 1868, v. 1, p. 60.

Answer to inquiry of T. W. Holt, jr.; means against Dermestidæ.

- 463. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. Weed; enemies of plant-lice and bark-lice; foodplant and characters of the larva of *Papilio turnus*.
- 464. [Walsh, B. D., and C. V. Riley.] Grapes spoiled by something. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. Wood; grapes injured from some unknown cause; juice from wounded grapes as food for moths.
- 465. [Walsh, B. D., and C. V. Riley.] White-pine worms. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. T. Little; food-plants of larva of Lophyrus abbotii; transformations of the same.
- 466. [Walsh, B. D., and C. V. Riley.] The Tarantula of Texas. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. Bell; manner in which *Pompilus* [= *Pepsis*] *formosa* prepares *Mygale hentzii* for food for its larva.
- 467. [Walsh, B. D., and C. V. Riley.] Insect foes of the apple-tree. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of G. C. Brackett; characters of larva of Carpocapsa pomonella; means against Datana ministra and Pemphigus pyri [=Schizoneura lanigèra].
- 468. [Walsh, B. D., and C. V. Riley.] Grape-leaf folders eaten by spiders. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of A. A. Hilliard; larva of Desmia maculalis destroyed by a spider.
- 469. [Walsh, B. D., and C. V. Riley.] Caterpillar of the Troilus butterfly. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of A. S. Fuller; food-habits of larva of Papilio troilus.
- 470. [Walsh, B. D., and C. V. Riley.] Potato beetles. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of P. H. Foster; identification of *Epicanta vittata* and *Macrobasis unicolor* as enemics of the potato.
- 471. [Walsh, B. D., and C. V. Riley.] Holes round the roots of young ash trees in the nursery. 'Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of I. Hicks; characters and food-habits of the larva of Xyloryctes satyrus.
- 472. [Walsh, B. D., and C. V. Riley.] Locust borers. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of U. Bruner; destructiveness of the larva of Clytus [= Cyllene] robiniæ.

Lip was well

- 473. [Walsh, B. D., and C. V. Riley.] The hellgrammite fly. <Amer. Ent., December, 1868, v. 1, pp. 61-62, figs. 56-57.
 - Habits of Corydalus cornutus; figures the larva, pupa, and imago; description and figure of the eggs of Belostoma americanum mistaken for those of Corydalus.
- 474. [Walsh, B. D., and C. V. Riley.] The periodical Cicada. < Amer. Ent., December, 1868, v. 1, pp. 63-72, figs. 58-64. Extract: < Op. cit., June, 1869, v. 1, p. 202.
 - Characterization of the 13-year brood of Cicada as a new species, C. = Tibicen tredecim; dimorphism of the same and of C. = T. septendecim; seasons, natural history, transformations, enemies, sting, and injuries of these species; chronological history of their several known broods; figures the several stages of C. = T. septendecim, the towers made by the pupa and twigs with eggs.
- 475. [Walsh, B. D., and C. V. Riley.] The hateful or Colorado grass-hopper, *Caloptenus spretus*, Uhler and Walsh. <Amer. Ent., December, 1868, v. 1, pp. 73-76, fig. 65.
 - Comparative characters and figures of Caloptenus spretus and C. femur-rubrum; native habitat, migrations, distribution, and ravages of the former; harmlessness of its second generation in the Mississippi Valley as compared with the first generation; improbability that it will ever become a permanent resident in that region.
- 476. [Walsh, B. D., and C. V. Riley.] The twig-girdler, *Oncideres cingulatus*, Say. <Amer. Ent., December, 1868, v. 1, pp. 76–77, fig. 66.
 - Oviposition and food habits of Oncideres cingulata; figure of imago and of amputated twig.
- 477. [Walsh, B. D., and C. V. Riley.] An Ichneumon fly mistaken for a wasp. < Amer. Ent., December, 1868, v. 1, p. 77.
 - Review of paper in Christian Advocate on "The stiletto wasp"; Pimpla [= Thalessa] atrata probably mistaken for a wasp.
- 478. [Walsh, B. D., and C. V. Riley.] A plant growing out of an insect. <Amer. Ent., December, 1868, v. 1, p. 77.
 - See Nos. 332, 403, 436; extract from Sedalia Press, with comments; larvæ of Lachnosterna fusca infested with Cordyceps; larvæ supposed to have eaten poisonous seed, which has germinated after killing the larvæ; sowing the seed as a means against the larvæ.
- 479. [Walsh, B. D., and C. V. Riley.] On our table. <Amer. Ent., December, 1868, v. 1, p. 78.
 - Notices of The butterflies of North America by W. H. Edwards.—The Canadian Entomologist.—Guide to the study of insects by A. S. Packard, jr.
- 480. [Walsh, B. D., and C. V. Riley.] Eggs of the white-marked tussock moth. Amer. Ent., December, 1868, v. 1, p. 79, fig. 67.
 - Answer to inquiry of J. M. Hannah; characters of egg and larva of Orgyia leucostigma; sexual differences; parasites; figure of the larva.
- 481..[Walsh, B. D., and C. V. Riley.] Current borers. < Amer. Ent., December, 1868, v. 1, p. 79.
 - Answer to inquiry of B. N. McKinstry; means against Trochilium [= Egeria] tipuliformis, T. [= Alcatha] caudatum, and Psenocerus supernotatus.

- 482. [Walsh, B. D., and C. V. Riley.] Insect infesting grape seed. <Amer. Ent., December, 1868, v. 1, p. 79, fig. 68.
 - Answer to inquiry of A. S. Fuller; characters and figure of larva of *Isosoma* vitis infesting grape seed.
- 483. [Walsh, B. D., and C. V. Riley.] Museum pests again. < Amer. Ent., December, 1868, v. 1, p. 79.
 - Answer to inquiry of J. Huggins; means against Anthreni in collections of natural history.
- 484. [Walsh, B. D., and C. V. Riley.] Grape-vine leaf-hoppers. <Amer. Ent., December, 1868, v. 1, p. 79.
 - Answer to inquiry of P. C. Holmes; means against grape-vine leaf-hoppers.
- 485. [Walsh, B. D., and C. V. Riley.] Apple-tree borer. Amer. Ent., December, 1868, v. 1, p. 80, fig. 69.

 Answer to inquiries of J. T. Zimmerman, C. H. Roberts, and C. R. Babbitt; habits, ravages, and figure of Bostrichus [= Amphicerus] bicaudatus.
- 486. [Walsh, B. D., and C. V. Riley.] The murky ground beetle. <
 - Answer to inquiry of E. T. Dale; fcod-habits and bombardier discharges of Harpalus caliginosus.
- 487. [Walsh, B. D., and C. V. Riley.] Curculio and bark-lice. Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of D. A. Compton; hibernation of Conotrachelus nenuphar; soft soap as a means against Coccidæ.
- 488. [Walsh, B. D., and C. V. Riley.] Insects to be named. <Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of E. B. Beach; identification of several lepidopterous larvæ; food-habits of *Dryocampa stigma* and *D. pellucida* [= virginieusis].
- 489. [Walsh, B. D., and C. V. Riley.] Young pecan trees girdled. Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of Mann and Redmond; means against Oucideres cingulata.
- 490. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of E. T. Dale; identification of several Colcoptera; supposed food-habits of *Tragidion fulvipenne*; character of cocoons of two species of *Microgaster* infesting *Protoparce celens*.
- 491. [Walsh, B. D., and C. V. Riley.] Eggs of true bugs. < Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of S. C. Thornton; characters of eggs of *Prionidus cristatus*. See Amer. Ent., v. 1, pp. 96 and 187.
- 492. [Walsh, B. D., and C. V. Riley.] Museum pest once more. <Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of G. M. L.; larvæ of Anthrenus musworum injuring whalebone, woolen goods, etc.
- 493. [Walsh, B. D., and C. V. Riley.] The Tarantula of Texas again. <Amer. Ent., December, 1868, v. 1, p. 80.

Answer to inquiry of C. Peabody; distribution of Mygale heutzii in Missouri.

- 494. [Walsh, B. D., and C. V. Riley.] The Hessian fly on seed wheat. < Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of U. Scott; broods and times of transformation of Cecidomyia destructor.
- 495. [Walsh, B. D., and C. V. Riley.] The apple-root plant-louse. Eriosoma (Pemphigus) pyri, Fitch. <Amer. Ent., January, 1869, v. 1, pp. 81–84, figs. 70–72.
 - Habits, ravages, description, enemies, and parasites of, and means against Eriosoma pyri [=Schizoneura lanigera]; figures injured root, larva, and adult with details of structure; figures adult plant-louse found on cottonwood; description and figure of larva, puparium, and image of Pipiza radicum n. sp.
- 496. [Walsh, B. D., and C. V. Riley.] Ants' nests in gardens. <Amer. Ent., January, 1869, v. 1, p. 84.

 Means against ants.
- 497. [Walsh, B. D., and C. V. Riley.] The parasites of the human animal. <Amer. Ent., January, 1869, v. 1, pp. 84-88, figs. 73-74.
 - Brief accounts of Pediculus humanus [= vestimenti], P. cervicalis [= capilis], P. [= Pthirius] pubis, Estrus [= Dermatobia] hominis, Pulex irritans, P. [= Sarcopsylla] penetrans, Acanthia lectularia, Conorhinus sanguisuga, and Acarus [= Sarcoptes] scabiei; figures and habits of Reduvius [= Opsicatus] personatus and of Pirates [= Rasahus] biguttatus; figure of Conorhinus sanguisuga; classificatory relations of Pediculina and Mallophaga.
- 498. [Walsh, B. D., and C. V. Riley.] The coffee borer. <Amer. Ent., January, 1869, v. 1, p. 88.
 Unnamed coffee borer injurious to coffee trees in Madras.
- 499. [Walsh, B. D., and C. V. Riley.] Strawberry worms. < Amer. Ent., January, 1869, v. 1, pp. 89-91, figs. 75-76.
 - Description, natural history, rayages, means against, and figures of larva and imago of Anchylopera [= Phoxopteris] fragariæ, n. sp.; figures all stages of Emphytus [= Harpiphorus] maculatus; description of its larva and pupa, its natural history, and means against its rayages.
- 500. [Walsh, B. D., and C. V. Riley.] Fungoid growths. < Amer. Ent., January, 1869, v. 1, pp. 91-92.
 - Mention of instances of the growth of fungi on living plants and animals; letter of S. H. Y. Early on the occurrence of fungoid growths on the larvæ of Lachnosterna fusca.
- 501. [Walsh, B. D., and C. V. Riley.] Plums for the million. < Amer. Ent., January, 1869, v. 1, pp. 92-93.
 - Means against Conotrachelus nenuphar; notes on varieties of plum exempt from the attacks of the same.
- 502. [Walsh, B. D., and C. V. Riley.] The "Colorado grasshopper." <Amer. Ent., January, 1869, v. 1, pp. 95-96.
 - Answer to inquiries of W. N. Byers and V. Devinny; specific names indicating particular districts can not be changed because the insect is found in other regions.

- 503. [Walsh, B. D., and C. V. Riley.] Universal remedies. <Amer. Ent., January, 1869, v. 1, p. 97.
 - Criticism of an advertisement of "Best's patent fruit tree and vine invigorator."
- 504. [WALSH, B. D., and C. V. RILEY.] Complimentary. <Amer. Ent., January, 1869, v. 1, p. 98.

 Notices of several notices of the American Entomologist.
- 505. [Walsh, B. D., and C. V. Riley.] Paper-makers. < Amer. Ent., January, 1869, v. 1, p. 98.

Comparison of the manufacture of paper by man and by the Vespida.

- 506. [WALSH, B. D., and C. V. RILEY.] On our table. <Amer. Ent., January, 1869, v. 1, p. 98.

 Notices of L'Insectologie agricole—Cecil's books of natural history.
- 507. [WALSH, B. D., and C. V. RILEY.] Look out for the eggs of the apple-tree plant-louse. <Amer. Ent., January, 1869, v. 1, p. 99. Abundance of eggs of Aphis mali in winter of 1868-'69, in Missouri and Illinois; means against the same.
- 508. [WALSH, B. D., and C. V. RILEY.] Greenhouse plants [= pests]. <Amer. Ent., January, 1869, v. 1, p. 99.

 Means against Aphididæ, Coccidæ, and Tetranychus telarius.
- 509. [Walsh, B. D., and C. V. Riley.] How to hatch pupe. <Amer. Ent., January, 1869, v. 1, p. 99.

 Answer to inquiry of D. P. Smith; methods of raising insects.
- 510. [WALSH, B. D., and C. V. RILEY.] Drug-store pests. < Amer. Ent., January, 1869, v. 1, p. 99.

 Answer to inquiry of J. M. Good; characters of larva of undertermined ptinid; food-habits of *Ptinus brunneus*; food-habits, characters, and means
- 511. [Walsh, B. D., and C. V. Riley.] Apple-tree worms. < Amer. Ent., January, 1869, v. 1, p. 99.

against Calandra oryzæ and C. granaria. (See No. 551.)

- Answer to inquiry of J. J. Thomas; habits, food-plants, and means against Phycita nebulo [=Acrobasis indiginella].
- 512. [WALSH, B. D., and C. V. RILEY.] Crane-flylarvæ. Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiries of R. D. Alexander and Hickman; habits of *Tipula* sp.; food-habits of *T. trivittata*.
- 513. [Walsh, B. D., and C. V. Riley.] Punctured grape canes. <Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of B. L. Kingsbury; description of punctures in grape canes probably caused by *Orocharis saltator*; means against tree-crickets.
- 514. [Walsh, B. D., and C. V. Riley.] Insects to be named. < Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of W. R. Marine; identification of several insects found in apple and peach nurseries.
- 515. [Walsh, B. D., and C. V. Riley.] Supposed cause of yellows in peach trees. <Amer. Ent., January, 1869, v. 1, p. 100, fig. 77.

 Answer to inquiry of C. H. Roberts; yellows of peach-trees a vegetable disease; figure of healthy and diseased limb; affected trees more subject to

the attacks of insects than healthy trees.

- 516. [Walsh, B. D., and C. V. Riley.] Eggs of katydid. < Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of A. A. Hilliard; character of the eggs of *Platyphyllum* [= Cyrtophyllus] concavus.
- 517. [Walsh, B. D., and C. V. Riley.] Apple-twig borer. <Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of J. T. Zimmerman; ravages of Bostrichus [= Amphicerus] bicaudatus.
- 518. [Walsh, B. D., and C. V. Riley.] Galls and their architects. <Amer. Ent., February, 1869, v. 1, pp. 101–110, figs. 78–90.
 - Definition, classification, and variation of galls; accounts of some galls made by Cynipidæ, Cecidomyidæ, and Aphididæ and of the gall-makers; natural history, descriptions, and figures of Cynips [= Amphibolips] q.-spongifica, C. [= A.] q.-inanis, C. [= A.] q.-prnnus n. sp., Cecidomyia s.-strobiloides, C. s.-brassicoides, C. v.-pomum n. sp., C. v.-coryloides n. sp., Pemphigus ragabundus, P. rhois and Colopha ulmicola; description of the image of Cynips q.-prunus and of Pemphigus ulmifusus n. sp., and of the larvæ of the new species of Cecidomyia; dimorphism of gall-makers; presence of inquilines and parasites in galls. (See No. 821.)
- 519. [Walsh, B. D., and C. V. Riley.] The bogus Colorado potatobug, Doryphora juncta, Germar. <Amer. Ent., February, 1869, v. 1, p. 110.

Food-habits of D. juncta.

- 520. [Walsh, B. D., and C. V. Riley.] Ants and aphides. <Amer. Ent., February, 1869, v. 1, p. 110.
 - Inquiry of W. Batchelor, with answer; secretion of honey-dew by Aphidide and harvesting of the same by ants.
- 521. [Walsh, B. D., and C. V. Riley.] The Tarantula of Texas, Mygale hentzii, Girard. Samer. Ent., February, 1869, v. 1, p. 11, fig. 91.
 - Figure of Mygale hentzii; habits of Pompilus [=Pepsis] formosa; quotes from G. Lincecum's "The Tarantula killers of Texas" (Amer. Nat., v. 1, pp. 137-141).
- 522. [Walsh, B. D., and C. V. Riley.] The melancholy chafer. <Amer. Ent., February, 1869, v. 1, p. 111.

Ravages and figure of the image of Euphoria melancholica.

- 523. [Walsh, B. D., and C. V. Riley.] Apple worms (Carpocapsa pomonella, Linn.). <Amer. Ent., February, 1869, v. 1, pp. 112–114, fig. 93.
 - Natural history, ravages, and means against Carpocapsa pomonella; figures injured apple, larva, pupa, and imago.
- 524. [Walsh, B. D., and C. V. Riley.] The asparagus beetle (*Crioceris asparagi*, Linn.). <Amer. Ent., February, 1869, v. 1, pp. 114-115, fig. 94.
 - Introduction into the United States; natural history, ravages, parasites of, and means against *Crioceris asparagi*; figure of eggs, larvæ, and imago of the same.

- 525. [WALSH, B. D., and C. V. RILEY.] A popular delusion. < Amer. Ent., February, 1869, v. 1, p. 116.
 - No insect passes through all the stages of its growth within one day; life history of *Ephemeridæ*.
- 526. [Walsh, B. D., and C. V. Riley.] The squirrel bot. < Amer. Ent., February, 1869, v. 1, v. 117.
 - Comments on paper by S. S. Rathvon; emasculation of the striped squirrel by Cuterebra buceata.
- 527. [Walsh, B. D., and C. V. Riley.] [Periodical Cicada.] < Amer. Ent., February, 1869, v. 1, p. 117.
 - Comments on letter of R. H. Warder; oviposition of Cicada [= Tibicen] septendecim in evergreens.
- 528. [Walsh, B. D., and C. V. Riley.] Transformations of insects. <Amer. Ent., February, 1869, v. 1, p. 118.

 Brief statement of the stages of growth of insects.
- 529. [Walsh, B. D., and C. V. Riley.] Do bees injure raspberries? <Amer. Ent., February, 1869, v. 1, p. 118.
 - Comments on discussion before New York Frnit Growers' Club as to the effects of bees on flowers and frnits.
- 530. [Walsh, B. D., and C. V. Riley.] Best's fruit-tree invigorator again. <Amer. Ent., February, 1869, v. 1, p. 119.

 Notice of invention of a new "invigorator" by B. Best.
- 531. [WALSH, B. D., and C. V. RILEY.] The insect extinguisher, by Joseph Treat, N. J. < Amer. Ent., February, 1869, v. 1, p. 119. Critical review of pamphlet by J. Treat.
- 532. [Walsh, B. D., and C. V. Riley.] More good words. <Amer. Ent., February, 1869, v. 1, p. 119.

 Notice of favorable notices of the American Entomologist.
- 533. [Walsh, B. D., and C. V. Riley.] Lacewing fly. < Amer. Ent., February, 1869, v. 1, p. 119.
 - Answer to inquiry of J. Huggins; characters of Chrysopa sp.; its hibernation as a pupa and imago.
- 534. [Walsh, B. D., and C. V. Riley.] Gigantic water-bug. < Amer. Ent., February, 1869, v. 1, p. 119.
 - Answer to inquiry of S. E. Munford; food-habits of Belostoma grandis [=americanum].
- 535. [Walsh, B. D., and C. V. Riley.] The white-marked tussock moth again. <Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of A. S. Fuller; mention of seven parasites attacking Orgyia leueostigma.
- 536. [Walsh, B. D., and C. V. Riley.] Grasshopper eggs—will they hatch? <Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of C. J. Jones; frosts not likely to prevent the hatching of the eggs of Caloptenus spretus.
- 537. [Walsh, B. D., and C. V. Riley.] Injured apple-trees. < Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of G. C. Broadhead; means against Chrysobothris femorata,

- 538. [Walsh, B. D., and C. V. Riley.] Bag worms. < Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of W. W. Butterfield; means against Thyridopteryx ephemer wform is.
- 539. [Walsh, B. D., and C. V. Riley.] Eggs of the apple-tree plant-louse. <Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of M. W. Seaman; effect of frosts upon the cggs and larva of Aphis mali.
- 540. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of J. B. Merwin; identification of insects sent; characters of Vanessa antiopa and Cynthia [=Pyrameis] atalanta.
- 541. [WALSH, B. D., and C. V. RILEY.] White grubs. < Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of W. C. Holmes; undetermined white-grub injuring grass and osage orange.
- 542. [Walsh, B. D., and C. V. Riley.] The polyphemus moth, Attacus polyphemus, Linn. <Amer. Ent., March, 1869, v. 1, pp. 121-122, fig. 95.
 - Attaeus [= Telea] polyphemus, A. cynthia, and Bombyx [=Seriearia] mori as silk-producers; food-plants, description of larva, habits, and seasons of A. [= T.] polyphemus; figure of the imago; ravages of A. cecropia; antennæ mistaken for wings; mention of Lepidoptera having fissured wings.
 - 543. [Walsh, B. D., and C. V. Riley.] Wasps and their habits. <Amer. Ent., March, 1869, v. 1, pp. 122-143, figs. 96-112.
 - Structure, classification, and habits of North American predatory Hymenoptera; comparative characters of the digger and true wasps; figure to show the folding of the wing in true wasps; habits and figures of images of Chlorion caruleum, Bembex fasciata, Sphex iehneumonea, Ammophila pictipennis, Pepsis formosa, Stizus grandis [=Sphecius speciosus], S. [=S.] speciosus, Pelopaus lunatus [=cementarius], Agenia bombycina, Trypoxylon albitarse, Ceropales rufiventris, Eumenes fraternus, Vespa maculata, and Polistes rubiginosus; figure of image of Cryptus [=Linoceras] juneeus and of several nests of wasps. (Sec No. 375.)
- 544. [WALSH, B. D., and C. V. RILEY.] Do toads eat worker bees? <Amer. Ent., March, 1869, v. 1, p. 144.
 - Reprint and review of article by C. Dadant, and of comments of editors of American Bee Journal; usefulness of toads.
- 545. [WALSH, B. D., and C. V. RILEY.] Best's invigorator once again. <Amer. Ent., March, 1869, v. 1, p. 145.
 - Condemnation of B. Best's patent fruit tree and vine invigorator.
- 546. [Walsh, B. D., and C. V. Riley.] On our table. < Amer. Ent., March, 1869, v. 1, p. 146.
 - Notices of Nos. 1 and 2 of Le Naturaliste Canadien.
- 547. [Walsh, B. D., and C. V. Riley.] Sugar-tree borer. <Δmer. Ent., March, 1869, v. 1, p. 146.
 - Answer to inquiry of E. Simms; characters and means against Arhopalus [=Plagionotus] speciosus.

No. 510.

- 548. [Walsh, B. D., and C. V. Riley.] Swellings on apple scions. <Amer. Ent., March, 1869, v. 1, p. 146.
 - Answer to inquiry of W. Colwell; dipterous galls on apple scions and basswood twigs.
- 549. [Walsh, B. D., and C. V. Riley.] Gas-waste vs. Curculio. <Amer. Ent., March, 1869, v. 1, p. 147.

Answer to inquiry of H.; merits of gas-waste as an insect destroyer.

- 550. [Walsh, B. D., and C. V. Riley.] Eggs of the apple-tree plant-louse again. < Amer. Ent., March, 1869, v. 1, p. 147.

 Answers to inquiries of C. Williams and W. L. French; means against Aphis mali.
- 551. [Walsh, B. D., and C. V. Riley.] Drug-store pests. < Amer. Ent., March, 1869, v. 1, p. 147.

 Answer to inquiry of J. M. Good; food-habits of Lasioderma serricorne. Sec
- 552. [Walsh, B. D., and C. V. Riley.] Small galls and minings on apple-twigs. <Amer. Ent., March, 1869, v. 1, p. 147.

 Answer to inquiry of O. O. A. Gardner; characters of undetermined galls and borings in twigs of apple-trees.
 - 553. [Walsh, B. D., and C. V. Riley.] Stinking bugs. < Amer. Ent., March, 1869, v. 1, p. 147.

 Answer tooinquiry of C. L. Janney; habits of an undescribed Brachyrhynchus.
 - 554. [Walsh, B. D., and C. V. Riley.] Eggs in peach-twigs. < Amer. Ent., March, 1869, v. 1, p. 147.

 Answer to inquiry of G. Fisher; character of eggs of *Ecanthus niveus*.
 - 555. [Walsh, B. D., and C. V. Riley.] Flat-headed apple-tree borer. <Amer. Ent., March, 1869, v. 1, p. 147. Answer to inquiry of B. F. Mudge; means against *Chrysobothris femorata*.
- 556. [Walsh, B. D., and C. V. Riley.] A bundle of entomological queries. <Amer. Ent., March, 1869, v. 1, p. 148.

 Answer to inquiry of W. W. Butterfield; mentions several desirable entomological books.
- 557. [Walsh, B. D., and C. V. Riley.] Grape-berry moth. < Amer. Ent., March, 1869, v. i, p. 148.

 Answer to inquiry of M. C. Read; means against *Penthina vitivorana* [= Eudemis botrana].
- 558. [Walsh, B. D., and C. V. Riley.] Hairy caterpillar. <Amer. Ent., March, 1869, v. 1, p. 148.

 Answer to inquiry of T. S. Gold; characters of the larva and image of Arctia [=Pyrrharctia] isabella.
- 559. [WALSH, B. D., and C. V. RILEY.] Borer in plum-twigs. <Amer. Ent., March, 1869, v. 1, p. 148.

 Answer to inquiry of W. Coldwell: undetermined herer allied to the cold.

Answer to inquiry of W. Caldwell; undetermined borer, allied to the oak-pruner, in plum-twigs. See No. 606.

Smithsternes ,

- 560. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., March, 1869, v. 1, p. 148.
 - Answer to inquiry of W. W. Butterfield; identification of insects sent; comparative characters of Arhopalus [= Cyllene] pictus and A. [= C.] robiniæ; variation in Clytus [= Neoclytus] capræa; generie characters of Arhopalus and Clytus. (See No. 582.)
- 561. [Walsh, B. D., and C. V. Riley.] Pear-root borer. < Amer. Eut., March, 1869, v. 1, p. 148.
 - Answer to inquiry of P. Earle; larva of Prionus laticollis injurious to the pear and grape.
- 562. [WALSH, B. D., and C. V. RILEY.] Insects named. < Amer. Ent., March, 1869, v. 1, p. 148.
 - Answer to inquiry of M. H. Boye; Gryllotalpa longipennis [= columbia] a dimorphic form of G. borealis.
- 563. [WALSH, B. D., and C. V. RILEY.] The joint-worm (*Isosoma hordei* Harris). <Amer. Ent., April, 1869, v. 1, pp. 149-158, figs. 113-118.
 - Natural history, variation, parasites, ravages, and means against *Isosoma hordei*; unity of habits in insects; generic classification of the joint-worm fly; figures male and female imago; figures \mathcal{E} and \mathcal{Q} imagos of *Semiotellus chalcidiphagus* n. sp., *Eurytoma* sp., and *Decatoma* sp., and of \mathcal{Q} imago of *Antigaster* [=*Eupelmus*] *mirabilis*.
- 564. [Walsh, B. D., and C. V. Riley.] The wavy-striped flea-beetle. (Haltica [Phyllotreta] striolata Illiger). <Amer. Ent., April, 1869, v. 1, pp. 158-159, fig. 119.
 - Description, habits, and ravages of *Phyllotreta striolata* [=vittata]; figures larva, pupa, and imago of the same; quotes from H. Shimer's "The wavy-striped flea-beetle" (Amcr. Nat., December, 1868, v. 2, pp. 514-517).
- 565. [Walsh, B. D., and C. V. Riley.] Concerning certain smart bugs. <Amer. Ent., April, 1869, v. 1, p. 160.
 - Preferences of Phylloxera vitifolia [=vastatrix] and Macrodactylus subspinosus for certain varieties of grape-vines, of Doryphora 10-lineata and Lema trilineata for certain varieties of potato-vines, and of Mytilaspis pomicorticis [=pomorum] and Carpocapsa pomonella for certain varieties of apple.
- 566. [Walsh, B. D., and C. V. Riley.] Curculio remedies. < Amer. Ent., April, 1869, v. 1, p. 161.
 - Comments on essay of L. C. Francis on the plum; coal oil not effective against Conotrachelus nenuphar.
- 567. [Walsh, B. D., and C. V. Riley.] How great wits jump together. <Amer. Ent., April, 1869, v. 1, p. 161.
 - Stopping holes with hard soap ineffective against borers.
- 568. [Walsh, B. D., and C. V. Riley.] On our table. <Amer. Ent., April, 1869, v. 1, p. 165.
 - Notices of Harris's Insects injurious to Vegetation, L'Insectologie agricole, and several other works not entomological.

- 569. [Walsh, B. D., and C. V. Riley.] Eggs of the oblong-winged katydid. Amer. Ent., April, 1869, v..1, p. 166, fig. 120.
 - Answer to inquiries of T. A. Thop, J. L. Rice, B. J. Campbell, H. Cheeney, and A. MeMoore; eggs of *Phylloptera* [= Amblycorypha] oblongifolia eompared with those of *Platyphyllum* [= Cyrtophyllus] concavum; figures of the eggs of A. oblongifolia.
- 570. [Walsh, B. D., and C. V. Riley.] Mossy rose gall. < Amer. Ent., April, 1869, v. 1, p. 166.

Answer to inquiry of F. W. Collins; characters of Rhodites rosa and its gall.

- 571. [Walsh, B. D., and C. V. Riley.] Caterpillars on lombardy poplars. <Amer. Ent., April, 1869, v. 1, p. 166.
 - Answer to inquiry of J. F. Gurley; characters of larva, habits, food-plants, and parasites of Acronycta accricola [= americana].
- 572. [Walsh, B. D., and C. V. Riley.] Museum pests. < Amer. Ent., April, 1869, v. 1, p. 166, fig. 121.
 - Answer to inquiries of F. W. Hoit, jr., G. M. L., and J. Huggins; figures larva, pupa, and imago of Anthrenus sp.; A. varius compared with A. museovum.
- 573. [Walsh, B. D., and C. V. Riley.] Cockroach eggs. < Amer. Ent., April, 1869, v. 1, p. 166.
 - Answer to inquiry of H. C. Freeman; character of the egg-cases and adult of *Platamodes unicolor*; occurrence of *Ectobia germanica* in Illinois; ravages of species of *Blattidæ*.
- 574. [Walsh, B. D., and C. V. Riley.] Apple-tree leaf-crumplers. Amer. Ent., April, 1869, v. 1, p. 166.
 - Answer to inquiry of J. F. Jones; result of the work of the larvæ of *Phycita* nebulo [= Acrobasis indiginella] on the growth of apple-trees.
- 575. [Walsh, B. D., and C. V. Riley.] A most precious bug. < Amer. Ent., April, 1869, v. 1, p. 167.
 - Answer to inquiry of E. S. Holmes; characters and variety of *Hylecætus* lugubris; destructiveness of Lymexylon navale; use of the maxillary palpi in the 3 3 of Lymexylidæ.
- 576. [Walsh, B. D., and C. V. Riley.] Blackberry-cane borers. <Amer. Ent., April, 1869, v. 1, p. 167.
 - Answer to inquiry of C. Parry; characters and habits of larva of unnamed borer [= Bembecia marginata]; plants affected by species of Ægeriadæ.
- 577. [WALSH, B. D., and C. V. RILEY.] Rows of eggs in pear-twigs. <Amer. Ent., April, 1869, v. 1, p. 167.
 - Answer to inquiry of H. C. Freeman; characters of unknown eggs found in pear-twigs.
- 578. [Walsh, B. D., and C. V. Riley.] An orchard giving out. <Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. M. Clemens; habits and means against Saperda bivittata [= candida] and Chrysobothris femorata.
- 579. [Walsh, B. D., and C. V. Riley.] Plant-louse eggs on apple and mountain ash. <Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. Stewart; eggs of Aphis mali on apple-twigs and probably on those of mountain ash; Aspidiotus harrisii [= Chionaspis furfurus] infests both trees.

- 580. [Walsh, B. D., and C. V. Riley.] Gigantic rhinoceros beetle. <Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of F. G. Smith; characters of Dynastes tityrus.
- 581. [Walsh, B. D., and C. V. Riley.] Bee queries. <Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. R. Howard; Galleria cereana always injurious; a new swarm composed of both old and new bees.
- 582. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. W. Butterfield; characters of Clytus [=Neoclytus] capræa. (See No. 560.)
- 583. [WALSH, B. D., and C. V. RILEY.] Elm-tree borer. < Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. M. Gregory; Saperda lateralis and Dryobius sexfasciatus attack the elm in the larva state. (See No. 696.)
- 584. [Walsh, B. D., and C. V. Riley.] The chinch-bug (*Micropus leucopterus* Say). <Amer. Ent., 1869, v. 1, May, pp. 169–177, fig. 122; June, pp. 194–199, figs. 135–139. Reprint: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 15–37, figs. 1–10. See No. 1127 for synopsis of contents.
- 585. [Walsh, B. D., and C. V. Riley.] The grape-berry moth (*Penthina vitivorana*, Packard). <Amer. Ent., May, 1869, v. 1, pp. 177–179, figs. 123–125.
 - Natural history, ravages, and means against *Penthina vitivorana* [= *Eudemis botrana*]; figures, larva, pupa, eocoon, imago, and injured grape.
- 586. [Walsh, B. D., and C. V. Riley.] Poisonous flour. < Amer. Ent., May, 1869, v. 1, p. 179.
 - Application of the term weevil; vesicatory properties of Sitophilus [= Calandra] granaria; poisonous nature of flour made from wheat infested by these insects.
- 587. [WALSH, B. D., and C. V. RILEY.] Mounding peach-trees. < Amer. Ent., May, 1869, v. 1, pp. 180–181, fig. 126.
 - Mounding as a means against $Sannina\ exitiosa$; figures \mathcal{J} and \mathcal{Q} of the same.
- 588. [Walsh, B. D., and C. V. Riley.] Going it blind. <Amer. Ent., May, 1869, v. 1, pp. 182-183.

 Remarks on the failure of persons to observe correctly.
- 589. [Walsh, B. D., and C. V. Riley.] Another new Curculio humbug. <Amer. Ent., May, 1869, v. 1, p. 183.
 - Extract from "agricultural paper," with comments on proposed means against Conotrachelus nenuphar.
- 590. [Walsh, B. D., and C. V. Riley.] Rear-horses vs. grasshoppers. <Amer. Ent., May, 1869, v. 1, p. 184, figs. 127–128.
 - Vernacular names of Mautis [= Phasmomantis] carolina; its usefulness; figures of eggs and \mathcal{E} and \mathcal{E} imago.
- 591. [Walsh, B. D., and C. V. Riley.] Apple-tree plant-lice. (Aphis mali, Linn.) < Amer. Ent., May, 1869, v. 1, p. 184.
 - Abundance of Aphis mali in several localities; its comparative harmlessness; means against it.

- 592. [Walsh, B. D., and C. V. Riley.] Crack-jaw names. < Amer. Ent., May, 1869, v. 1, p. 184.
 - Combination of vernacular and technical names for the accommodation of different classes of readers.
- 593. [Walsh, B. D., and C. V. Riley.] Send plenty of specimens. <Amer. Ent., May, 1869, v. 1, p. 185.
 - Reasons why several specimens of insects should be sent for examination.
- 594. [Walsh, B. D., and C. V. Riley.] White-grub fungus. < Amer. Ent., May, 1869, v. 1, p. 186, fig. 129.
 - Answer to inquiries of J. Smith and of T. J. Freeman; larva of *Lachnosterna* fusca infested with *Cordyceps ravenelii*; figure of infested larva.
- 595. [Walsh, B. D., and C. V. Riley.] "Buck fly." < Amer. Ent., May, 1869, v. 1, p. 186.
 - Answer to inquiry of G. W. Copley; characters of larva and image of Saturnia [= Hemileuca] maia; food-plants of the larva.
- 596. [Walsh, B. D., and C. V. Riley.] Swarms of minute flies in rooms. <Amer. Ent., May, 1869, v. 1, p. 186.

 Answer to inquiry of S. S. Rathvon; habitat of larva of Sciara sp.
- 597. [WALSH, B. D., and C. V. RILEY.] Worms in osage orange seed. <Amer. Ent., May, 1869, v. 1, p. 186.

 Answer to inquiry of A. Plant; habits of the larva of an undetermined dip
 - teron found among osage orange seed.
- 598. [WALSH, B. D., and C. V. RILEY.] Insects to be named. <Amer. Ent., May, 1869, v. 1, p. 186.

 Answer to inquiry of X. Q. Z.; value of names to a collection of insects.
- 599. [Walsh, B. D., and C. V. Riley.] The spotted lady-bird. < Amer. Ent., May, 1869, v. 1, p. 186, fig. 130.
 - Answer to inquiry of R. Seevers; value of *Hippodamia* [= *Megilla*] *maculata* as a destroyer of noxious insects; figure of the same.
- 600. [Walsh, B. D., and C. V. Riley.] Eggs of the white-marked tussock moth. <Amer. Ent., May, 1869, v. 1, p. 186.
 - Answer to inquiry of S. G. Knight; characters of eggs of Orgyia leucostigma.
- 601. [WALSH, B. D., and C. V. RILEY.] Owl's pellets. <Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of C. H. G.; presence of injurious insects in the pellets disgorged by owls [= hawks]. See No. 643.
- 602. [Walsh, B. D., and C. V. Riley.] Crab-apple borers. < Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of J. Huggins; characters of undetermined lepidopterous borer infesting crab-apple trees.
- 603. [Walsh, B. D., and C. V. Riley.] Mosquitoes. < Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of A. M. Abbott; life-habits of Culicida; benefits derived from the same.
- 604. [WALSH, B. D., and C. V. RILEY.] Large silken cocoon. < Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of W. W. Butterfield; food-habits of Attacus promethea.

- 605. [Walsh, B. D., and C. V. Riley.] Bugs in alcohol. <Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of D. P. Smith; alcohol as a means of preserving insects.
- 606. [Walsh, B. D., and C. V. Riley.] Borer in plum-twig. < Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of W. Colwell; *Elaphidion parallelum* [=villosum] bred from plum-twigs; it does not prune the twig. See No. 559.
- 607. [Walsh, B. D., and C. V. Riley.] Eggs of cut-worm moth. <Amer. Ent., May, 1869, v. 1, p. 188, fig. 131.
 - Answer to inquiries of G. Pauls, T. A. Thorp, and E. S. Foster; characters and figure of eggs of *Agrotis inermis* [= saucia]; characters and habits of the larva.
- 608. [Walsh, B. D., and C. V. Riley.] Snow fleas. <Amer. Ent., May, 1869, v. 1, p. 188.
 - Answer to inquiry of H. H. G. Bradt; habits and food of *Podura* [= Achorutes] nivicola.
- 609. [Walsh, B. D., and C. V. Riley.] Fuzzy galls on blackberry-twigs. <Amer. Ent., May, 1869, v. 1, p. 188.
 - Answer to inquiry of J. Huggins; characters of Diastrophus cuscutæformis and its gall.
- 610. [Walsh, B. D., and C. V. Riley.] Pithy galls on blackberry-twigs. <Amer. Ent., May, 1869, v. 1, p. 188.
 - Answer to inquiry of T. W. Gordon; characters of the gall made by Diastro-phus nebulosus.
- 611. [Walsh, B. D., and C. V. Riley.] Moth eggs. <Amer. Ent., May, 1869, v. 1, p. 188.
 - Answer to inquiries of A. M. Shute and of J. Huggins; characters f the eggs of an undetermined moth.
- 612. [Walsh, B. D., and C. V. Riley.] Horse-hair snakes. < Amer. Ent., May, 1869, v. 1, p. 188.
 - Answer to inquiry of A. M. Abbott; characters and habitats of Gordiacea.
- 613. [Walsh, B. D., and C. V. Riley.] Imitative butterflies. <Amer. Ent., June, 1869, v. 1, pp. 189–193, figs. 132–134.
 - Immunity of Danaidæ from and liability of Pieridæ to the attacks of predatory animals; mimicry of Danaidæ by Pieridæ and of Danais archippus by Limenitis disippus; hibernating habits and description of the larva of the latter, with figures of its larva, chrysalis, imago, and hibernaculum; figure of Danais archippus; theory of the origin of mimicry.
- 614. [Walsh, B. D., and C. V. Riley.] Cabbage-worms upon gilly-flowers. <Amer. Ent., June, 1869, v. 1, p. 199.

 Habits and food-plants, seasons, and synonyms of Plutella cruciferarum.
- 615. [Walsh, B. D., and C. V. Riley.] "Wasps and their habits." <Amer. Ent., June, 1869, v. 1, p. 200.
 - Comments on letter of S. S. Rathvon; species having essentially different habits though externally indistinguishable should be considered specifically distinct.
- 616. [Walsh, B. D., and C. V. Riley.] The social wasps. < Amer. Ent., June, 1869, v. 1, p. 201.
 - Comments on paper by D. A. A. Nichols; habits of Vespa crabro.

- 617. [Walsh, B. D., and C. V. Riley.] Mounding peach-trees. < Amer. Ent., June, 1869, v. 1, pp. 201–202.
 - Comments on letter of R. L. Wells; success of the mounding system against Egeria [= Sannina] exitiosa.
- 618. [Walsh, B. D., and C. V. Riley.] Out of evil there cometh good. <Amer. Ent., June, 1869, v. 1, p. 202.
 - Probable abundance of the fruit crop in southern Illinois and iu Missouri in 1869 due to the pruning of the trees by *Tibicen septendecim* in 1868.
- 619. [Walsh, B.D., and C. V. Riley.] The periodical Cicada. < Amer. Ent., June, 1869, v. 1, p. 202.
 - Extract from No. 474; request for information of the appearance of *Cicada* [= *Tibicen*] septendecim in any part of the United States in 1869.
- 620. [Walsh, B. D., and C. V. Riley.] The Curculio scarcer than last year. <Amer. Ent., June, 1869, v. 1, p. 202.
 - Extract from letter of A. M. Brown; comparative scarcity of Conotrachelus nenuphar in 1869.
- 621. [Walsh, B. D., and C. V. Riley.] The American Entomological Society. <Amer. Ent., June, 1869, v. 1, p. 203.
 - Notice of the formation, publications, aims, and needs of the American Eutomological Society; proposition for the raising of a fund for the support of the society.
- 622. [Walsh, B. D., and C. V. Riley.] Remarkable peculiarity in the insect world. <Amer. Ent., June, 1869, v. 1, p. 204.
 - Insects contrasted with animals of other groups; as a rule they produce but one brood of offspring in the course of their lives.
- 623. [Walsh, B. D., and C. V. Riley.] On our table. <Amer. Ent., June, 1869, v. 1, pp. 204-205.
 - Notices of: The Harris correspondence; The butterflies of North America, by W. H. Edwards; Guide to the study of insects, by A. S. Packard, jr.
- 624. [Walsh, B. D., and C. V. Riley.] Cannibal mites. <Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of C. S. Davis; *Trombidium* sp. found preying on grass-hopper eggs.
- 625. [Walsh, B. D., and C. V. Riley.] Gnats. < Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of W. O. Hiskey; distinctive characters of Culex and Chironomus; appearance of "clouds" of Chironomus.
- 626. [Walsh, B. D., and C. V. Riley.] Cut-worms severing cabbage plants. <Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of N. C. Burch; characters of the larva of Agrotis telifera [= ypsilon].
- 627. [Walsh, B. D., and C. V. Riley.] Beetle named. < Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of J. M. Shaffer; Anisodactylus baltimorensis flying in great numbers at Fairfield, Iowa.
- 628. [WALSH, B. D., and C. V. RILEY.] Tiger beetles. < Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of J. M. Shaffer; habitat of Cicindela vulgaris.

- 629. [Walsh, B. D., and C. V. Riley.] Cocoons and chrysalids named. < Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of A. S. Fuller; characters of larva and image of Cerato-campa [= Citheronia] regalis; food-plants of its larva; characters of the cocoons of Attacus promethea, A. cecropia, A. [= Telea] polyphemus. and of Thyridopteryx ephemeræformis.
- 630. [Walsh, B. D., and C. V. Riley.] White-lined morning Sphinx. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of S. Blanchard; characters, transformations, habits, and distribution of *Deilephila lineata*; food-plants of its larva.
- 631. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of W. W. Butterfield; identification of insects sent; differences between the spring and autumn broods of *Drasteria erechtea*.
- 632. [Walsh, B. D., and C. V. Riley.] Peach-twig borer. < Amer. Ent., June, 1869, v. 1, p. 206, fig. 140.
 - Answer to inquiry of W. Muir; characters, habits, and figure of the larva of Gortyna nitela.
- 633. [Walsh, B. D., and C. V. Riley.] The apple-twig borer. < Amer. Ent., June, 1869, v. 1, p. 206, fig. 141.
 - Answer to inquiry of A. Hinckley; habits, sexual differences, and figure of Bostrichus [= Amphicerus] bicaudatus.
- 634. [Walsh, B. D., and C. V. Riley.] Apple-tree borers on south side of trees. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of J. F. Wielandy; preference shown for the south and southwest sides of trees by the larva of *Chrysobothris femorata*.
- 635. [Walsh, B. D., and C. V. Riley.] Cocoons of the Cecropia moth. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of T. W. Gordon; comparison between the cocoons of Telea polyphemus and Attacus cecropia.
- 636. [Walsh, B. D., and C. V. Riley.] Flea-beetles. < Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of F. Hecker; characters of an undescribed *Longitarsus* infesting wheat fields.
- 637. [Walsh, B. D., and C. V. Riley.] Strawberry bugs. <Amer. Ent., June, 1869, v. 1, p. 207.
 - Answer to inquiry of J. M. Pearson; characters, ravages, and means against Corimelana pulicaria.
- 638. [Walsh, B. D., and C. V. Riley.] Eggs on apple-trees. < Amer. Ent., June, 1869, v. 1, p. 207, fig. 142.
 - Answer to inquiry of H. Compton; characters of eggs of Sinea diadema; figure of the imago; its value as a destroyer of canker-worms.
- 639. [WALSH, B. D., and C. V. RILEY.] Lady-bird larvæ. <Amer. Ent., June, 1869, v. 1, p. 207, fig. 143.
 - Answer to inquiry of E. S. Foster; characters of the larva of *Hippodamia convergens*; figures its larva, pupa, and imago; usefulness of *Coccinellidæ*.

- 640. [Walsh, B. D., and C. V. Riley.] White-grub fungus, again. <Amer. Ent., June, 1869, v. 1, p. 207, fig. 144.
 - Answer to inquiry of W. C. Holmes; figures larva of Lachnosterna fusca attacked by Cordyceps ravenelii.
- 641. [Walsh, B. D., and C. V. Riley.] Bag-worms. < Amer. Ent., June, 1869, v. 1, p. 207.
 - Answer to inquiry of C. Parry; means against Thyridopteryx ephemeræformis.
- 642. [Walsh, B. D., and C. V. Riley.] Tent-caterpillar. < Amer. Ent., June, 1869, v. 1, p. 208, fig. 145.
 - Answer to inquiry of S. Blanchard; characters of the imago and figures of the early stages of Clisiocampa americana; means against the same.
- 643. [Walsh, B. D., and C. V. Riley.] Hawk's pellets. < Amer. Ent., June, 1869, v. 1, p. 208.
 - Answer to inquiry of C. H. G.; insectivorous habits of *Buteo pennsylvanicus*. See No. 601.
- 644. [Walsh, B.D., and C. V. Riley.] Plant-lice on berberry. < Amer. Ent., June, 1869, v. 1, p. 208.

 Answer to inquiry of J. R. Preston; means against Aphididæ.
- 645. [Walsh, B. D., and C. V. Riley.] Tent-caterpillar of the forest. <Amer. Ent., June, 1869, v. 1, p. 208, fig. 146.
 - Answer to inquiry of G. Whitcomb; characters, habits, parasites, means against, and figure of larva of Clisiocampa sylvatica [= disstria].
- 616. [Walsh, B. D., and C. V. Riley.] Cotton insects. < Amer. Ent., July, 1869, v. 1, pp. 209-214, fig. 147-151.
 - Descriptions and figures of all stages of Aletia xylina and of Heliothis armigera; habits, seasons, ravages, food-plants of, and means against the same; figures larva of Clisiocampa sylvatica [= disstria].
- 647. [Walsh, B. D., and C. V. Riley.] The true Army-worm (*Leucania unipuncta* Haworth). <Amer. Ent., July, 1869, v. 1, pp. 214–217, fig. 152–155.
 - Descriptions and figures of larva, pupa, and imago of *Leucania unipuncta*; seasons, ravages, and enemies of the same; figure of *Exorista militaris* [= Nemoræa leucaniæ].
- 648. [Walsh, B. D., and C. V. Riley.] Belated individuals of the periodical Cicada. <Amer. Ent., July, 1869, v. 1, p. 217.
 - Occurrence of scattering individuals of Tibicon septendecim in years before or after their regular period.
- 649. [Walsh, B. D., and C. V. Riley.] Is the Curculio scarcer than it was last year? <Amer. Ent., July, 1869, v. 1, pp. 217, 218. Record of observations to prove the comparative scarcity of *Conotrachelus nenuphar* during 1869.
- 650. [Walsh, B. D., and C. V. Riley.] To destroy Colorado potato bugs. <Amer. Ent., July, 1869, v. 1, p. 219.
 - Reprint of communication of G. Liddle, with comments; Paris green as a means against *Doryphora* 10-lineata.
- 651. [Walsh, B. D., and C. V. Riley.] Dr. Hull's Curculio-catcher. Amer. Ent., June, 1869, v. 1, pp. 220–221, fig. 156.

 Description and figure of Hull's Curculio-catcher.

- 652. [Walsh, B. D., and C. V. Riley.] The New York weevil (*Ithycerus noveboracensis*, Forster). <Amer. Ent., July, 1869, v. 1, pp. 221–222, fig. 157.
 - Habits, food-plants, distribution, description, and figure of larva and imago of *Ithycerus novcboracensis*; means against the same.
- 653. [Walsh, B. D., and C. V. Riley.] Mounding peach-trees again. <Amer. Ent., July, 1869, v. 1, p. 223.
 - Letter of A. Dean, with comments; characters and habitat of Mycetophila persicæ.
- 654. [Walsh, B. D., and C. V. Riley.] Overcrowded. <Amer. Ent., July, 1869, v. 1, p. 223.
 - Duties of a State entomologist; impossibility of answering inquiries received during the past month.
- 655. [Walsh, B. D., and C. V. Riley.] No air-holes needed in sending insects. Amer. Ent., July, 1869, v. 1, p. 223.

 Directions for sending living insects.
- 656. [Walsh, B. D., and C. V. Riley.] Plum-leaf worms. < Amer. Ent., July, 1869, v. 1, p. 223.

 Answer to inquiry of W. D. Hiskey; characters of larva of Lyda sp.
- 657. [Walsh, B. D., and C. V. Riley.] Seed-corn maggot. <Amer. Ent., July, 1869, v. 1, p. 224, figs. 138-139.

 Answer to inquiry of G. Pauls; characters, ravages, means against, and figure of larva of Anthomyia zeas; characters of the imago; figure of the puparium.
- 658. [Walsh, B. D., and C. V. Riley.] Cut-worms. < Amer. Ent., July, 1869, v. 1, p. 224.

 Answer to inquiry of N. C. Birch; ravages of Agrotis telifera [= ypsilon].
- 659. [WALSH, B. D., and C. V. RILEY.] New York weevil. <Amer. Ent., July, 1869, v. 1, p. 224.

 Answer to inquiries of D. H. Kauffman and of W. D. Turrill; ravages of Ithycerus noveboracensis.
- 661. [Walsh, B. D., and C. V. Riley.] Large green worm in a peach. <Amer. Ent., July, 1869, v. 1, p. 224.

 Answer to inquiry of G. Wilgins; characters and food-habits of an undetermined larva found in a peach.
- 662. [Walsh, B. D., and C. V. Riley.] Ichneumon flies. <Amer. Ent., July, 1869, v. 1, p. 224.

 Answer to inquiry of H. Klinehaus; characters and figure of cocoons of Microgaster sp.
- 663. [WALSH, B. D., and C. V. RILEY.] Raspberry worms. < Amer. Ent., July, 1869, v. 1, p. 224.

 Answer to inquiry of B. Borden; characters and ravages of Sclandria [=Mo-

nophadnus] rubi.

- 664. [Walsh, B. D., and C. V. Riley.] Hairy grape-leaf folders. <Amer. Ent., July, 1869, v. 1, p. 224.
 - Answer to inquiry of A. C. Davis; characters of larva and image of *Pterophorus* [= Oxyptilus] periscelidactylus.
- •665. [Walsh, B. D., and C. V. Riley.] Row of eggs in maple-twigs. <Amer. Ent., July, 1869, v. 1, p. 224.

Answer to inquiry of J. Bower; characters of eggs of undetermined katydid.

- 666. [Walsh, B. D., and C. V. Riley.] Butterfly named. < Amer. Ent., July, 1869, v. 1, p. 224.
 - Answer to inquiry of A. R. Bodley; characters of *Papilio marcellus*; foodplants of its larva.
- 667. [Walsh, B. D., and C. V. Riley.] Grasshopper³s eggs. < Amer. Ent., July, 1869, v. 1, p. 224.
 - Answer to inquiry of E. P. Burlingame; characters of the eggs of an undetermined grasshopper and of the image of *Œdipoda* [= *Dissosteria*] carolina.
- 668. [Walsh, B. D., and C. V. Riley.] Asilus fly larva. < Amer. Ent., July, 1869, v. 1, p. 225, figs. 161-162.
 - Answer to inquiry of G. Pauls; characters, habits, and figure of larva of Asilus sp.; food-habits of larva and imago of A. sericeus; figure of the imago of the same; Trupanea [= Promachus] apivorus as a destroyer of bees.
- 669. [Walsh, B. D., and C. V. Riley.] New insect-foe of the black-berry. <Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of C. Parry; comparative characters of *Aphididæ* and *Psyllidæ*; habits of *Psylla rubi* [= *Trioza tripunctata*].
- 670. [WALSH, B. D., and C. V. RILEY.] Army-worm. < Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of J. H. Butts; ravages and food-plants of *Leucania uni-puncta*.
- 671. [WALSH, B. D., and C. V. RILEY.] Green grape-vine worm. <Amer. Ent., July, 1869, v. 1, p. 225, fig. 163.
 - Answer to inquiry of G. Pauls; characters, food-plants, and figure of larva of Pyrophila pyramidoides.
- 672. [Walsh, B. D., and C. V. Riley.] Rose slug. <Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiries of G. W. Copley and B. S. Morris; characters, ravages, and means against Selandria [= Monostegia] rosw.
- 673. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of J. Weed; food-habits of Leptostylus aculiferus and of Podabrus modestus.
- 674. [Walsh, B. D., and C. V. Riley.] Chrysalis of the virgin tiger moth. <Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of C. Mallinckrodt; characters of Arctia virgo; supposed food-plants of its larva.
- 675. [WALSH, B. D., and C. V. RILEY.] Clover-worms. <Amer. Ent., July, 1869, v. 1, p. 226, fig. 164.
 - Answer to inquiry of E. P. Flanders; characters of Asopia costalis; habits of its larva; figures larva, pupa, cocoon, and imago.

- 676. [Walsh, B. D., and C. V. Riley.] Raspberry brand. < Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiry of J. M. Beecher; larva of small gnat feeding on raspberry-brand.
- 677. [WALSH, B. D., and C. V. RILEY.] Rose-worms. < Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiry of G. S. Grover; character, ravages, food-plants, and means against *Heliothis margidens* [= Pyrrhia exprimens].
- 678. [Walsh, B. D., and C. V. Riley.] Plum-tree plant-lice. <Amer. Ent., July, 1869, v. 1, p. 226, figs. 165-167.
 - Answer to inquiry of T. W. Gordon; characters of Aphis prunifolia and A. [=Myzus] cerasi; ravages, enemies, and means against Aphidida; figures larva of coccinellid, syrphid and hemerobil.
- 679. [WALSH, B. D., and C. V. RILEY.] Ichneumon-flies. < Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiries of S. J. Throp and J. E. Trabue; life-habits of Microgaster sp.?
- 680. [Walsh, B. D., and C. V. Riley.] A new Curculio humbug. <Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiry of E.P. Flanders; uselessness of a patent lamp for destroying Conotrachelus nenuphar.
- 681. [WALSH, B. D., and C. V. RILEY.] Sweet-potato beetles. <Amer. Ent., July, 1869, v. 1; p. 227.
 - Answer to inquiries of Subscriber and A. E. Trabne; food-habits and characters of larva and imago of *Coptocycla aurichalcea* and of *C.* [= Cassida] bivittata.
- 682. [Walsh, B. D., and C. V. Riley.] Injured strawberry and grapevines. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of W. P. Pierson; ravages of Capsus oblineatus [= Lygus pratensis].
- 683. [Walsh, B. D., and C. V. Riley.] Plum-tree insects. < Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of J. F. Waters; characters of Bibio albipennis; food-habits of its larva.
- 684. [Walsh, B. D., and C. V. Riley.] Apple-tree bugs. < Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of W. L. Youse; character of Brochymcna annulata.
- 685. [Walsh, B. D., and C. V. Riley.] Hickory-stem gall-louse. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of B. F. Long; characters of galls of *Phylloxera caryw-caulis*; food-habits of *Thrips*.
- 686. [Walsh, B.D., and C. V. Riley.] Grape-vine leaf-hopper. < Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of R. M. Copeland; means against Tettigonia [= Typhlocyba] vitis.
- 687. [Walsh, B. D., and C. V. Riley.] Twelve-spotted Diabrotica. Amer. Ent., July, 1869, v. 1, p. 227, fig. 168.
 - Answer to inquiries of E. S. Foster and R. D. Parker; characters and means against *Diabrotica* 12-punctata; figure of the same.

- 688. [Walsh, B. D., and C. V. Riley.] The caterpillar of the forest. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of M. McKenzie; food-habits and means against Clisio-campa sylvatica [= disstria].
- 689. [Walsh, B. D., and C. V. Riley.] Strawberry destroyer. < Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of G. W. Copley; habits of unknown strawberry destroyer.
- 690. [Walsh, B. D., and C. V. Riley.] Eggs of bugs on strawberry. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of A. S. Fuller; characters of eggs of unknown reduvid attacked by parasites.
- 691. [Walsh, B. D., and C. V. Riley.] Frog-spittle insects. < Amer. Ent., July, 1869, v. 1, p. 228. Reprint: < Cultivator and Country Gentleman, 29 July, 1869, v. 34, p. 82.
 - Answer to inquiry of J. B. Hartwell; habits, characters, and injuries of Aphrophora quadrangularis.
- 692. [Walsh, B. D., and C. V. Riley.] Eggs of ground-beetle. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of E. J. Ayres; characters of eggs of undetermined groundbeetle found under bark of pear-twigs.
- 693. [Walsh, B. D., and C. V. Riley.] New insect-foe of the potato. <Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of I. Hicks; food-habits of Cassida [= Coptocycla] clavata.
- 694. [Walsh, B. D., and C. V. Riley.] Tomato-stalk borer. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of E. J. Ayres; Gortyna nitela injurious to tomato-stalks.
- 695. [Walsh, B. D., and C. V. Riley.] Breeding cages. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of H. S. Redney; description of cage for breeding insects.
- 696. [Walsh, B. D., and C. V. Riley.] Elm-tree borer. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of W. M. Gregory; characters and food-habits of *Physocnemum brevilineum*. See No. 583.
- 697. [Walsh, B. D., and C. V. Riley.] Peach-twig borer. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of G. Fisher; means against undetermined lepidopterous borer in twigs of peach.
- 698. [Walsh, B. D., and C. V. Riley.] Eggs of periodical Cicada in savin-twig. <Amer. Ent., July 1809, v. 1, p. 228.
 - Answer to inquiry of J. A. Greason; Tibicen septendecim ovipositing in twigs of Juniperus sabina.
- 699. [Walsh, B. D., and C. V. Riley.] Elm-tree saw-fly. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of A. R. Whitney; characters of larva and imago of Cimbex laportei [= americana]; food-plants of its larva.
- 700. [Walsh, B. D., and C. V. Riley.] Snout-beetle. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of H. Kleinhaus; supposed food-habits of Hylobius con-fusus.

701. [Walsh, B. D., and C. V. Riley.] The close of the first volume. <Amer. Ent., August, 1869, v. 1, p. 229.

Prospectus of the second volume of the American Entomologist.

- 702. [Walsh, B. D., and C. V. Riley.] The Royal horned caterpillar. (*Ceratocampa (Citheronia) regalis*, Fabr.). <Amer. Ent., August, 1869, v. 1, pp. 230-231, pl. 1.
 - Habits, seasons, sexual characters, food-plants, and vernacular names of *Citheronia regalis*; descriptions and figures of larva and pupa; figure of larva, pupa, and imago.
- 703. [Walsh, B. D., and C. V. Riley.] Comparative scarcity of the Curculio again. Amer. Ent., August, 1869, v. 1, p. 241.

 Observations on the comparative abundance of Conotrachelus nenuphar during

the early summer of 1869.

- 704. [WALSH, B. D., and C. V. RILEY.] A possible cause of the bee disease. <Amer. Ent., August, 1869, v. 1, pp. 241–242.

 Reprint of article by P. H. Philbrook (Amer. Bee Journal, May, 1869), with comments; dipterous enemies of the honey-bce.
- 705. [WALSH, B.D., and C. V. RILEY.] Ash and mountain ash. < Amer. Ent., August, 1869, v. 1, pp. 243-244.

 Criticisms of paper of H. Shimer (Trans. Ill. State Hortic. Soc., 1868).
- 706. [Walsh, B. D., and C. V. Riley.] How the Curculio flies by night. <Amer. Ent., August, 1869, v. 1, p. 244.

 Extract from lecture by I. P. Trimble, with comment.
- 707. [Walsh, B. D., and C. V. Riley.] The periodical Cicada; our first brood established. Amer. Ent., August, 1869, v. 1, p. 244. Appearance in Connecticut of a brood of Tibicen septendecim in 1869.
- 708. [Walsh, B. D., and C. V. Riley.] Be on the guard. Amer.
 Ent., August, 1869, v. 1, p. 244.

 Necd of care in the transportation of living insects; accidental introduction of Doryphora 10-lineata.
- 709. [WALSH, B. D., and C. V. RILEY.] Prophecy fulfilled. <Amer. Ent., August, 1869, v. 1, p. 244.

 Spread of Doryphora 10-lineata through Michigan.
- 710. [Walsh, B. D., and C. V. Riley.] A poisonous worm. <Amer. Ent., August, 1869, v. 1, p. 245.

 Extract from exchange; larva of *Protoparce celeus* not poisonous.
- 711. [Walsh, B. D., and C. V. Riley.] Wheat midge, alias milk weevil, alias red weevil. <Amer. Ent., August, 1869, v. 1, p. 245.
 - Answer to inquiry of C. Corbit; characters of larva of Cecidomyia [= Di-plosis] tritici; change in heads of wheat.
- 712. [Walsh, B. D., and C. V. Riley.] Large fish-fly. < Amer. Ent., August, 1869, v. 1, p. 245.

Answer to inquiry of A. R. McCutchen; characters of Chauliodes pectinicornis; habits of the larva of C. rastrucornis.

6 ENT

- 713. [Walsh, B. D., and C. V. Riley.] Cottonwood leaf-galls. < Amer. Ent., August, 1869, v. 1, p. 245.
 - Answer to inquiry of J. B. Taylor; habits of *Pemphigus populicaulis*; characters of its gall.
- 714. [WALSH, B. D., and C. V. RILEY.] Insects named., < Amer. Ent., August, 1869, v. 1, p. 245.
 - Answer to inquiry of J. G. Goodrich; characters of Saperda bivittata [= can-dida].
- 715. [Walsh, B. D., and C. V. Riley.] Four-lined leaf-bug on current. <Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of M. B. Bateman; characters, habits, and means against Capsus 4-vittatus [= Pæcilocapsus lineatus].
- 716. [Walsh, B. D., and C. V. Riley.] Bee moth. < Amer. Ent., August, 1869, v. 1, p. 246, fig. 182.
 - Answer to inquiry of S. Blanchard; ravages and means against Galleria cercana; figures larva, pupa, cocoon, and imago of the same.
- 717. [Walsh, B. D., and C. V. Riley.] Canker-worm parasites. <Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of J. Petit; mention of Microgaster sp., parasitic on Anisopteryx.
- 718. [Walsh, B. D., and C. V. Riley.] Beetles swarming about the lawn. < Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of S. Thompson; characters of Gymnetis [=Allorhina] nitida; habits of its larva.
- 719. [WALSH, B. D., and C. V. RILEY.] Tiger-beetle larva. < Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of R. J. Dodge; characters and habits of larva of Cicindelide.
- 720. [Walsh, B. D., and C. V. Riley.] Large compound gall on grapevine. <Amer. Ent., August, 1869, v. 1, p. 247, fig. 183.
 - Answer to inquiries of A. S. Fuller and D. W. Kauffman; characters and figure of gall of *Lasioptera vitis*; habits of its larva; enemy and parasite of the same.
- 721. [WALSH, B. D., and C. V. RILEY.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 247.
 - Answer to inquiry of M. Treat; habits of larva of Temnochila [= Trogosita] virescens, of Aplodes [= Synchlora] rubivora, and of Calosoma calidum.
- 722. [Walsh, B. D., and C. V. Riley.] Destructive larvæ. < Amer. Ent., August, 1869, v. 1, p. 247.
 - Answer to inquiry of B. F. Lee; ravages of an undetermined larva; parasites of Saturnia [= Hemileuca] maia.
- 723. [WALSH, B. D., and C. V. RILEY.] Eggs of tree-cricket on grape-vine. <Amer. Ent., August, 1869, v. 1, p. 247.
 - Answer to inquiry of B. F. Lee; characters of eggs of *Œcanthus niveus*; carnivorous habits of the larva of the same.

- 724. [Walsh, B. D., and C. V. Riley.] Grape-vine leaf-gall. < Amer. Ent, August, 1869, v. 1, p. 248, fig. 184.
 - Answer to inquiry of A. A. Hilliard; characters and figure of the leaf-galls of *Phylloxera vitifoliae* [= vastatrix]; its enemies and means against them; varietics of grape infested; identity of the leaf- and root-galls; criticism of the new genera and families of H. Shimer.
- 725. [Walsh, B. D., and C. V. Riley.] Grape-vine insects. < Amer. Ent., August, 1869, v. 1, p. 248.
 - Answer to inquiry of G. Pauls; characters of *Pelidnota punctata*; its larval habits; characters, habits, and food-plants of *Enchophyllum* [= *Enchenopa*] binotata.
- 726. [WALSH, B. D., and C. V. RILEY.] Museum pests. < Amer. Ent., August, 1869, v. 1, p. 248.

 Answer to inquiry of C. P. Faulkner; ravages of Dermestes lardarius.
- 727. [WALSH, B. D., and C. V. RILEY.] Wheat maggots. < Amer. Ent., August, 1869, v. 1, p. 248.
 - Answer to inquiry of S. K. Faulkner; characters of larva of Meromyza americana injurious to heads of wheat.
- 728. [Walsh, B. D., and C. V. Riley.] Parasites on "hateful grass-hopper." < Amer. Ent., August, 1869, v. 1, p. 249.
 - Answer to inquiry of S. K. Faulkner; characters and habits of Astoma [= Trombidium] locustarum.
- 729. [Walsh, B. D., and C. V. Riley.] Crippled moths. < Amer. Ent., August, 1869, v. 1, p. 248.
 - Answer to inquiry of C. P. Faulkner; conditions needed to enable moths to expand their wings; means by which insects walk on smooth surfaces.
- 730. [Walsh, B. D., and C. V. Riley.] Insects on the oleander. <Amer. Ent., August, 1869, v. 1, p. 249, fig. 185.
 - Answer to inquiry of T. W. Gordon; means against undetermined Coccid on oleander; habits and figure of Chilocorus bivulnerus.
- 731. [WALSH, B. D., and C. V. RILEY.] Insects found on apple-trees. Amer. Ent., August, 1869, v. 1, p. 249.
 - Answer to inquiry of J. W. Waters; identification of the eggs of Reduvius raptatorius [= Sinea diadema]; habits of Chilocorus bivulnerus.
- 732. [Walsh, B. D., and C. V. Riley.] Beetles named. < Amer. Ent., August, 1869, v. 1, p. 249.
 - Answer to inquiry of J. M. Shaffer; Lytta atrata [= Epicauta pennsylvanica] caught on rag-weed.
- 733. [Walsh, B. D., and C. V. Riley.] Gigantic water-bug. < Amer. Ent., August, 1869, v. 1, p. 249, fig. 186.
 - Answer to inquiry of E. M. Downing; habits and figure of Belostoma grandis [= americanum].
- 734. [Walsh, B. D., and C. V. Riley.] Worm eating into green tomatoes. <Amer. Ent., August, 1869, v. 1, p. 249. Answer to inquiry of D. L. Hall; food-habits of Gortyna nitela.

- 735. [WALSH, B. D., and C. V. RILEY.] Miscellaneous. < Amer. Ent., August, 1869, v. 1, p. 249.
 - Answer to inquiry of G. W. Copley; food-habits of Chrysochus auratus, Hemileuca maia, and Aphis [= Myzus] ribis; habits of Lozotania [= Cacacia] rosaceana.
- 736. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of H. A. Munger; food-habits of Cassida pallida [= Coptocycla aurichalcea], Lytta murina [= Macrobasis unicolor], and of the larvæ of the species of Prionus.
- 737. [Walsh, B. D., and C. V. Riley.] Lightning-hoppers. < Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of T. W. Gordon; characters and habits of Pæciloptera pruinosa; vernacular names of Fulgoridæ, Membracidæ, and Jassidæ.
- 738. [Walsh, B. D., and C. V. Riley.] Bag-worms. < Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiries of T. W. Gordon and S. Thompson; food-habits of Thyridopteryx ephemeræformis.
- 739. [Walsh, B. D., and C. V. Riley.] Woolly gall on white oak. Amer. Ent., August, 1869, v. 1, p. 250, fig. 187.
 - Answer to inquiry of A. S. Fuller; characters and figure of the gall of *Cynips* [=Andricus] seminator.
- 740. [Walsh, B. D., and C. V. Riley.] Cabbage pests. < Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of J. A. Williams; characters and habits of an undetermined elaterid larva; means against the larvæ of *Elateridæ*.
- 741. [Walsh, B. D., and C. V. Riley.] Bugs gathering on pear shoots. <Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of E. J. Ayres; characters and habits of Corimelana pulicaria.
- 742. [Walsh, B. D., and C. V. Riley.] Potato-bug. <Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of J. B. Cartwell; food-habits of Cassida [= Coptocycla] clavata.
- 743. [WALSH, B. D., and C. V. RILEY.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of H. T. Birch; Trochilium [= Ægeria] acerni bred from maple.
- 744. [Walsh, B. D., and C. V. Riley.] Apple-tree worms. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of C. Waters; abundance of undetermined moth (*Corycia vestaliata?*); larva of the same on apple-trees.
- 745. [Walsh, B. D., and C. V. Riley.] Oak-fig gall. <Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of A. Fendler; characters of the gall of Cynips [= Bio-rhiza] forticornis and of a guest-fly, Ceroptres ficus, inhabiting the same.

- 746. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of D. L. Phares; irregular appearance of Cicada tredecim [= Tibicen septendecim]; food-habits of Oncideres cingulata.
- 747. [Walsh, B.D., and C. V. Riley.] Small apple-leaf worms. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of H. Compton; means against Spilonota oculana [= Tme-tocera ocellana].
- 748. [Walsh, B. D., and C. V. Riley.] Rose bug on apples. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of A. Dean; food-plants and means against *Macrodactylus* subspinosus.
- 749. [Walsh, B. D., and C. V. Riley.] Unicorn apple-tree caterpillar. <Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of G. C. Brodhead; characters of the larva and imago of Notodonta [= Cwlodasys] unicornis; food-plants of the larva of the same.
- 750. [Walsh, B. D., and C. V. Riley.] Large water beetle. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of S. E. Mumford; characters of Cybister fimbriolatus. See No. 816.
- 751. [Walsh, B. D., and C. V. Riley.] Beetles around peach-trees. <Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of E. Hollister, jr.; characters and habits of *Helops pullus* [= arcus].
 - 752. [Walsh, B. D., and C. V. Riley.] A quick traveler. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of G. C. Brodhead; characters and harmlessness of Cermatia forceps.
 - 753. [Walsh, B. D., and C. V. Riley.] Unknown moth. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of W.G. Barton; distribution of Junonia lavinia in Illinois.
 - 754. [Walsh, B. D., and C. V. Riley.] Corn-borer. <Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of F. M. N.; identification of the larva of Gortyna nitela injurious to corn.
 - 755. [Walsh, B. D., and C. V. Riley.] Horns of stag-beetle. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of C. R. Edwards; larval habits of Lucanus elaphus.
 - 756. [Walsh, B. D., and C. V. Riley.] Imported gooseberry worms. <Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of C. P. Faulkner; means against Nematus ribesii.
 - 757. [WALSH, B. D., and C. V. RILEY.] Worm on bark of walnut-tree. <Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of F. S. Fuller; undetermined notodontoid larva on the bark of walnut.

- 758. [Walsh, B. D., and C. V. Riley.] Rotten root. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of J. M. Beecher; scavenger habits of undetermined larva feeding on dead roots of apple-trees.
- 759. [Walsh, B. D., and C. V. Riley.] Large dragon-fly. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of G. S. Grover; characters and usefulness of *Æschna constricta*.
- 760. [Walsh, B. D., and C. V. Riley.] Stinging larvæ. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of E. H. King; characters and urticating properties of Saturnia [= Hemileuca] maia; peach-blow potatoes avoided by Doryphora 10-lineata.
- 761. [Walsh, B. D., and C. V. Riley.] Raspberry worms. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of E. H. Beebe; characters of larva of Selandria [=Mo-nophadnus] rubi.
- 762. [WALSH, B. D., and C. V. RILEY.] Leaf-galls and caterpillars on the sugarberry. <Amer. Ent., August, 1869, v. 1, p. 252.

 Answer to inquiry of S. L. Scofield; undetermined gall on leaves of Celtis occidentalis; Orgyia leucostigma feeding on the leaves of the same.
- 763. [Walsh, B. D., and C. V. Riley.] Dark grape-worm. < Amer. Ent., August, 1869, v. 1, p. 252.
- Answer to inquiry of T. W. G.; characters and food-plants of *Thyrcus abbotii*.

 764. [WALSH, B. D., and C. V. RILEY.] Too fond of honey. <Amer.
- Ent., August, 1869, v. 1, p. 252.

 Answer to inquiry of T. W. Gordon; Cermatia forceps found in a jar of honey.
- 765. [Walsh, B. D., and C. V. Riley.] Caterpillar of polyphemus moth. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of A. De Wyl; larva of Telea polyphemus feeding on plums.
- 766. [Walsh, B. D., and C. V. Riley.] Why noxious insects increase upon us. <Amer. Ent., September-October, 1869, v. 2, pp. 1-2. Reasons why noxious insects increase; review of articles by E. S. Hull, H. W. Beecher, and Puritan.
- 767. [Walsh, B. D., and C. V. Riley.] Tortoise-beetles. < Amer. Ent., September-October, 1869, v. 2, pp. 2–5, figs. 1–3.
 - Resemblances of insects to other animals or to their surroundings; distribution, food-plants, and figure of Deloyala [= Coptocycla] clavata; description and figure of Physonota quinque-punctata n. sp. [= unipunctata]; figure of the larva of the same; correspondence of structural differences in larva with those in imagos of Cassidida, and with differences in food-plants; list of insects injurious to Solanum; figure of pupa and imago of Chelymorpha cribraria [= argus].
- 768. [Walsh, B. D., and C. V. Riley.] Scientific nomenclature. Amer.Ent., September-October, 1869, v. 2, pp. 5-8.
 - Rules observed in giving specific names to animals and plants; the law of priority; nature and extent of generic subdivisions.

- 769. [WALSH, B. D., and C. V. RILEY.] Killing apple-worms by machinery. <Amer. Ent., September-October, 1869, v. 2, p. 9.
 - Extract from article of L. P. Haskell, with comment; use of rags in place of hay-bands as traps for *Carpocapsa pomonella*; destruction of the insects caught by means of a clothes-wringer.
- 770. [Walsh, B. D., and C. V. Riley.] A potter wasp (Odynerus flavipes? Fabr.). <Amer. Ent., September-October, 1869, v. 2, p. 10, fig. 4.
 - Method employed by Eumenidæ to provision their nests; construction of the same by species of Odynerus; habits of O. flavipes in provisioning a nest with several species of larvæ; figure of the imago of the same and of the nest of a species of Odynerus; structure of wings and habits of Vespidæ, Eumenidæ, and fossorial wasps.
- 771. [WALSH, B. D., and C. V. RILEY.] Tomato-worms not poisonous. <Amer. Ent., September-October, 1869, v. 2, p. 11.
 - Prejndices in regard to certain animals; presence of the horn on the larvæ of almost all Sphingidæ; larva of Protoparce celeus not poisonous.
- 772. [Walsh, B. D., and C. V. Riley.] Gooseberry and current worms. <Amer. Ent., September-October, 1869, v. 2, pp. 12-22, figs. 5-11.
 - Need of precision in nomenclature; relations and distribution of the North American species of Ribes; insect commics of the same; natural history and description of Ellopia [= Eufitchia] ribearia, Nematus ventricosus [= ribesii], and Pristiphora grossulariæ; figures larvæ and imagos of the three species and the pupa of the Eufitchia.
- 773. [Walsh, B. D., and C. V. Riley.] Striped cucumber beetle. <a href="Amer. Ent., September-October, 1869, v. 2, p. 24, figs. 17-19.

 Answer to inquiry of M. M. Gray; description of larva, and habits, ravages of and means against the larva and imago of Diabrotica vittata; figures larva, pupa, and imago of the same.
- 774. [WALSH, B. D., and C. V. RILEY.] Leafy oak gall. < Amer. Eut., September-October, 1869, v. 2, p. 25, fig. 20.

 Answer to inquiry of B. H. B.; description and figure of the gall of Cynips q.-frondosa.
- 775. [Walsh, B. D., and C. V. Riley.] Drop of gold. <Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of B. H. B.; egg of Citheronia regalis? on shellbark-hick-ory; characters of eggs and oviposition of Metapodius nasulus [= femoratus].
- 776. [WALSH, B. D., and C. V. RILEY.] The luna moth. <Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of G. W. Kinney; characters of Attacus [= Actias] luna; food-plants of the larva of the same.
- 777. [Walsh, B. D., and C. V. Riley.] Hag-moth larva. < Amer. Ent., September-October, 1869, v. 2, p. 25, fig. 21.
 - Answer to inquiries of C. T. Farrell and M. B. Baldwin; characters of cocoon and image of Limacodes [= Phobetron] pithecium; number of broods of the moth in the year; figure of the larva; occurrence of Harpacter [= Milyas] cinctus in Illinois.

- 778. [Walsh, B. D., and C. V. Rilley.] Stinging bug. <Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of J. M. Shaffer; habits and characters of Phymata erosa.
- 779. [Walsh, B. D., and C. V. Riley.] Pear-tree worms. < Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of B. Hathaway; larva of Notodonta [= Œdemasia] concinna found on pear-tree leaves.
- 780. [Walsh, B. D., and C. V. Riley.] "Dobson." <Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of Fisherman; ignorance as to what the larva called "Dobson" is.
- 781. [Walsh, B. D., and C. V. Riley.] White-pine weevil. < Amer. Ent., September-October, 1869. v. 2, p. 26, fig. 22.
 - Answer to inquiry of A. S. Fuller; seasons, ravages, and means against *Pissodes strobi*; figures larva, pupa, and imago of the same.
- 782. [Walsh, B. D., and C. V. Riley.] Unnatural secretion of wax. Amer. Ent., September-October, 1869, v. 2, p. 26.
 - Answer to inquiry of F. Brewer; description of a case of excessive secretion of wax by Apis mellifica.
- 783. [Walsh, B. D., and C. V. Riley.] Raspberry borer. < Amer. Ent., September-October, 1869, v. 2, p. 26.
 - Answer to inquiry of F. A. Gates; ravages of the larva of Oberea perspicillata [=bimaculata] in blackberry and raspberry bushes; characters of the imago of Dryocampa scnatoria; food-plants of the larva of the same.
- 784. [Walsh, B. D., and C. V. Riley.] Cocoon of horn-bug. < Amer. Ent., September-October, 1869, v. 2, p. 26.
 - Answer to inquiry of A. R. McClutchen; characters of cocoon of *Lucanus* dama?
- 785. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., September-October, 1869, v. 2, p. 26, fig. 23.
 - Answer to inquiry of J. R. Muhleman; characters and figure of Amphipyra [=Pyrophila] pyramidoides; food-plants and larva of the same and of A. pyramidea of Europe; characters of the larva and imago of Agnomonia anilis; supposed food-plants of its larva.
- 786. [Walsh, B. D., and C. V. Riley.] Cecropia moth caterpillar. <Amer. Ent., September-October, 1869, v. 2, p. 26.
 - Answer to inquiries of H. G. Lewelling and S. H. I. Green; characters and food-plants of the larva of Attacus cecropia.
- 787. [Walsh, B. D., and C. V. Riley.] How cut-worms originate. <Amer. Ent., September–October, 1869, v. 2, p. 26.
 - Answer to inquiry of T. W. Gordon; cut-worms are larvæ produced from eggs of certain *Noctuidæ*.
- 788. [Walsh, B.D., and C. V. Riley.] Red-humped caterpillar. < Amer. Ent., September-October, 1869, v. 2, p. 27, figs. 24-26.
 - Answer to inquiry of D. W. Kauffman; habits, characters, food-plants, and means against larva of *Notodonta* [= *Œdemasia*] *concinna*; figures larva, pupa, and imago of the same; poisonousness of the fluids of certain insects.

- 789. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., September-October, 1869, v. 2, p. 27.
 - Answer to inquiry of T. W. G[ordou]; characters and food-plants of *Thelia bimaculata* and of larva of *Procris* [= Harrisina] americana and Eudamus tityrus; characters of the image of the last.
- 790. [Walsh, B. D., and C. V. Riley.] Gilt gold-beetle. < Amer. Ent., September-October, 1869, v. 2, p. 27.
 - Answer to inquiry of W. H. Martin; characters and food-plants of Chrysochus auratus.
- 791. [Walsh, B. D., and C. V. Riley.] The trumpet grape-gall. <Amer. Ent., September-October, 1869, v. 2, p. 28, fig. 27.
 - Answer to inquiry of D. McClaine; description and figure of galls of Cecidomyia vitislituus [=viticola]; occurrence of similar galls on several varieties of grape-vines and on leaves of hickory and hackberry.
- 792. [Walsh, B. D., and C. V. Riley.] Grape-berry moth. < Amer. Ent., September-October, 1869, v. 2, p. 28.
 - Answer to inquiry of H. C. Barnard; ravages of Penthina vitivorana [= Eudemis botraua].
- 793. [Walsh, B. D., and C. V. Riley.] Oak pruner. < Amer. Ent., September-October, 1869, v. 2, p. 28.
 - Answer to inquiry of T. J. Plumb; occurrence of Elaphidion putator [=villosum] at Madison, Wis.
- 794. [Walsh, B. D., and C. V. Riley.] Potato-bugs. <Amer. Ent., September-October, 1869, v. 2, p. 28.
 - Answer to inquiry of W. R. Shelmire; ravages and food-plants of Lytta [= Epicauta] vittata; means against potato-eating Meloidæ; characters of an unknown lepidopterous larva boring in a potato-stalk; directions for packing insects.
- 795. [Walsh, B. D., and C. V. Riley.] Blood-sucking cone-nose. <Amer. Ent., September-October, 1849, v. 2, p. 28.
 - Answer to inquiry of G. W. C.; effect of the "bite" of Conorhinus sanguisuga; food-habits of the same.
- 796. [Walsh, B. D., and C. V. Riley.] Woolly slug-like worm on apple. <Amer. Ent., September-October, 1869, v. 2, p. 29.
 - Answer to inquiry of H. A. Green; food-plants and characters of larva of Lagoa opercularis; characters of the image of the same; improper method of packing living insects.
- 797. [Walsh, B. D., and C. V. Riley.] A water-bug. <Amer. Ent., September-October, 1869, v. 2, p. 29.
 - Answer to inquiry of W. V. Smith; characters of Ranatra fusca; habits and habitat of Nepidæ.
- 798. [Walsh, B. D., and C. V. Riley.] Goldenrod galls. < Amer. Ent., September-October, 1869, v. 2, p. 29.
 - Answer to inquiry of G. W. C.; characters of galls of Trypeta solidaginis and Cecidomyia solidaginis.

- 799. [Walsh, B. D., and C. V. Riley.] Oak-leaf gall. < Amer. Ent., September-October, 1869, v. 2, p. 29.
 - Answer to inquiry of B. H. Broadnox; description of galls of Cecidomyia quercus-pilulæ and C. q.-symmetrica; Cynipidæ inquilinous in galls of Cecidomyidæ; differences between larvæ of Cynipidæ and Cccidomyidæ; transformations of C. q.-pilulæ and of the Cynips sp., inquilinous in its gall; distinction between groups of oaks.
- 800. [Walsh, B. D., and C. V. Riley.] Humble bees. < Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiry of C. S. Davis; number of species and distribution of the genus *Bombus* in North America; habits of and differences between the several forms composing a society of social insects; habits of *Bombus pennsylvanicus*, *Halictus* sp., and *Andrena* sp.
- 801. [Walsh, B. D., and C. V. Riley.] Can land be insured against cut-worms and other insects? <Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiry of A. Willis; means against larve of cut-worms.
- 802. [Walsh, B.D., and C. V. Riley.] Beetles named. < Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiry of T. W. Hoyt, jr.; characters of Cassida [=Coptocycla] aurichalcea and Brachinus americanus; effect of the discharge made by Brachinus sp. upon the inside of the human mouth.
- 803. [Walsh, B. D., and C. V. Riley.] Royal horned-caterpillar. <Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiries of W. C. Holmes and M. G. Kern; larva of Citheronia regalis found feeding on Syringa.
- 804. [Walsh, B. D., and C. V. Riley.] Parsnip caterpillar. < Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiry of T. W. Hoyt, jr.; characters of larva of Papilio asterias.
- 805. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., September-October, 1869, v. 2, p. 31.
 - Answer to inquiry of C. P. Fanlkner; identification of several beetles; habits of Creophilus villosus, Listotrophus cingulatus, Scarites subterraneus, and Uloma impressa.
- 806. [Walsh, B. D., and C. V. Riley.] Beetle named. < Amer. Ent., September-October, 1869, v. 2, p. 31.
 - Answer to inquiry of W. Keyes; habits and characters of larva of Calopteron terminale; distinctness of C. reticulatum from C. terminale.
- 807. [Walsh, B. D., and C. V. Riley.] Moth named. < Amer. Ent., September-October, 1869, v. 2, p. 31.
 - Answer to inquiry of W. G. Barton; food-plant of larva and characters of imago of Alaria [= Rhodophora] florida.
- 808. [Walsh, B. D., and C. V. Riley.] Worm boring into cucumber.
 - Answer to inquiries of G. W. C., O. L. Barler, and E. S. Smith; characters of larva and imago of *Phakellura* [=Eudioptis] nitidalis; food-plants of larva; characters of undetermined larva found boring in cucumbers.

- 809. [Walsh, B. D., and C. V. Riley.] Caterpillar of the io moth. <a href="mailto: Amer. Ent., September-October, 1869, v. 2, p. 31.
 - Answer to inquiry of Mrs. Tildesley; characters of larva and image of Saturnia [= Hyperchiria] io; food-plant and urticating properties of the larva.
- 810. [Walsh, B. D., and C. V. Riley.] Apple-tree worms. < Amer. Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of H. K. Vickroy; habits and characters of larva of Acrobasis [= Pempelia] hammondi; habits of larva of Phycita nebulo [= Acrobasis indiginella].
- 811. [Walsh, B. D., and C. V. Riley.] Stinging larva. < Amer. Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of J. C. Falls; meaning of the vernacular and technical names of *Empretia stimulea*; urticating properties of lepidopterons larvæ.
- 812. [WALSH, B. D., and C. V. RILEY.] Lappet caterpillar on appletree. <Amer. Ent., September-October, 1869, v. 2, p. 32.

 Answer to inquiry of W. Stark; characters of larva of Gastropacha americana.
- 813. [WALSH, B. D., and C. V. RILEY.] Spined spider. <Amer. Ent., September-October, 1869, v. 2, p. 32.

 Answer to inquiries of G. W. Kinney and T. W. Gordon; characters and synonymy of Epeira [= Acrosoma] spinea.
- 814. [WALSH, B. D., and C. V. RILEY.] Dangerous looking. <Amer Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of M. M. Kenzic; characters and sting of Mutilla coccinea [= Sphærophthalma occidentalis].
- 815. [Walsh, B. D., and C. V. Riley.] Bag-worms again. < Amer. Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of T. C. Tipton; ravages of Thyridopteryx ephemeræformis; harmlessness of larva of Protoparce celeus; habitat of larva of Musca domestica.
- 816. [Walsh, B. D., and C. V. Riley.] Large water-beetle. < Amer. Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of S. E. Munford; secondary sexual characters in elytra of Cybister fimbriolatus. See No. 750.
- 817. [Walsh, B. D., and C. V. Riley.] Beetles under dead fish. Amer. Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of T. Ferrell; food-habits and characters of Silpha peltata [=americana].
- 818. [Walsh, B. D., and C. V. Riley.] Universal remedies. <Amer. Ent., November, 1869, v. 2, pp. 33-35.

 Worthlessness of any one substance as a means against insects.
- 819. [Walsh, B. D., and C. V. Riley.] Tent-caterpillars and fall-web-worms. < Amer. Ent., November, 1869, v. 2, p. 39.
 - Critical review of article in Western Rural, August 26, 1869; seasons, habits, food-plants, and characters of Clisiocampa americana and Hyphantria textor [=cunea].

- 820. [Walsh, B. D., and C. V. Riley.] The boll-worm or corn-worm. 2d article. (*Heliothis armigera*, Hübner.) <Amer. Ent., November, 1869, v. 2, pp. 42–44, fig. 29.
 - Seasous, food-plants, ravages of, and means against *Heliothis armigera*; figures of larvæ, pupa, cocoon, and imago of the same; food-plants of *Gortyna nitela*.
- 821. [Walsh, B. D., and C. V. Riley.] Galls and their architects. 2d article. <Amer. Ent., 1869–1870, v. 2: November, pp. 45–50, figs. 30–32; December–January, pp. 70–74, figs. 45–47; February, pp. 103–106, figs. 68–71.
 - See No. 518; definition and classification of galls; descriptions and figures of galls and larvæ of Nematus salicis-pomum, Euura s.-ovum, and E. s.-gemma [=orbitalis]; habits, scasons, and descriptions of the same; habits and seasons of Anthonomus sycophanta, Batrachedra salicipomonetla, and Nematus mendicus; differences between gall-makers and guest-flies; occurrence of distinct genera of gall-insects on plants of distinct genera; descriptions and figures of galls of Cynips [= Andricus] quercus-seminator and C. q.-frondosa; synoptic table of North American genera of Cynipidæ Psenides; description of Antistrophus n. g. and of A. lygodesmiæ-pisum n. sp. and its gall; description and figure of the larva and imago of Agrilus ruficollis and its gall; habits, food-plants of, and means against, the same; figure and description of Baridius [= Ampeloglypter] sesostris and its gall; habits, food-plants of, and means against, the same; characters of Madarus ampelopsidos [= Ampeloglypter ater] and its gall; habits of Buprestidæ.
- 822. [Walsh, B. D., and C. V. Riley.] Toads in gardens. < Amer. Ent., November, 1869, v. 2, p. 50.
 Value of toads as a means against noxious insects, etc., in gardens.
- 823. [Walsh, B. D., and C. V. Riley.] Notes on the Tarantula-killer. <Amer. Ent., November, 1869, v. 2, p. 52.
 - Comments on note of C. Peabody; occurrence of Mygale hentzii and Pepsis formosa in Missouri.
- 824. [Walsh, B. D., and C. V. Riley.] Swarms of lady-birds. < Amer. Ent., November, 1869, v. 2, p. 55.
 - Occurrence of countless millions of Coccinellidæ in England; their origin and movements.
- 825. [Walsh, B. D., and C. V. Riley.] The squash-bug does not touch the white bush scollop. <Amer. Ent., November, 1869, v. 2, p. 55.
 - Coreus [= Anasa] tristis does not attack the white bush scollop variety of the squash-vine; means against the same.
- 826. [Walsh, B. D., and C. V. Riley.] [Scientific names.] < Amer. Ent., November, 1869, v. 2, p. 57.

 Use of English and scientific names of insects.
- 827. [Walsh, B. D., and C. V. Riley.] On our table. < Amer. Ent., November, 1869, v. 2, pp. 57-58.
 - Notices of: Record of American entomology for the year 1868.—The Canadian entomologist.—The butterflies of North America, by W. H. Edwards.—Guide to the study of insects, by A. S. Packard, jr., etc.

- 828. [Walsh, B. D., and C. V. Riley.] Locust borer. <Amer. Ent., November, 1869, v. 2, p. 58.
 - Answer to inquiry of J. Bagby; $Arhopalus [= Cyllene] \ robiniw \ \ \ \$ undistinguishable from $A. [= C.] \ pictus \ \ \ \$.
- 829. [Walsh, B. D., and C. V. Riley.] Saddle-back larva. < Amer. Ent., November, 1869, v. 2, p. 59, fig. 36.
 - Answer to inquiry of G. T. Cost; figure of the larva of *Empretia stimulea* found on Indian corn.
- 830. [Walsh, B. D., and C. V. Riley.] Silk spiders. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiries of G. Howe and C. W. Spaulding; characters of *Epeira* [= Argiope] riparia and Nephila plumipes.
- 831. [Walsh, B. D., and C. V. Riley.] Entomological works. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of S. W. Cowles; mention of works containing descriptions of North American Colcoptera and Lepidoptera; food-plants of the larva of *Endryas unio*.
- 832. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of A. H. R. Bryant; characters of the cocoon of Attacus cecropia and the image of Mutilla coccinea [= Sphærophthalma occidentalis].
- 833. [Walsh, B. D., and C. V. Riley.] A new bee enemy. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of F. Brewer; characters of an undetermined carabid larva found eating Apis mellifica.
- 834. [Walsh, B. D., and C. V. Riley.] Thousand-legged worms. <Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of J. W. Merchant; characters and poisonousness of Scolopendra castaneiceps; Julus sp. and Polydesmus sp. injurious to strawberries.
- 835. [Walsh, B. D., and C. V. Riley.] Cabbage-worms. < Amer. Ent., November, 1869, v. 2, p. 60, figs. 37–38.
 - Answer to inquiry of W. C. Holmes; figures larva, pupa, and imago of *Pieris* protodice; means against cabbage-worms.
- 836. [Walsh, B. D., and C. V. Riley.] The rape butterfly. Amer. Ent., November, 1869, v. 2, p. 60.
 - Answer to inquiry of J. E. Chase; occurrence of Pieris rapæ in Bangor, Me.
- 837. [Walsh, B. D., and C. V. Riley.] Bad packing. < Amer. Ent., November, 1869, v. 2, p. 60.
 - Answer to inquiry of H. C. Beardslee; characters and food-plants of *Empretia stimulea*; food-plant of *Darapsa* [= Ampelophaga] myron; directions for sending larvæ by mail.
- 838. [Walsh, B.D., and C.V.Riley.] Granddaddy long-legs. < Amer. Ent., November, 1869, v. 2, p. 60.
 - Answer to inquiry of W. R. Howard; vernacular names and habits of *Phalangida*.
- 839. [Walsh, B. D., and C. V. Riley.] Borer in apple-twig. < Amer. Ent., November, 1869, v. 2, p. 60.
 - Answer to inquiry of G. C. Brackett; habits of Bostrichus [= Amphicerus] bicaudatus and Elaphidion parallelum [=villosum].

- 840. [Walsh, B.D., and C.V. Riley.] Grape-vine leaf-galls. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of W. T. Heildrup; habits and means against *Phylloxera* vastatrix; varieties of grape infested by the same.
- 841. [Walsh, B. D., and C. V. Riley.] Maple-worms. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of H. K. Vickroy; characters and ravages of *Dryocampa* rubicunda and *Acronycta americana*; food-plants of *Telea polyphemus* and *Attacus cecropia*.
- 842. [Walsh, B. D., and C. V. Riley.] Melancholy chafer in apples. <Amer. Ent., November, 1869, v. 2, p. 61, fig. 39.
 - Answer to inquiry of J. F. Fulton; figure of Euryomia [=Euphoria] melan-cholica found boring in apples.
- 843. [Walsh, B. D., and C. V. Riley.] Worms boring in cucumbers. <Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of W. B. Ramson; food-habits of *Phacellura* [= *Eudioptis*] nitidalis.
- 844. [Walsh, B. D., and C. V. Riley.] Lilac-borer. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of T. J. Freeman; characters, affinities, and means against Ægeria [= Podosesia] syringæ.
- 845. [Walsh, B. D., and C. V. Riley.] Burying beetles. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of J. H. Osborn; habits of Necrophorus marginatus and of Silphidæ generally; characters of Hylobius stupidus [= Pachylobius picivorus] found on plum-trees.
- 846. [Walsh, B. D., and C. V. Riley.] Wire-worms in potatoes. <Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of W. R. Shelmire; characters of larva (*Melanotus incertus?*) boring in potatoes; food-habits and means against larvæ of *Elateridæ*.
- 847. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of J. F. Waters; unknown tortricid and *Limacodes* sp., found on apple-tree; characters of *Chariesterus antennator*.
- 848. [Walsh, B. D., and C. V. Riley.] Girdled pear twigs. < Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of T. A. Thorp; pear twigs girdled by Oncideres cingulata.
- 849. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of E. T. Dale; food-habits of Hippodamia glacialis, Strachia [=Murgantia] histrionica and Blepharida rhois; figure of Hippodamia glacialis.
- 850. [Walsh, B. D., and C. V. Riley.] Tomato-feeding worm. < Amer. Ent., November, 1869, v. 2, p. 62, fig. 41.
 - Answer to inquiry of A. C. Davis; description, seasons, and food-plants of larva of *Prodenia commelina*; figure of dorsal surface of a segment of the same.

- 851. [Walsh, B. D., and C. V. Riley.] Cocoons of Ichneumon flies. <Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of C. Mitchell; cocoons of *Microgaster* sp. found on larva of *Protoparce celcus*.
- 852. [Walsh, B. D., and C. V. Riley.] Gall on spotted touch-me-not. <Amer. Ent., November, 1869, v. 2, p. 63, fig. 42.
 - Answer to inquiry of A. N. Prentiss; description and figures of the gall of Cecidomyia impatientis found on Impatiens fulva.
- 853. [Walsh, B. D., and C. V. Riley.] Unknown larvæ. < Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of J. M. Harold; characters of unknown larva allied to Lagoa.
- 854. [Walsh, B. D., and C. V. Riley.] Blood-sucking cone nose. <Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of D. B. Watson; occurrence of Conorhinus sanguisugus at Saint Louis, Mo.
- 855. [Walsh, B. D., and C. V. Riley.] Eggs on a grape-cane. < Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of J. Cochrane; difficulty of identifying eggs of insects.
- 856. [Walsh, B. D., and C. V. Riley.] Gregarious willow-worms. <Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of G. C. Brackett; characters, habits, and food-plants of the larva of Clostera americana [= Ichthyura inclusa].
- 857. [Walsh, B. D., and C. V. Riley.] Caterpillars named. < Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiries of G. W. Copley and W. D. Butler; food-habits of larvæ of *Eudamus tityrus* and *Papilio troilus*.
- 858. [WALSH, B. D., and C. V. RILEY.] Insects named. <Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of L. G. Saffer; vernacular name of Mutilla coccinca [= Spharophthalma occidentalis]; characters of Stizus [= Sphceius] speciosus.
- 859. [Walsh, B. D., and C. V. Riley.] Range of the rear-horse. <Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of V. T. Chambers; northern range of Mantis [= Phasmomantis] earolina.
- 860. [Walsh, B. D., and C. V. Riley.] Royal horned-caterpillar. <Amer. Ent., November, 1869, v. 2, p. 64.
 - Answer to inquiry of D. L. Phares; Citheronia regalis more abundant in the South than in the North.
- 861. [Walsh, B. D., and C. V. Riley.] Hair-worm or hair-snake. <Amer. Ent., November, 1869, v. 2, p. 64.
 - Answer to inquiry of E. H. King; Gordius sp. parasitic in the pupa of Orchelimum sp.
- 862. [Walsh, B. D., and C. V. Riley.] Dahlia-stalk borer. < Amer. Ent., November, 1869, v. 2, p. 64.
 - Answer to inquiry of G. C. Broadhead; larva of Gortyna nitela boring in stalk of dahlia.
- 863. [Walsh, B. D., and C. V. Riley.] Parsnip worm. < Amer. Ent., November, 1869, v. 2, p. 64.
 - Answer to inquiry of J. Adams; larva of Papilio asterias feeds on parsnip.



U. S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

 \mathbf{OF}

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

BY

SAMUEL HENSHAW.

PART III.


THE MORE IMPORTANT WRITINGS

 \mathbf{OF}

CHARLES VALENTINE RILEY.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1889.

7 ENT



BIBLIOGRAPHY OF THE MORE IMPORTANT CONTRIBUTIONS TO AMERICAN ECONOMIC ENTOMOLOGY.

PART III.

- 864. R[ILEY], C. [V.] The house-fly. < Prairie Farmer, 2 May, 1863, [v. 27], n. s., v. 11, pp. 276-277. S.-b. No. 1, p. 9.
 - Stomoxys calcitrans distinguished from Musca domestica; habits of each; transformations of the latter; habit of larvæ of collecting underneath boards.
- 865. R[ILEY], C. [V.] The May-beetle. < Prairie Farmer, 6 June, 1863, [v. 27], n. s., v. 11, p. 356. S.-b. No. 1, p. 7.
 - Ravages of Phyllophaga quercina [= Lachnosterna fusca]; description and habits of larvæ and imagos; remedies.
- \$66. R[ILEY], C. V. Larvæ of the ten-striped spearman. < Prairie Farmer, 8 August, 1863, [v. 28], n. s., v. 12, pp. 85-86, fig. S.-b. No. 1, p. 11.
 - Description of eggs; figure and first description of larva of *Doryphora* 10-lineata; habits of larva; seasons; remedies; enemies of larva.
- 867. R[ILEY], C. V. The squash-bug. <Prairie Farmer, 8 August, 1863, [v. 28], n. s., v. 12, p. 86. S.-b. No. 1, p. 11.
 - Description of images and larvæ of Coreus [= Anasa] tristis; habits, seasons, remedies.
- 868. RILEY, C. V. The cut-worm. < Prairie Farmer, 12 March, 1864, [v. 29], n. s., v. 13, p. 169.
 - Answer to communication of G. R. Huffman; habits and transformations of Agrotididæ; means against them.
- 869. [RILEY, C. V.] Entomological. <Prairie Farmer, 21 May, 1864, [v. 29], n. s., v. 13, p. 361.
 - Answer to communication of J. S. Lawver; ravages of unknown caterpillars, probably cut-worms, and of *Ithycerus noveboracensis* on young appletrees; means against these insects.
- 870. [RILEY, C. V.] Apple-borers. < Prairie Farmer, 14 January, 1865, [v. 31], n. s., v. 15, p. 21, 5 figs. S.-b. No. 1, p. 30.
 - Description and figure of larva and image of Buprestis [= Chrysobothris] femorata; injury done by the larva; remedies. Figure of image of Saperda bivittata [= candida].

- 871. [RILEY, C. V.] Peach-tree borers. < Prairie Farmer, 25 February, 1865, [v. 31], n. s., v. 15, pp. 122–123, 6 figs. S.-b. No. 1, pp. 30–31.
 - Descriptions and figures of larva, pupa, cocoon, and ∂ and Q imagos of Egeria[=Sannina] exitiosa; description of egg; seasons and habits of larva and imago; remedics. Figure of imago of Dicerca divarienta.
- 872. [RILEY, C. V.] Entomology. < Prairie Farmer, 22 April, 1865, [v. 31], n. s., v. 15, p. 306. S.-b. No. 1, pp. 32-33.
 - Coming forth of insects in spring; desirability of observing noxious insects and of recording observations; abundance of eggs of *Orgyia* this season; methods of destroying them.
- 873. [RILEY, C. V.] Flea-beetles and Curculio. <Prairie Farmer, 27 May, 1865, [v. 31], n. s., v. 15, p. 418, fig. S.-b. No. 1, p. 33.

 Prevalence of and means against *Halticide*; habits of and means against

Conotrachelus nenuphar; figure of pupa.

- 874. [RILEY, C. V.] Apple-tree caterpillars. < Prairie Farmer, 3 June, 1865, [v. 31], n. s., v. 15, pp. 437–438. S.-b. No. 1, p. 34.

 Answer to communication from J. C. Brown; descriptions of larva, pupa, and imago, of Chatochilus pometellus; habits of the same; other caterpillars injurious to the leaves of apple trees and means against them.
- 875. [RILEY, C. V.] Curculio catcher. <Prairie Farmer, 10 June, 1865 [v. 31], n. s., v. 15, p. 457, fig.
 Figure of E. S. Hull's Curculio catcher, with explanatory text.
- 876. [RILEY, C. V.] The army-worm. < Prairie Farmer, 8 July, 1865, [v. 32], n. s., v. 16, pp. 3-4, 3 figs. S.-b. No. 1, pp. 36-37.

 Answer to communication from W. R.; descriptions and figures of larva, pupa, and imago of Leucania unipuncta; number of broods unknown; remedies.
- 877. [RILEY, C. V.] The current-worm. < Prairie Farmer, 15 July, 1865, [v. 32], n. s., v. 16, p. 27, 4 figs. S.-b. No. 1, p. 36.

 Description of eggs; descriptions and figures of larva, pupa, and image of Euftchia ribearia; habits, seasons, remedies.
- 878. [RILEY, C. V.] Swallows. <Prairie Farmer, 15 July, 1865, [v. 32], n. s., v. 16, p. 27. S.-b. No. 1, p. 36.
 Usefulness of swallows in the destruction of noxions insects.
- 880. [RILEY, C. V.] Singular caterpillar. <Prairie Farmer, 22 July, 1865, [v. 32], n. s., v. 16, p. 50. S.-b. No. 1, p. 35.

 Descriptions of larva, pupa, and image of Notodonta [= Calodasys] unicornis; habits; food-plants.
- 881. [RILEY, C. V.] Collecting and preserving insects. < Prairie Farmer, 5 August, 1865, [v. 32], n. s., v. 16, pp. 101–102, fig.

 Interest and importance of entomology; directions for the construction of nets and cabinets, and for the capture, killing, preparation, and preservation of insects.

- 822. RILEY, C. V. The grass-bug and its habits: Currant-worms. <Cultivator and Country Gentleman, 10 August, 1865, v. 26, p. 98. Reprint: <Boston Cultivator, 19 August, 1865, v. 37, p. 259. S.-b. No. 1, p. 60.
 - Habits of Cercopidida; nature of the image of eurrant-worms.
- 883. [RILEY, C. V.] Apple plant-louse. <Prairie Farmer, 19 August, 1865, [v. 32], n. s., v. 16, p. 127. S.-b. No. 1, p. 40.

 Answer to inquiry of J. Doron; habits and fecundity of Aphis mali; means against it.
- 884. R[ILEY], C. [V.] Seventeen-year locust. < Prairie Farmer, 19 August, 1865, [v. 32.], n. s., v. 16, p. 127. S.-b. No. 1, pp. 40-41. Agrees with S. P. G. in doubting that Cicada [= Tibicen] septendecim lives seventeen years immature, and gives reasons for his doubt.
- 885. [RILEY, C. V.] Tobacco-worm. < Prairie Farmer, 2 September, 1865, [v. 32], n. s., v. 16, p. 165. S.-b. No. 1, p. 41.

 Answer to inquiry of A. B. Knowlton; habits of Macrosila quinquemaculata [= Protoparce celeus].
- 887. RILEY, C. V. The sheep gad-fly. <Prairie Farmer, 14 October, 1865, [v. 32], n. s., v. 16, pp. 288–289, figs. 1–6. S.-b. No. 1, pp. 48–49.
 - Answer to inquiries of a subscriber; figures and descriptions of larva, pupaease, and imago of Cephalemyia [= Estrus] ovis; habits of and means against it; imago viviparous.
- 888. [RILEY, C. V.] Chinch-bug not in seed grain. <Prairie Farmer, 21 October, 1865, [v. 32], n. s., v. 16, p. 308. S.-b. No. 1, p. 4. Supposed evidence that the eggs of Blissus leucopterus are not deposited and do not winter in seed wheat.
- 890. [RILEY, C. V.] The chinch-bug once more. A reply to D. H. Sherman. <Waukegan [Ill.] Gazette, 20 January, 1866, v. 16, No. 18, p. 4.
 - Critical review of article of D. H. Sherman.
- 891. RILEY, C. V. The chinch-bug. <Prairie Farmer, 3 March, 1866, [v. 33], n. s., v. 17, p. 133. S.-b. No. 1, p. 71.

 Answer to inquiry of W. R. Everett; means against Blissus leucopterus; habits, hibernation, and oviposition.
- 892. [RILEY, C. V.] Wire-worms. < Prairie Farmer, 3 March, 1866, [v. 33], n. s., v. 17, p. 133, figs. 4-5. S.-b. No. 1, p. 71.

 Answer to inquiry of Creswell; means against larve of Elateridæ.

- 893. R[ILEY], C. V. Entomological. < Prairie Farmer, 24 March, 1866, [v. 33], n. s., v. 17, p. 192, figs. 5-6. S.-b. No. 1, p. 68.
 - Answers to inquiries of E. T. Nelson, J. Doron, and H. W. Shore; habits and figures of *Amphicerus bicaudatus*; remedies. Apple-twigs picrced by one of the *Uroceridæ*.
- 894. RILEY, C. V. The chinch-bug. < Pract. Ent., 26 March, 1866, v. 1, pp. 47-48.
 - Criticism of D. H. Sherman's communication on this insect; Blissus leucopterus does not oviposit in the fuzzy end of wheat-grain; brief life-history.
- 895. RILEY, C. V. The chinch-bug once more. <Waukegan [Ill.] Gazette, 31 March, 1866, v. 16, No. 28, p. 4. S.-b. No. 1, p. 70.
 - Reply to D. H. Sherman's second communication; Blissus leucopterus does not deposit its eggs in the grain of wheat; means against it.
- 896. [RILEY, C. V.] Bark-lice remedy. < Prairie Farmer, 7 April, 1866, [v. 33], n. s., v. 17, p. 229. S.-b. No. 1, p. 79.
 - Answers to inquiries of a subscriber and of S. J. B.; means against *Mytilas-* pis pomicorticis [= pomorum].
- 897. [RILEY, C. V.] "Bug" on melon, etc. < Prairie Farmer, 7 April, 1866, [v. 33], n. s., v. 17, p. 229. S. b. No. 1, p. 79.

 Answer to inquiry of M. E. W.; means against Diabrotica vittata.
- 898. [RILEY, C. V.] Warbles. < Prairie Farmer, 21 April, 1866, [v. 33], n. s., v. 17, p. 276.
 - The occurrence of lumps on the back of cattle caused by larvæ of *Hypoderma* bovis is not unusual or alarming.
- 899. RILEY, C. V. The ailanthus silk-worm. < Prairie Farmer, 28
 April, 1866, [v. 33], n. s., v. 17, p. 289, fig. S.-b. No. 1, p. 93.
 Partial reprint: < New York Tribune, 29 May, 1866, v. 26, No. 7844, p. 7. S.-b. No. 1, p. 92.
 - Describes attempts to introduce Bombyx arrindia and Samia [=Attacus] cynthia into Europe for commercial purposes. Describes and figures egg, larva, and imago, and figures pupa of Samia [= Attacus] cynthia; habits, seasons, and hardiness, and disadvantages for purposes of silk culture of this species; superiority of native species.
- 900. [RILEY, C. V.] Lice on pigs. < Prairie Farmer, 28 April, 1866, [v. 33], n. s., v. 17, p. 292. Means against.
- 901. RILEY, C. V. Novel facts about cut-worms. < Prairie Farmer, 2 June, 1866, [v. 33], n. s., v. 17, pp. 371, 372, figs. 3–8. S.-b. No. 1, p. 89.
 - * Climbing habits of the larvæ of Agrotididæ; figures and descriptions of the larvæ of Agrotis cochranii [= A. messoria], A. clandestina, and ? Hadena subjuncta; enemies of cut-worms.
- 902. [RILEY, C. V.] The canker-worm. < Prairie Farmer, 16 June, 1866, [v. 33], n. s., v. 17, p. 412. S.-b. No. 1, p. 96.

 Answer to inquiry of a subscriber; means against *Anisopteryx*.
- 903. R[ILEY], C. V. Large fish-fly. < Prairie Farmer, 16 June, 1866, [v. 33], n. s., v. 17, p. 412. S. b. No. 1, p. 96.

Answer to inquiry of A. B. Knowlton; habits and distribution of Perla dorsata.

- 904. RILEY, C. V. The bee-moth. < Cultivator and Country Gentleman, 21 June, 1866, v. 27, p. 399. S.-b. No. 1, p. 97.

 Answer to inquiry of E. S. Fowler; habits of Galleria cercana.
- 905. RILEY, C. V. The potato-bug. <Prairie Farmer, 23 June, 1866, [v. 33], n. s., v. 17, p. 432. S.-b. No. 1, p. 97.

 Auswer to inquiry of J. D. Eilington; eastward spreading of and means
- against Doryphora decemlineata.

 906. [RILEY, C. V.] Army-worm. < Prairie Farmer, 23 June, 1866, [v. 33], n. s., v. 17, p. 432. S.-b. No. 1, p. 97.

Occurrence of Leucania unipuncta in various parts of Illinois.

- 907. RILEY, C. V. White willow insects. < Prairie Farmer, 30 June, 1866, [v. 33], n. s., v. 17, p. 452. S.-b. No. 1, p. 101.
 - Answer to inquiry of L. G. H.; descriptions of larva, pupa, and image of Vanessa antiopa; description of larva of Nematus ventralis; habits of and means against these.
- 908. RILEY, C. V. Still they come. < Prairie Farmer, 30 June, 1866, [v. 33], n. s., v. 17, p. 452. S.-b. No. 1, p. 101.

 Answer to inquiry of S. D.; means against Doryphora decemlineata.
- 909. RILEY, C. V. The worm question. < Ohio Farmer, 7 July, 1866, v. 15, p. 209. S.-b. No. 2, p. 17.
 - Comments on a controversy between J. K. and J. Brocket concerning *Pyrrharctia isabella*; molting, pupation, and cocoon-making of this and other caterpillars.
- 910. RILEY, C. V. Army-worm and canker-worm wisdom. < Prairie Farmer, 21 July, 1866, [v. 34], n. s., v. 18, p. 38. S.-b. No. 1, p. 101; No. 2, p. 40.
 - Comments on entomological ignorance amongst members of the Fruit Grower's Society of Western New York; efficacy of means against Anisopteryx.
- 911. R[ILEY], C. V. [Attacus cecropia.] < Prairie Farmer, 21 July, 1866, [v.34], n. s., v. 18, p. 40. S.-b. No. 1, p. 104; No. 2, p. 40. Answer to inquiry of F. T.; habits of Attacus eccropia; its nsefulness for silk.
- 912. R[ILEY], C. V. [Cantharis cinerea.] < Prairie Farmer, 21 July, 1866, [v. 34], n. s., v. 18, p. 40. S.-b. No. 1, p. 104; No. 2, p. 40. Answer to inquiry of W. D. Hoord; criticizes the figure of Lytta fabricii [=Macrobasis unicolor] in Harris' Insects; usefulness of the Meloidæ as vesicants; habits of and means against M. unicolor.
- 913. RILEY, C. V. [*Edema albifrons.*] < Prairie Farmer, 1 August 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

Answer to inquiry of H. Kennedy; probability of unknown notodontoid proving *Edema albifrons*.

- 914. RILEY, C. V. Practical entomology in reality. < Maine Farmer, 2 August, 1866. S.-b. No. 2, p. 21.
 - Critical review of Brackett's Practical Entomology No. 9; some eut-worms do climb trees; Estrus ovis is viviparous.
- 915. RILEY, C. V. [Clytus speciosus.] < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.
 - Answer to inquiry of C. L. Edwards; habits of and means against Clytus [= Plagionotus] speciosus injuring maple-trees; description of larva of Anisota [= Dryocampa] rubicunda which feeds upon maple leaves.

- 916. RILEY, C. V. Grape-leaf louse. < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

 Answer to inquiries of "Subscriber"; description, habits and development of Pemphigus vitifoliæ [= Phylloxera vastatrix].
- 917. RILEY, C. V. White willow worm. < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

 Descriptions of larva and imago of Nematus ventralis; food-plants; ravages and number of broods.
- 918. RILEY, C. V. Insects in timber. < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110. Unknown species of *Tortricidæ* injurions to forests in Brown and Schuyler Counties, Illinois, in 1865.
- 919. RILEY, C. V. Der neue Kartoffel-Käfer. < Deutsche Prairie Farmer, August, 1866, 2 figs. S.-b. No. 2, p. 14.

 Descriptions and figures of larva and image of *Doryphora decemlineata*; its eastern migrations; habits; means against it.
- 920. RILEY, C.'V. Cicadas and walking-sticks. < Prairie Farmer, 1
 September, 1866, [v. 34], n. s., v. 18, p. 136. S.-b. No. 2, p. 41.
 Answer to inquiry of J. D. Swain; condensed account of Cicada canicularis
 [=tibicen]; comparison with C. [= Tibicen] septendecim; mentions Spectrum [= Diapheromera] femoratum.
- 921. [RILEY, C. V.] Locust-borer. <Prairie Farmer, 1 September, 1866, [v. 34], n. s., v. 18, p. 140.

 Answer to inquiry of O. H. S.; nearly all of the locust groves in the West have been destroyed by Clytus [= Cyllene] robiniæ; no remedy known.
- 922. RILEY, C. V. Apple-tree caterpillars. <Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152. S.-b. No. 2, p. 41.

 Answer to inquiries of S. Caverno and H. M. Lyman; habits, descriptions of larvæ, and images of Datana ministra and Notodonta [= Œdemasia] concinna; means against them; description of pupa of Datana ministra.
- 923. RILEY, C. V. Joint-worm. < Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152. S.-b. No. 2, p. 41.

 Answer to inquiry of G. W. Conklin; habits of and means against Isosoma
- 924. RILEY, C. V. [Nematus ventralis.] < Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152.

 Answer to inquiry of G. E. W.; ravages of Nematus ventralis in Kalmar, Minn.

hordei.

- 925. [RILEY, C. V.] Ten-lined potato-beetle. <Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152. S.-b. No. 2, p. 42.

 Larva of Lema trilineata on potato plants in Maine mistaken for that of Doryphora decemlineata.
- 926. [RILEY, C. V.] Locusts. < Prairie Farmer, 3 November, 1866, [v. 34], n. s., v. 18, p. 290.

 Occurrence and ravages of Calopterius spretus in portions of Kansas and west-

ern United States in 1866.

927. RILEY, C. V. Caterpillars on the pine. <Prairie Farmer, 10 November, 1866, [v. 34], n. s., v. 18, p. 301. S.-b. No. 2, p. 42.

Answer to inquiry of N. R. Strong and A. L. Jones; detailed description of larva of Lophyrus abbotii; habits of larva; means against the insect.

928. RILEY, C. V. Elm- and pear-tree borer. <Prairie Farmer, 10 November, 1866, [v. 34], n. s., v. 18, p. 301. S.-b. No. 2, p. 42.

Answer to inquiry of James Matteson; description and habits of *Tremex*

columba.

- 929. RILEY, C. V. Grasshoppers and locusts. < Prairie Farmer, 24 November, 1866, [v. 34], n. s., v. 18, p. 333. S.-b. No. 2, pp. 43–44.
 - Answer to inquiry of J. N.; distinguishes grasshoppers from locusts; characterizes Achetadæ [= Gryllidæ], Gryllidæ [= Locustidæ], and Locustidæ [= Acrididæ]; oviposition, transformations, ravages, and migrations of Acrididæ; recognition of Caloptenus spretus.
- 930. RILEY, C. V. Black-knot once more. Sardeners Mo. and Hortic., November, 1866, v. 8, pp. 331–332. S.-b. No. 2, p. 31. Fungoid origin of black-knot [Sphæria morbosa]; larvæ of Conotrachelus nenuphar present in the swellings.
- 931. R[ILEY], C. V. Brimstone for borers. <Prairie Farmer, 8 December, 1866, [v. 34], n. s., v. 18, p. 365. S.-b. No. 2, p. 44. Criticism of a communication by W.; sulphur inserted in trunks of trees ineffectual as a means against insects.
- 932. RILEY, C. V. The wire-worm. <Cultivator and Country Gentleman, 27 December, 1866, v. 28, p. 414. S.-b. No. 2, p. 34. Critical review of J. D. Gros' "The wire-worm;" myriapods and the larvæ of *Elateridæ* are called wire-worms; separable by the number of their legs; means against them.
- 933. [RILEY, C. V.] Tilden tomato and the tobacco-worm. < Prairie Farmer, 5 January, 1867, [v. 35], n. s., v. 19, p. 5.

 Remarks on communication of T. B.; tomato plants eaten by tobacco-worms [Protoparce carolina] in preference to tobacco plants; possibly the worms observed were tomato-worms [P. celeus].
- 934. [RILEY, C. V.] Remarks on Saperda, Chrysobothris, Carpocapsa, and Conotrachelus. < Prairie Farmer, 12 January, 1867, [v. 35], n. s., v. 19, p. 23.
 - Report of remarks made at the annual meeting of Southern Illinois Fruit Growers' Association.
- 935. [RILEY, C. V.] Bark-lice. <Prairie Farmer, 12 January, 1867, [v. 35], n. s., v. 19, p. 24.
 Washing apple-trees with lye effective against bark-lice.
- 936. RILEY, C. V. Insects in the flower garden. A troublous time. <Prairie Farmer, 19 January, 1867, [v. 35], n. s., v. 19, p. 37. S.-b. No. 2, p. 45.
 - Answer to communication by Kate Sherman; describes larva of [Heliothis phlogophagus] injuring Phlox.
- 937. [RILEY, C. V.] Salt and vinegar for insects. <Prairie Farmer, 19 January, 1867, [v. 35], n. s., v. 19, p. 37. S.-b. No. 2, p. 45. Review of T. Glover's Report of the U. S. Entomologist for 1865, T. Glover's Entomological exhibition in Paris, and W. C. Lodge's Fruits and fruit trees of the Middle States; use of salt and vinegar as means against insects; criticism of errors. See Nos. 222, 939, 942.

- 938. RILEY, C. V. Hickory bark borer, *Scolytus caryæ*, n. sp. < Prairie Farmer, 2 February, 1867, [v. 35], n. s., v. 19, pp. 68, 69, 6 figs. S.-b. No. 2, p. 50.
 - Descriptions and figures of larva and image of Scolytus caryæn. sp. [=4-spinosus] and of its borings in bark and wood of Carya; habits and ravages; figure of the pupa of S. destructor of Europe.
- 939. [RILEY, C. V.] Correction. < Prairie Farmer, 2 February, 1867, [v. 35], n. s., v. 19, p. 69. S.-b. No. 2, p. 50.
 - Correction of statement made in No. 937; T. Glover not responsible for calling Saperda bivittata [=candida] a butterfly. See Nos. 222, 937, 942.
- 940. RILEY, C. V. Dahlia and aster stalk-borer, Gortyna nitela Guenée. <Prairie Farmer, 23 February, 1867, [v. 35], n. s., v. 19, p. 116, 2 figs. S.-b. No. 2, p. 58.
 - Description of larva, pupa, and image of Gortyna nitela; habits of and means against this insect.
- 941. R[ILEY], C. V. Borers and canker-worms. < Prairie Farmer, 9 March, 1867, [v. 35], n. s., v. 19, p. 151. S. b. No. 2, p. 61.
 - Critical review of E. H. C.'s "Remedy for the borer" and of J. Huggins' "Canker-worm;" indefiniteness of the word borer; borers confined to particular trees according to species; relations between Clytus [= Cyllene] pictus and C. [= C.] robinia; habits of and means against canker-worms. See No. 954.
- 942. [RILEY, C. V.] The critic criticised. < Prairie Farmer, 16 March, 1867, [v. 35], n. s., v. 19, p. 169. S.-b. No. 2, p. 58.
 - Reply to B. D. Walsh's "The critic criticised;" insists upon the correctness of one criticism made; acknowledges incorrectness of another. See Nos. 222, 937, 939.
- 943. RILEY, C. V. [Solenobia.] < Prairie Farmer, 16 March, 1867, [v. 35], n. s., v. 19, p. 169. S.-b. No. 2, p. 58.
 - Answer to inquiry of J. C. Plumb; brief description of the larva of ? Solenobia; habits of the genus.
- 944. RILEY, C. V. Bark-lice. Their history, together with sundry remedies. <Prairie Farmer, 23 March, 1867, [v. 35], n. s., v. 19, p. 184. S.-b. No. 2, pp. 61–62. Extract: <Cultivator and Country Gentleman, 23 May, 1867, v. 29, p. 334. <Pract. Ent., April, 1867, v. 2, pp. 81–82.
 - Criticises several patent remedies for ravages of bark-lice, Coccidæ; ignorance of entomology amongst intelligent writers; natural history of Mytilaspis pomicorticis [= pomorum].
- 945. RILEY, C.V. The phlox-worm. < Prairie Farmer, 6 April, 1867, [v. 35], n. s., v. 19, p. 219, 2 figs. S.-b. No. 2, pp. 63, 64.
 - Describes and figures larva and image of *Heliothis phloxiphaga*; describes pupa; seasons and habits.
- 946. RILEY, C. V. The potato-beetle. < Prairie Farmer, 6 April, 1867, [v. 35], n. s., v. 19, p. 219. S.-b. No. 2, p. 64.
 - Answer to inquiries of H. Tilden; Doryphora decemlineata remains permanently in regions invaded by it; hibernates as an imago under ground.

947. RILEY, C. V. Meadow-worms. < Prairie Farmer, 6 April, 1867, [v. 35], n. s., v. 19, p. 219. S.-b. No. 2, p. 64.

Answer to inquiry of B. S.; habits of Tipula in all stages.

- 948. RILEY, C. V. Clover-worms. < Prairie Farmer, 1867, [v. 35], n. s., v. 19, 20 April, pp. 260–261, 10 figs.; 27 April, p. 279. S.-b. No. 2, pp. 67, 68.
 - Habits and ravages of *Pyralis olinalis* [= Asopia costalis]; figures and detailed descriptions of larva, pupa, cocoon, and imago.
- 949. RILEY, C. V. The apple leaf crumpler. <Prairie Farmer, 27 April, 1867, [v. 35], n. s., v. 19, p. 279, 4 figs. S.-b. No. 2, p. 68. Answer to inquiry of A. H.; description of pupa, figure of larva-case and imago of Phycita nebulo [= Acrobasis indiginella]; habits and means against the same.
- 950. RILEY, C. V. Cocoons on the flowering ash. <Prairie Farmer, 27 April, 1867, [v. 35], n. s., v. 19, p. 279. S.-b. No. 2, p. 68. Answer to inquiry of A. B. Pierce; descriptions of larva, cocoon, and imago of Attacus promethea; food-plants of larva; method of emergence from cocoon.
- 951. RILEY, C. V. Scarred apple-trees. Prairie Farmer, 27 April, 1867, [v.35], n. s., v. 19, p. 279. S.-b. No. 2, p. 68.

 Answer to inquiry of P. M. Williamson; descriptions of the injuries to apple-trees caused by Tettigonia sp., Coccus harrisii [=Chionaspis furfurus], and Mytilaspis pomicorticis [=pomorum].
- 952. RILEY, C. V. Aphides. < Prairie Farmer, 18 May, 1867, [v. 35,], n. s., v. 19, p. 332. S.-b. No. 2, p. 73.

 Answer to inquiry of J. Taylor; color and propagation of Aphis mali; means against the same.
- 953. RILEY, C. V. Tree-cricket. Prairie Farmer, 18 May, 1867, [v. 35], n. s., v. 19, p. 332. S.-b. No. 2, p. 73.

 Answer to inquiry of A. N. Prentiss; habits of *Œcauthus niveus*.
- 954. RILEY, C. V. Note. < Prairie Farmer, 18 May, 1867, [v. 35], n. s., v. 19, p. 332. S.-b. No. 2, p. 73.

 Correction of No. 941; criticism unfounded.
- 955. RILEY, C. V. The strawberry-worm, *Emphytus maculatus* Norton. < Prairie Farmer, 25 May, 1867, [v. 35], n. s., v. 19, p. 348, 9 figs. S.-b. No. 2, p. 72.
 - Descriptions and figures of all stages, geographical distribution, seasons, habits, and oviposition of and means against *Emphytus* [=Harpiphorus] maculatus.
- 956. RILEY, C. V. White-pine worm: *Lophyrus abbotii*. < Prairie Farmer, 25 May, 1867, [v. 35], n. s., v. 19, p. 348, 7 figs. S.-b. No. 2. p. 72.
 - Description of image of *Lophyrus abbotii*; figures larva, pupa, cocoon, ♀ image and ♂ and ♀ antennæ; habits and scasons of larvæ.
- 957. [RILEY, C. V.] Stag-beetle. < Prairie Farmer, 25 May, 1867, [v. 35], n. s., v. 19, p. 348. S.-b. No. 2, p. 72.

 Answer to inquiry of S. Barrier; habits of Lucanus elaphus.

- 958. [RILEY, C. V.] The Curculio. < Prairie Farmer, 1 June, 1867, [v. 35], n. s., v. 19, p. 368. S.-b. No. 2, p. 73.
 - Successful results from the use of machines for destroying Conotrachelus nenuphar; this insect hibernates in the image state.
- 959. RILEY, C. V. Fifteen-spotted lady-bird. <Prairie Farmer, 8 June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Answer to inquiry of A. B. Knowlton; food, variable coloration of imago, and usefulness of Mysia [Anatis] 15-punctata.
- 960. RILEY, C. V. Cherry Aphis. < Prairie Farmer, 8 June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Answer to inquiry of G. Lee; habits, ravages of and means against Myzus cerasi.
- 961. RILEY, C. V. Tree-cricket. < Prairie Farmer, 8 June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Eggs of *Œcanthus nivens* deposited in raspberry canes cause the death of the wood above them.
- 962. RILEY, C. V. Apple-tree borer. < Prairie Farmer, 8 June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Answer to inquiry of A. B. Campbell; commends D. B. Wier's remedy against the apple-tree borer; Saperda bivittata [= candida] goes through its transformations in two years, though commonly believed to require three years.
- 963. RILEY, C. V. Insects affecting apple-tree roots. < Prairie Farmer, 15 June, 1867, [v. 35], n. s., v. 19, p. 397. S.-b. No. 2, p. 100.
 - Letter from O. B. Galusha, with answer; descriptions of young and mature wingless individuals of the root-inhabiting form, Pemphigus pyri, of Schizoneura lanigera; habits, seasons, and reproduction of this form; its gallmaking and means against it; description of larva of Cecidomyia species found with the lice; and of larva and imago of Mycetophila persica [n. sp.?]; seasons of the latter; larva of Saperda calcarata bores in roots of apple; larva of Helops micans feeds on decaying apples; Julus [Spirobolus] marginatus infested with Gamasus juloides.
- 964. RILEY, C. V. A chapter on cut-worms. <Prairie Farmer, 22 June, 1867, [v. 35], n. s., v. 19, pp. 413-414, 7 figs. S.-b. No. 2, pp. 79-80.
 - Extract from J. Townley's "Do cut-worms destroy tree buds?". Buds of fruit trees destroyed by the larvæ of Agrotididæ; other ravages and means against the same; descriptions of the larvæ of Agrotis subgothica [= A. herilis and A. tricosa], A. telifera [= A. ypsilon], and Celæna [= Hadena] renigera; description of A. cochranis n. sp. [= A. messoria]; figures larva and imago of A. cochranis, A. telifera, and Celæna renigera; and imago of A. subgothica; habits, seasons, and vernacular names of Agrotididæ.
- 965. [RILEY, C. V.] Strawberry-worm. < Prairie Farmer, 22 June, 1867, [v. 35], n. s., v. 19, p. 414. S.-b. No. 2, p. 80.
 - Answer to inquiry of A. R. Whitney; geographical distribution of *Emphytus* [= Harpiphorus] maculatus.
- 966. RILEY, C. V. Insects stripping the bur-oak. <Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, p. 81. Answer to inquiry of A. B. Price; habits of Lachnosterna pilosicollis [= tristis.]

- 967. RILEY, C. V. Borers. < Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, p. 81.
 - Answer to inquiry of J. Wentworth; means against Clytus [== Cyllene] robiniw and Xylcutes [= Cossus] robiniw.
- 968. RILEY, C. V. Potato-beetle. < Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, pp. 81-82.
 - Answer to inquiries of J. L. W. and G. L. Merriwether; Doryphora 10-lineata three-brooded; its occurrence at Shipman, Ill.
- 969. [RILEY, C. V.] Smith's patent Curculio trap. < Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, p. 82. Communication from N. C. Coffman, with reply; condemns J. Smith's Curculio trap.
- 970. RILEY, C. V. [Aphis ribis.] < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81. Answer to inquiry of C. J. Eckhart; means against Myzus ribis.
- 971. RILEY, C. V. Mantis carolina. < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81. Answer to communication of J. H. Graves; eggs of Mantis [= Phasmomantis] carolina found in Ogle County, Ill.
- 972. RILEY, C. V. Lappet caterpillars on the apple. < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81. Answer to communication of Perkins and Congdon; description of the larva. pupa, cocoon, and imago of Gastropacha [== Tolype] velleda.
- 973. RILEY, C. V. Apple-bark lice on pears. < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81. Mytilaspis pomicorticis [= pomorum] on the fruit of pcar-trees; said to occur on currants, plum, and black oak.
- 974. RILEY, C. V. Currant bush borer. < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81. Answer to inquiry of H. B. Kinne; habits, description, and means against · Egevia tipuliformis.
- 975. RILEY, C. V. Apple-leaf crumpler. <Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S. b. No. 2, p. 81. Answer to inquiry of H. C. Clock; habits and means against Phycita uebulo [=Acrobasis indiginella].
- 976. RILEY, C. V. Tomato-stalk borer. < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81. Answer to inquiry of O. G. Nevins; food-plants of Gortyna nitela.
- 977. RILEY, C. V. A few errors corrected. < Wisc. Farmer, 17 August, 1867. S.-b. No. 2, p. 83. Habits of Doryphora 10-lineata and of Carpocapsa pomonella.
- 978. RILEY, C. V. Curculio. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82. Answer to inquiry of J. Shearer; means against Conotrachelus nenuphar.
- 979. RILEY, C. V. Hop-vine caterpillars. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82. Communication from J. H. Graves, with answer; Mantis [= Phasmomantis]

carolina breeds as far north as Ogle County, Ill. Description of larva and imago of Hypena humuli [=scabra]; means against it.

- 980. RILEY, C. V. Wheat-worms. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

 Communication from C. Campbell, with answer; description of larva and image and habits of Panalia [— Associal faving line reverses of Tanahia makes.]
 - image and habits of Pyralis [=Asopia] farinalis; ravages of Tenebrio molitor; means against both species.
- 981. RILEY, C. V. [Root-borer.] < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82. Communication from S. T. Kelsey, with answer; ravages of an unknown
- gerambycid larva which destroys the roots of various plants.

 982. RILEY, C. V. Bark-lice on the pear. <Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

Letter from T. D. Plumb, with answer; Mytilaspis pomicorticis [=pomorum] lives on fruit of pear and crab-apple; the scale a secretion; extent of injuries caused by this insect.

- 983. RILEY, C. V. A nuisance made useful. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

 Means against Doryphora 10-lineata discredited.
- 984. RILEY, C. V. Strawberry leaf-roller. <Prairie Farmer, 5 October, 1867, [v. 36], n. s., v. 20, p. 212. S.-b. No. 2, p. 100.

 Answer to inquiry of N. R. Strong; Anchylopera [= Phoxopteris] fragariæ injures strawberry leaves; larvæ of Selandria [= Monostegia] rosæ also eat leaves of the strawberry.
- 985. RILEY, C. V. False caterpillars on the pine. < Prairie Farmer,
 5 October, 1867, [v. 36], n. s., v. 20, p. 212. S.-b. No. 2, p. 101.
 Answer to inquiry of A. S. Fuller; habits and description of Lophyrus lecontei.
- 986. RILEY, C. V. An unknown worm. < Prairie Farmer, 5 October, 1867, [v. 36], n. s., v. 20, p. 212. S.-b. No. 2, p. 101. Extract from Beaver Dam Citizen, with comments; habits and description of Eristalis sp.
- 987. [RILEY, C. V.] Bark-louse. < Prairie Farmer, 19 October, 1867, [v. 36], n. s., v. 20, p. —. S.-b. No. 2, p. 75.

 Benzine and soap as a means against bark-lice.
- 988. [RILEY, C. V.] Editorial excursion to the Rocky Mountains. <Prairie Farmer, 7 December, 1867, [v. 36], n. s., v. 20, pp. 353–354. S.-b. No. 2, pp. 112–113. List of plants, insects, and mammals observed.
- 989. RILEY, C. V. Bark-lice. <Prairie Farmer, 21 December, 1867, [v. 36], n. s., v. 20, p. 389. S.-b. No. 2, p. 112.

 Answer to inquiries of G. Backster and J. C. Cobbey; means against Mytilaspis pomicorticis [= pomorum].
- 990. RILEY, C. V. Root Aphis. < Prairie Farmer, 21 December, 1867, [v. 36], n. s., v. 20, p. 389. S.-b. No. 2, p. 112.

 Answer to inquiry of J. M. Jordan; means against the root and stem forms of Schizoneura lanigera.
- 991. [RILEY, C. V.] The Colorado potato-beetle. <Prairie Farmer, 21 December, 1867, [v. 36], n. s., v. 20, p. 389. S.-b. No. 2, p. 112. Communications from C. W. Murtfeldt asserting the possession of poisonous qualities by the larvæ of *Doryphora* 10-lineata.

- 992. RILEY, C. V. Cut-worm. Fall and spring plowing. <Moore's Rural New Yorker, ———, 1867. S.-b. No. 2, p. 94. Criticism of article by Agricola.
- 993. RILEY, C. V. Entomology. <Prairie Farmer Annual [No. 1 for 1868], 1867, pp. 53-59, 6 figs. S.-b. No. 2, pp. 125-126; No. 14, pp. 216-219.
 - Descriptions and figures of larva, pupa, and imago of Coptocycla [= Cassida] bivittata; habits, seasons, and means against it; list of Cassididæ injurious to sweet-potato. Descriptions of larva, nidus, and imago of Desmia maculalis, figures pupa; habits and seasons of and means against it. Figures Colaspis flavida; describes its supposed habits. Food-plants of Heliothis armigera. Figures larva and imago of Gortyna nitela; its food-plants, seasons, and hibernation. Figures larva and imago of Heliothis phlogophagus; synonymy, geographical distribution, and means against it; remarks on the balance of nature. Figures larva, pupa, cocoon, and imago of Asopia costalis; its synonymy and ravages.
- 994. RILEY, C. V. Bark-lice. < Prairie Farmer, 15 February, 1868, [v. 37], n. s., v. 21, p. 100. S.-b. No. 2, p. 118.
 - Answer to inquiry of J. Hawkins; natural limitation and parasites of *Mytilaspis pomicorticis* [=pomorum]; it attacks healthy and unhealthy trees; means against it; experience with washes.
- 995. RILEY, C. V. Potato beetle. < Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117. S.-b. No. 2, p. 119; No. 3, p. 58.
 - Answer to inquiry of F. T. Moore; invention of a machine for killing *Dory-phora* 10-lineata; hand-picking and heavy mulching the most available remedies.
 - 996. RILEY, C. V. Apple-tree plant-lice. <Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117. S.-b. No. 2, p. 119; No. 3, p. 58.
 - Letter from L. M., with answer; habits of and means against Aphis mali.
 - 997. RILEY, C. V. Apple-root blight. <Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117. S.-b. No. 2, p. 119; No. 3, p. 58.
 - Answer to inquiry of C. S. J.; Schizoneura lanigera not the cause of "rotten root."
 - 998. RILEY, C. V. Apple leaf-crumpler. < Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117. S.-p. No. 2, p. 119; No. 3, p. 58.
 - Answer to inquiry of J. M. Pearson; means against Phycita nebulo [= Acrobasis indiginella].
 - 999. RILEY, C. V. Tree-cricket. < Prairie Farmer, 14 March, 1868, [v. 37], n. s., v. 21, p. 164. S.-b. No. 2, p. 120; No. 3, p. 58.
 - Letter from J. J. Hnggins, with answer; place of oviposition of *Ecanthus niveus*; list of plants in which this species oviposits.
- 1000. RILEY, C. V. Egg-masses and cocoons on apple-trees. < Prairie Farmer, 14 March, 1868, [v. 37], n. s., v. 21, p. 164. S.-b. No. 2, p. 120; No. 3, p. 58.
 - Answer to inquiry of a correspondent; description of egg-masses and of larva of Orgyia leucostigma; beneficial influence and means of encouraging parasites.

1001. RILEY, C. V. Hop insects; Hop Aphis. < Prairie Farmer, 21

March, 1868, [v. 37], n. s., v. 21, p. 184. S.-b. No. 3, p. 58.

Brief account of and means against Human humalify and Photodom

Brief account of and means against Hypena humuli [= scabra] and Phorodon humuli.

1002. RILEY, C. V. Supposed eggs of the preying Mantis. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.

Answer to inquiry of J. H. Graves; cocoons of *Rhogas* sp. attached in a mass to twigs of an apple-tree.

- 1003. RILEY, C. V. Oak-tree borer. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Answer to inquiries of E. G. Mygatt; habits and food-plants of *Xyleutes* [= Cossus] robinia; means against it.
- 1004. RILEY, C. V. Maple-bark lice. <Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S. b. No. 2, p. 121; No. 3, p. 59.

Lecanium sp. found on twigs of sugar-maple.

- 1005. RILEY, C. V. Eggs of the katydid. <Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Answer to inquiry of W. Colwell; habits and food-plants of *Platyphyllum* [= Cyrtophyllus] concavus.
- 1006. RILEY, C. V. Eggs of tree-cricket in raspberry canes. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Letter from E. T. Nelson, with answer; oviposition of *Ecanthus niveus* in raspberry canes; means against this insect.
- 1007. RILEY, C. V. Bark-lice. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Answer to inquiry of C. D. Robinson; means against Mytilaspis pomicorticis [=pomorum].
- 1008. RILEY, C. V. Bark-lice again; the native species. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Letter from N. Coleman, with answer; comparative characters of *Mytilaspis* pomicorticis [=pomorum] and Diaspis harrisii [=Chionaspis furfurus]; food-plants of the latter.
- 1009. RILEY, C. V. Dahlia stalk borer. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.

 Answer to inquiry of S. M. Wierman; means against Gortyna nitela.
- 1010. RILEY, C. V. Apple-worm. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.

Answer to communication of S. J. C.; Carpocapsa pomonella hibernates as a larva within the cocoon.

- 1011. RILEY, C. V. False caterpillars on the Scotch and Austrian pines. <Prairie Farmer, 2 May, 1868, [v. 37], n. s., 21, p. 285, fig.

 Description and figures of larva, cocoou and imago, and of male antenna of
 - Lophyrus lecontei; food-plants and habits of larva; value of technical science.
- 1012. RILEY, C. V. Prevention of bark-lice. <Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.
 - Means against Mytilaspis pomicorticis [= pomorum] and Phycita nebulo [= Acrobasis indiginella]; figure of the latter.
- 1013. RILEY, C. V. Beetles in stomach of meadow-lark. < Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.
 - Fragments of Nottiglossa [= Oncometopia] undata and of Platynus excavatus or P. aruginosus found in the stomach of Sturnella magna.
- 1014. RILEY, C. V. Beetle on sugar-maple. < Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.

 Eburia quadrigeminata found under bark of a felled sugar-maple, also on honey-locust.
- 1015. RILEY, C. V. White worms in wells. <Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S. b. No. 3, p. 60. Fish or frogs placed in wells will free them from worms.
- 1016. RILEY, C. V. Tanzy for borers. < Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.

 Means against borers in apple- and peach-trees.
- 1017. RILEY, C. V. Peach-borer. <Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S. b. No. 3, p. 60.

 Refers to articles ou means against Ægeria [= Sannina] exitiosa.
- 1018. RILEY, C. V. Black grape-vine caterpillars. < Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60. Descriptions of larva and image of *Thyreus abbotii*; the abundance of parasites prevents serious injury by the larva.
- 1019. RILEY, C. V. The potato-beetle. < Cultivator and Country Gentleman, 21 May, 1868, v. 31, p. 378.

 Criticism of article on the "Colorado potato-bug" (Cultivator and Country Gentleman, 63 April, 1868, p. 309); use of hellehore against Decembers, 10
- Gentleman, 23 April, 1868, p. 309); use of hellebore against Doryphora 10-lineata.
- 1020. RILEY, C. V. The May-beetle; white grub. <Journ. of Agric., 1868, 4 figs. S.-b. No. 3, p. 10.

 Habits, food-plants, enemies, and means against Lachnosterna quercina [=fusca]; figures the larva, pupa, and imago of the same.
- 1021. RILEY, C. V. Canker-worm. < Colman's Rural World, 1868. S.-b. No. 3, p. 16.

 Habits of and means against Anisoptery's [= Paleacrita] vernata.
- 1022. RILEY, C. V. The apple-worm or codling-moth. <Colman's Rural World, 1868, fig. S.-b. No. 3, p. 16.
 - Seasons of and means against Carpocapsa pomonella; figures all stages except egg.

8 ENT

- 1023. RILEY, C. V. Remedy for the apple-borer. <Colman's Rural World, 1868. S.-b. No. 3, p. 16.

 Means against Saperda bivittata [=candida].
- 1024. RILEY, C. V. Grape-vine hoppers. < Colman's Rural World, 1868. S.-b. No. 3, p. 16.

 Tettigonia coagulata injurious to leaves of grape-vines.
- 1025. RILEY, C. V. Thousand-legged worm. <Colman's Rural World, 1868. S.-b. No. 3, p. 16.
 Habits of Julus [= Spirobolus] marginatus.
- June, 1868, [v. 37], n. s., v. 21, p. 397. S.-b. No. 3, p. 59.
 Habits, description of, and means against Spermophagus robiniæ.
- 1027. RILEY, C. V. Potato-beetle. <Prairie Farmer, 20 June, 1868, [v. 37], n. s., v. 21, p. 397. S.-b. No. 3, p. 59.
 Natural history of and means against Doryphora 10-lineata.
- 1028. RILEY, C. V. Large moth on apple-tree. <Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S.-b. No. 3, p. 60. The larva of Attacus cecropia spins its eoeoon on apple and other trees.
- 1029. RILEY, C. V. Will unimpregnated eggs hatch? < Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S.-b. No. 3, p. 60.
 - Remarks upon the statement by H. D., of Champaign, Ill., eoneerning the hatching of unimpregnated eggs of *Attacus cecropia*; mentions the families and genera of insects in which parthenogenesis occurs.
- 1030. [RILEY, C. V.] Wire-worms: Experiments in killing. | < Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S.-b. No. 3, p. 60.
 - Effect of several liquid preparations upon the larvæ of *Elateridæ* immersed in them; eorn soaked in turpentine grew unmolested by wire-worms.
- 1031. [RILEY, C. V.] [Pear-slug and current-worm.] < Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S.-b., No. 3, p. 60. Means against Selandria [= Eriocampa] cerast and Nematus ventricosus [=ri-besii].
- 1032. RILEY, C. V. What becomes of bumble-bees? < Cultivator and Country Gentleman, 2 July, 1868, v. 32, p. 18.

 Answer to Anna's "Bumble-bees;" only the queens of Bombus survive the winter; new colonies formed by hibernated queens.
- 1033. RILEY, C. V. Large gray straight-horned snout-beetle. < Prairie Farmer, 4 July, 1868, [v. 38], n. s., v. 22, p. 2–3, figs. S.-b. No. 3, pp. 31, 56.
 - Descriptions and figures of larva and imago of Ithycerus noveboracensis; method of oviposition; injuries to apple, oak, and other trees.
- 1034. RILEY, C. V. The seventeen-year Cicada. <Prairie Farmer, 4
 July, 1868, [v. 38], n. s., v. 22, p. 2. S.-b. No. 3, pp. 31, 56.
 Dates and localities of occurrences of Cicada [= Tibicen] septendecim; occur-

rence of C. [= T.] tredccim in Missouri.

- 1035. RILEY, C. V. Raspberry canes dying. <Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10, 1 fig. S.-b. No. 3, pp. 31, 56.

 Answer to inquiry of J. A. Jackson; death of raspberry canes discussed; in this case not the result of injury by insects.
- 1036. RILEY, C. V. Bag-worms. < Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10. S.-b. No. 3, pp. 31, 56.

 Letter from B. B. Warfield, with answer; habits, food-plants, and means against Thyridopteryx ephemeraformis; description of the female.
- 1037. RILEY, C. V. Oak and rose galls. <Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10. S. b. No. 3, pp. 31, 32, 56.

 Answer to inquiry of F. H. G.; descriptions of galls and images of Cynips [= Rhodites] bicolor on rose and of Cynips [= Andricus] seminator on white oak; occurrence of Cicada [= Tibicen] septendecim in Michigan.
- 1038. RILEY, C. V. Apple-borer and root Aphis. <Prairie Farmer, 11
 July, 1868, [v. 38], n. s., v. 22, p. 10. S.-b. No. 3, pp. 32, 56.

 Answer to inquiry of W. Colwell; ravages of Saperda bivittata [=candida]
 and of Schizoneura lanigera.
- 1039. RILEY, C. V. Evergreen plant-lice. <Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10. S.-b. No. 3, pp. 32, 56.

 Answer to inquiry of J. Cochrane; balsam-fir injured by Lachnus strobi?; means against it.
- 1040. RILEY, C. V. Bark-lice again. < Prairie Farmer, 18 July, 1868, [v. 38], n. s., v. 22, p. 18, 2 figs. S.-b. No. 3, pp. 34, 54.

 Answer to inquiries of J. Hodgeson, S. D., J. Sutherland, and J. H. G.; descriptions and figures of the scales of Mytilaspis pomicorticis [= pomorum] and Diaspis harrisii [= Chionaspis furfurus].
- 1041. RILEY, C. V. Larvæ of grape-vine flea-beetle. < Prairie Farmer, 18 July, 1868, [v. 38], n. s., v. 22, p. 18. S.-b. No. 3, pp. 34, 54. Answer to inquiry of J. A. Pettingil; leaves of grape-vine eaten by larvæ of Haltica chalybea; description of larva; habits and means against it.
- 1042. [RILEY, C. V.] A corn Curculio < Prairie Farmer, 25 July, 1868, [v. 38], n. s., v. 22, p. 26. S.-b. No. 3, p. 33.

 Communication from L. V. Smith, with answer; seasons and ravages of Sphenophorus sculptilis.
- 1043. RILEY, C. V. Ephemera flies; a hard story. < Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, pp. 36, 57. Extract from Peoria (Ill.) Transcript, with letter of J. Cochrane, of Havana, Ill.; notes on the above; appearance of swarms of Palingenia [= Hexagenia] bilineata in Illinois and of other may-flies in Europe; life-history of Ephemeridæ.
- 1044. RILEY, C. V. Driving potato-beetles. < Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, pp. 36.

 Doryphora 10-lineata can not be driven; Macrobasis unicolor can; care should be taken in sending the former through the mail, so as not to increase its distribution.
- 1045. RILEY, C. V. Gregarious walnut caterpillar. <Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, pp. 36, 57. Answer to inquiry of "a subscriber;" leaves of walnut destroyed by larvæ of Datana ministra; habits and means against the same.

- 1046. RILEY, C. V. Bugs on grape-vines. < Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, p. 36.

 Letter from J. H. Bingham, with answer; new growth of stems and leaves of grape-vines injured by Corimelana pulicaria; means against it.
- 1047. RILEY, C. V. Corn-worms. < Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, p. 36.

 Letter from E. Daggy, with answer; injury to maize by an undescribed species of Hadena?.
- 1048. RILEY, C. V. Large worm on apple-tree. <Prairie Farmer, 15
 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, p. 36.
 Answer to inquiry of J. M. Devore; description of *Papilio turnus*, Q, form glaucus, which pupated on trunk of apple-tree.
- 1049. RILEY, C. V. White-pine trees killed by borers. < Prairie Farmer, 26 September, 1868, [v. 38], n. s., v. 22, p. 98.
- 1050. RILEY, C. V. Swarms of butterflies. < Prairie Farmer, 26 September, 1868, [v. 38], n. s., v. 22, p. 98.
- 1051. RILEY, C. V. Worms feeding on the hawthorn. < Prairie Farmer, 26 September, 1868, [v. 38], n. s., v. 22, p. 98.
- 1052. RILEY, C. V. Twig borers, sack-bearers, etc. < Prairie Farmer.
 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1053. RILEY, C. V. Oil beetles. < Prairie Farmer, 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1054. RILEY, C. V. Apple-tree caterpillars. < Prairie Farmer, 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1055. RILEY, C. V. Twigs punctured by periodical Cicada. < Prairie Farmer, 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1056. RILEY, C. V. Report of committee on entomology. <Trans. Ill. Hortic. Soc. for 1867, 1868, n. s., v. 1, pp. 105-107, figs. 1-8.
 - Criticism of previous reports; effect of cold on Aphis mali; distribution, food-plants, and means against Doryphora 10-lineata; distinctions between D. 10-lineata and D. juncta; ravages of Lytta [= Epicauta] vittata, L. [=E.] cinerea, and L. atrata [=E. pennsylvanica] on potato, and L. [=Pomphopæa] enca on fruit-trees; means against them; ravages of Haltica [= Crepidodera] cucumeris on potato; habits, ravages, and specific characters of Cassididæ found on sweet potato; Saperda bivittata [=candida] biennial; history and development of Mytilaspis pemicorticis [=pomorum]; its food-plants, supposed hermaphroditism [parthenogenesis] and mode of production of scale of this species; difficulty of explaining the production of galls; habits, characters, food-plants, and ravages of Conotrachelus nenuphar, Carpocapsa pomonella, Heliothis armigera, Gortyna nitela, Mamestra picta, Procris [= Harrisina] americana, Selandria vitis [= Blennocampa pygmaa], Darapsa [= Ampelophaga myron, and Desmia maculalis; notes on other insects injurious to grape, apple, and wheat, and on Mycetophila persica, Emphytus [= Harpiphorus] maculatus, Agrotis [= Hadena] devastatrix, and Hadena juneta; note on recent writings by the author.

1057. RILEY, C. V. Entomology. <Prairie Farmer Annual [No. 2 for 1869], 1868, pp. 30-41, 6 figs. S.-b. No. 14, pp. 220-226.

Directions for collecting and preserving insects; descriptions of apparatus and cabinets; means against cabinet pests; descriptions and figures of all stages, except egg, of Lophyrus abbotii and L. lecontei; habits and foodplauts of and means against both species; figures of eggs, pupe, and imago of Cicada [= Tibicen] septendecim and of twig punctured by imago; dates and localities of occurrence of this species and of C. [= T.] tredecim; C. [= T.] cassinii a form of this insect; figures larva, puparium, and imago of Cephalemyia [= Estrus] ovis; habits and ravages of and means against it; imago viviparous in the nostrils of sheep.

1058. RILEY, C. V. The American Meromyza, Meromyza americana, Fitch. Attacking wheat just before it ripens. < Moore's Rural New Yorker, 30 January, 1869, v. 20, p. 71, fig. S.-b. No. 3, p. 83.

Habits and ravages of *Meromyza americana*; figures larva, pupa, and injured stalk of wheat; related Europeau species have similar habits; remedies and natural enemies.

1059. RILEY, C. V. First annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <4th Ann. Rept. State Board of Agric. for 1868, March, 1869, pp. 182–187, 98 figs. Separate: <Jefferson City, Mo., March, 1869, pp. 182–187, 2 pl., 98 figs. Notice: <Amer. Ent., April, 1869, v. 1, p. 165.

CONTENTS.

Page. Introductory NOXIOUS INSECTS. THE BARK-LICE OF THE APPLE-TREE 7 Two species known to occur in the United States, 7-Harris's barklouse not numerous enough to do material damage, 7. The oyster-shell bark-louse, Mytilaspis pomorum..... 7 Difference in the scales of the two species, 7-Introduction of the oyster-shell bark-louse from Europe and its spread in the United States, 8-Precautionary measures to prevent its introduction into Missouri, 8-Its habits studied by different observers in 1867, 9—Seasonal notes on the development of the insect, 10— Only one annual brood in Missouri, 12-Formation and nature of the scale, 12-Rare occurrence of males, 14-Difference of opinion amoug observers as to the mode of growth of the scale, 14-Difficulty of explaining the spread of the insect from one tree to another, 15-Its occurrence upon other kinds of trees, 15-Natural enemies, 16-Artificial remedies, 16-Examination of young trees before planting, 16-Washing with alkalics, etc., 17-Scrubbiug the branches with a stiff brush, 17—Fumigating, 17—Application of sheep-manure, 17-Washes in general ineffective, 17-The insect can most successfully be fought during three or four

days of the year only, 18.

1059. RILEY, C. V.—Continued.

It is more numerous in trees on high land than on low ground, 42—Extent of its injury, 43—Its larva, 43—Appearance of the imago, 43—The hole made by the young larva, 44—It remains nearly three years in the larva state, 44—Its pupa state, 44—Remedies, 45—Alkaline washes, 45—Killing the larva by hot water, 45—Cutting out the larva, 46. The flat-headed apple-tree borer, Chrysobothris femorata	THE PERIODICAL CICADA, Tibicon septendecim	18
broods that will appear in the next 17 years in the different States, 42. APPLE-TREE BORERS	Its singular life history, 18—Seventeen and thirteen year races, 19—The two races not distinct species, 19—Two distinct forms occurring in both broods, 20—Season of their appearance and disappearance, 22—Natural history and transformations, 22—Elevated chambers of the pupa, 22—The larvæ frequently occurring at great depth in the ground, 24—The operation of emerging from the pupa, 24—Only the males are capable of singing, 24—Trees in which the females deposit their eggs, 24—Mode of oviposition, 24—The newly hatched larva, 25—The W on the wings of the Cicada, 25—Enemics, 26—Fungus infesting the imago, 26—The sting of the Cicada, 26—Wide-spread fear of the insect on account of its supposed stinging powers, 26—Explanations of the sting, 27—Injury caused by the insect, 29—by the larva, 29—by the imago, 29—Fruitless attempts to stop the injury, 30—Chronological table of all well-ascertained broods in the United States, 30—The insect will appear during the next 17 years somewhere	18
The peach borer, Sannina exitiosa Its nature, 47—Differences in the sexes, 48—Remedies, 48—The mounding system the best remedy, 48—Testimony as to the value of the mounding system, 48—Other remedies, 49. The Plum Curculio, Conotrachelus nenuphar Difference of opinion among authors on some points in its natural history, 52—It causes the spread of the peach-rot, 52—Fruit trees attacked and those not attacked by it, 56—Reamonds of egg-laying, 54—It has one annual brood, 55—Walsh's experiments to show that it is two-brooded, 55—Natural remedies, 57; The subangular ground-beetle, 58; Ground-beetle larva, probably of the trees the most effectual method, 60—Dr. Hull's Curculio catcher, 60—	broods that will appear in the next 17 years in the different	
It is more numerous in trees on high land than on low ground, 42—Extent of its injury, 43—Its larva, 43—Appearance of the imago, 43—The hole made by the young larva, 44—It remains nearly three years in the larva state, 44—Its pupa state, 44—Remedies, 45—Alkaline washes, 45—Killing the larva by hot water, 45—Cutting out the larva, 46. The flat-headed apple-tree borer, Chrysobothris femorata	States, 42.	
It is more numerous in trees on high land than on low ground, 42—Extent of its injury, 43—Its larva, 43—Appearance of the imago, 43—The hole made by the young larva, 44—It remains nearly three years in the larva state, 44—Its pupa state, 44— Remedies, 45—Alkaline washes, 45—Killing the larva by hot water, 45—Cutting out the larva, 46. The flat-headed apple-tree borer, Chrysobothris femorata		42
The flat-headed apple-tree borer, Chrysobothris femorata	It is more numerous in trees on high land than on low ground, '42—Extent of its injury, 43—Its larva, 43—Appearance of the imago, 43—The hole made by the young larva, 44—It remains nearly three years in the larva state, 44—Its pupa state, 44—Remedies, 45—Alkaline washes, 45—Killing the larva by hot	42
Its nature, 47—Differences in the sexes, 48—Remedies, 48—The mounding system the best remedy, 48—Testimony as to the value of the mounding system, 48—Other remedies, 49. The plum Curculio, Conotrachelus nenuphar	The flat-headed apple-tree borer, Chrysobothris femorata. Differences between it and the foregoing species, 46—Habits of the bcetle, 47—Amount of injury caused by it, 47—Parasite attacking	.46
Its nature, 47—Differences in the sexes, 48—Remedies, 48—The mounding system the best remedy, 48—Testimony as to the value of the mounding system, 48—Other remedies, 49. The plum Curculio, Conotrachelus nenuphar	THE PEACH BORER, Sannina exitiosa	47
Difference of opinion among authors on some points in its natural history, 50—Reasons for this difference of opinion, 51—Facts in its natural history, 52—It causes the spread of the peach-rot, 52—Fruit trees attacked and those not attacked by it, 53—It may hibernate as larva or pupa, but does generally as imago, 53—Mode of egg-laying, 54—It has one annual brood, 55—Walsh's experiments to show that it is two-brooded, 55—Natural remedies, 56—No parasites known to infest it, 56—Enemics, 57; The Pennsylvania soldier-beetle and its larva, 57; Lacewing-larva, 57; The subangular ground-beetle, 58; Ground-beetle larva, probably of the Pennsylvania ground-beetle, 59—Hogs as Curculio destroyers, 59—Artificial remedies, 60—Jarring the trees the most effectual method, 60—Dr. Hull's Curculio catcher, 60—	Its nature, 47—Differences in the sexes, 48—Remedies, 48—The mounding system the best remedy, 48—Testimony as to the	
Difference of opinion among authors on some points in its natural history, 50—Reasons for this difference of opinion, 51—Facts in its natural history, 52—It causes the spread of the peach-rot, 52—Fruit trees attacked and those not attacked by it, 53—It may hibernate as larva or pupa, but does generally as imago, 53—Mode of egg-laying, 54—It has one annual brood, 55—Walsh's experiments to show that it is two-brooded, 55—Natural remedies, 56—No parasites known to infest it, 56—Enemics, 57; The Pennsylvania soldier-beetle and its larva, 57; Lacewing-larva, 57; The subangular ground-beetle, 58; Ground-beetle larva, probably of the Pennsylvania ground-beetle, 59—Hogs as Curculio destroyers, 59—Artificial remedies, 60—Jarring the trees the most effectual method, 60—Dr. Hull's Curculio catcher, 60—	THE PLUM CURCULIO Conotrachelus nenunhar	50
the most effectual method, 60—Dr. Hull's Curculio catcher, 60—	Difference of opinion among authors on some points in its natural history, 50—Reasons for this difference of opinion, 51—Facts in its natural history, 52—It causes the spread of the peach-rot, 52—Fruit trees attacked and those not attacked by it, 53—It may hibernate as larva or pupa, but does generally as imago, 53—Mode of egg-laying, 54—It has one annual brood, 55—Walsh's experiments to show that it is two-brooded, 55—Natural remedies, 56—No parasites known to infest it, 56—Enemics, 57; The Pennsylvania soldier-beetle and its larva, 57; Lacewing-larva, 57; The subangular ground-beetle, 58; Ground-beetle larva, probably of the Pennsylvania ground-beetle, 59—Hogs as Cur-	
	,	

·,	
1059. RILEY, C. V.—Continued.	
THE COOLING MOTH OR APPLE WORM, Carpocapsa pomonella	62
It is common wherever apples are grown, 62—Description of the	
insect in its different states, 63—Its life-history, 63—Other fruits	
attacked by it, 64—Remedies, 65—Picking up the fallen fruit,	
65—Entrapping the worms the best remedy, 66—Trimble's hay	
band system and how to apply it, 66—Attracting the moth by	
fires, 67.	
Cut-worms	67
The natural history of twelve distinct species, 67—Definition of	
the term "cut-worm," 67—Habits of cut-worms, 67—Their nat-	
ural history briefly given, 68—Difficulty of breeding them in	
captivity, 69—Climbing cut-worms, 69—Injury done by them to orchards, 69—Fruit trees and shrubs they attack, 70—They at-	
tack large trees, 71.	
The variegated cut-worm, Agrotis saucia	72
The full-grown larva, 72—The eggs, 72—Habits of the larva, 72—	• ~
Cut-worm moths deposit their eggs on the leaves and not on the	
ground, 73—The imago, 73—Description of the insect as larva,	
pupa, and imago, 74.	
The dark-sided cut-worm, Agrotis messoria	74
General characters of the larva, 74—Habits of, and injury done by	
it, 75—Description of imago, 75; of the larva and chrysalis, 76.	
The climbing cut-worm, Agrotis scandens	7 6
Injury done by the larva, 77—General characters of the larva, 77,	
of the moth, 78—Description of the larva, 78; of the imago, 78.	
The w-marked cut-worm, Agrotis clandestina	7 9
General characters of the larva, 79—Plants it attacks, 79—Char-	
acters of the moth, 79—Description of the larva and chrysalis, 79.	00
The greasy cut-worm, Agrotis ypsilon	80
tobacco plants, 80—General characters of the moth, 80—Descrip-	
tion of larva, chrysalis, and imago, 81.	
The western striped cut-worm, Agrotis herilis	81
Resemblance of its larva to that of the corn rustic, 81—General	D.
characters of the worm and moth, 82—Description of the larva,	•
82.	
The dingy cut-worm, Agrotis subgothica	82
Difference between it and the foregoing species, 82—General char-	
acters of pupa and imago, 82—At least three species of our cut-	
worms are difficult to distinguish, 83—Description of larva, chrys-	
alis, and imago, 83.	
The glassy cut-worm, Hadena devastatrix	83
Habits and general characteristics of the larva, 83—Characteris-	
tics of the moth, 84—Description of larva and chrysalis, 84.	0.4
The speckled cut-worm, Hadena subjuncta	84
Characteristics and habits of the insect, 84—Description of larva,	
chrysalis, and imago, 85.	86
The small white bristly cut-worm, Hadena renigera	00
tion of larva, chrysalis, and imago, 86.	
Other cut-worms	87
Fitch's account of the corn cut-worm and the yellow-headed cut-	
worm, 87.	
The wheat cut-worm	87
Injury caused by it, 87—Description of the larva, 88.	

1059. RILEY, C. V.—Continued.

illi, c. v.—continucu.	
Cut-worms—Continued.	
Remedies against cut-worms Natural enemies, 89; Microgaster militaris, 89; Paniscus geminatus, 89; The spined soldier-bug, 89; The cut-worm lion, 89—Other euemies, 90—Artificial remedies for climbing cut-worms, 90; for common field cut-worms, 91.	89
Insects infesting the potato	91
The stalk-borer, Gortyna nitela	92
The potato stalk-weevil, <i>Trichobaris trinotata</i>	93
The potato- or tomato-worm, Protoparce celeus. It can not sting with its horn, 95—Its chrysalis, 95—How the image differs from the tobacco-worm moth, 95—Remedies and parasites, 96.	95
Blister-beetles, Meloidæ The striped blister-beetle, 96—The ash-gray blister-beetle, 97— The black-rat blister-beetle, 98—The black blister-beetle, 98— The margined blister-beetle, 98—Synonymical remarks, 98— Remedies for blister-beetles, 99.	97
The three-lined leaf-beetle, Lema trilineata Merdigerous habit of the larva, 99—It has two annual broods, 100—Other notes on the habits of the insect, 100.	99
The cucumber flea-beetle, Crepidodera cucumeris	101
The Colorado potato-beetle, Doryphora 10-lineata	101
Its past history and future progress, 101—Its native home, 101—Its gradual spread eastward, 102—Its confusiou with the bogus Colorado potato-beetle, 103—How the two species differ in habits, 104; in their larval states, 104; in the egg state, 105—Description of the larva of Doryphora juncta, 106—Differences in the imagos of the two species, 106—Habits of the Colorado potato-beetle, 107—When it appears and disappears, 107—Number of eggs laid by each female, 107—Food-plants, 107—Singular fact that D. juncta has not acquired the habit of attacking the potato, 108—Natural remedies, 109—Complicated economy of nature, 109—Decrease in the number of potato-beetles on account of increase in the number of parasites, 109—The Colorado potato-beetle parasite, 111—Its general character and habits, 111—Description of Lydella doryphora, 111—Lady-birds and their larva, 112—The spined soldier-bug, 113—The common squash-bug, erroneously considered an enemy of the potato-beetle, 113—The bordered soldier-bug, 114—The many banded robber, 114—The rapacious soldier-bug, 114—The Virginia tiger-beetle, 115—The fiery ground-beetle, 115—Blister-beetles, 115—The larvae not	
touched by fowl, 115—Artificial remedies, 116—Ineffectiveness of mixtures tried, 116—Killing the beetle early in spring, 116—Pincers for crushing the insect, 116—Benson's machine, 116—Proper choice of varieties of potatoes, 117—The pest will overrun the Eastern States, 117—Carelessness in transmitting specimens of the beetle, 117.	

1059.	RILEY, C. V.—Continued.	
	The apple-root plant-louse, Schizoneura lanigera Three distinct kinds of rots affecting the roots of apple-trees, 118—The root-louse the cause of one of these rots, 118—The cause of the other rots still hidden, 119—The root-louse especially in- jurious in southerly latitudes, 119—It occurs also on other parts of the tree besides the root, 120—Description of the winged lonse, 120—Fitch's description of the winged form refers to another species, 120—The root-louse belongs to the genns Eriosoma, 121—Natural enemies, 121; Chalcis-fly, 121; the root-louse Syr- phns-fly, 121; Scymnus cervicalis, 122—Artificial remedies, 123.	118
	THE WOOLLY ELM-TREE LOUSE, Schizoneura rileyi Its general appearance and habits, 123—Description of the winged form, 124.	123
	Insects injurious to the grape-vine. The new grape-root borer, Prionus laticollis Reports on the damage caused by it, 124—Description of the larva, 126—It belongs probably to the cylindrical Orthosoma, 126—Former accounts of the natural history of this beetle, 127—Its injury known for several years, 127—Remedies, 128.	124 124
	The grape curculio, Craponius inequalis	128
	The grape-seed Curculio, <i>Isosoma vitis</i>	129
	The grape-cane gall Curculio, Ampeloglypter sesostris	131
	The grape-vine Fidia, Fidia viticida It is very injurious in Missouri, 132—Habits of the beetle, 132— Remedies, 133.	132
	The grape fruit-worm, Eudemis botrana. Amount and extent of the injury caused by it, 133—Characteristics of the larva, 134—Transformations, 134—Description of larva, chrysalis, and imago, 135—Remedies, 135.	133
	The eight-spotted forester, Alypia 8-maculata Characteristics of the larva, 136—It is not numerous enough to cause serious injury, 136—Other caterpillars resembling it, 136.	136
	The grape-vine plume, Oxyptilus periscelidactylus	137
•	The snowy tree-cricket, <i>Œcanthus niveus</i> Characteristics of the insects, 138—It is injurious, 138—Nature of the injury caused by it, 138—Remedy, 139.	138
,	THE RASPBERRY GEOMETER, Synchlora rubivoraria	139
	THE GOOSEBERRY FRUIT-WORM, Dakruma convolutella	140

1059.	RILEY, C. V.—Continued.	
	THE STRAWBERRY LEAF-ROLLER, Phoxopteris fragariæ Extent and nature of the damage caused by it, 142—Habits of the insect, 142—Accounts of its injury in Indiana and Illinois, 142—Remedy, 143—Description of the image and larva, 143.	142
	The white-marked tussock-moth,* Orgyia leucostigma. The cgg-mass, 144—The larva and larval changes, 144—The full-grown larva, 145—Habits of the larva, 145—Mode of casting off the larval skin, 145—The cocoon, 146—The imago, 146—Two an-	144
	unal broods, 146—Food-plants, 146—Remedies, 147. THE BAG-WORM, alias BASKET-WORM, alias DROP-WORM, Thyridopteryx ephemariformis	147
	Its geographical distribution, 148—Injury caused by it, 148—The egg, 148—The larva and its growth, 148—Habits of the larva, 149—The chrysalis, 149—The sex distinguishable in the chrysalis state, 149—The imago, 149—Food-plants, 150—Parasites, 150; Cryptus inquisitor, 150; Hemiteles thyridopterygis, n. sp., 150—Remedies, 151.	
	The AILANTHUS-WORM, Œta punctella. Injury done to the Ailanthus tree, 151—Habits of the larva, 151— The chrysalis, 151—The imago, 152—Geographical distribution, 152—Remedy, 152—Description of larva and chrysalis, 152; of	151
	the imago, 153. The Walnut Tortrix, Cacacia rileyana Habits of the larva, 153—General appearance of the moth, 153— Phytophagic form of the insect on snowberry, 153—Description of larva, chrysalis, and imago, 154; of the variety symphoricarpi, 154.	153
	THE SEED-CORN MAGGOT, Anthomyia zeas Accounts of damage caused by it, 154—The maggot, 155—Transformation, 155—Description of the imago, 155—Remedy, 155—Habits of Anthomyia larvæ, 156.	154
	The white grub, Lachnosterna fusca. Account of the damage caused by it, 156—Injury done by the perfect insect, 157—Résumé of its life-history, 157—Remedies, 157—Regularity in the appearance of the beetle, 158—Accounts of the fungus infesting the white grub, 158.	156
	THE AMERICAN MEROMYZA, Meromyza americana. Nature of the damage caused by it, 159—Characteristics of larva, chrysalis, and imago, 160—European Diptera with similar habits, 160—Remedics, 161.	159
	The sheep bot-fly of head-maggot, Estrus ovis. The insect in its different states, 161—Its larva, 162—Pupa, 162—Characteristics of the imago, 162—Fatal results of the presence of the maggot in the head of the sheep, 163—Rabbits attacked by gad-fly, 164—Testimony regarding the viviparous habits of the bot-fly, 164—Remedics, 165.	161
	Insect enemies of the honey-bee. The bec-moth or wax-worm, Galleria cereana General appearance of the moth, 166—There are no moth-proof bee-hives, 166—Habits of the worm, 167—How its presence in the hive may be recognized, 167—Prevention and remedy, 167.	166 166

^{*} Reprint: <Cultivator and Country Gentleman, 14 May, 1874, v. 39, pp. 310-311, 3 figs.

DIDLIGHARITY OF ECONOMIC ENTOMOLOGY.	120
1059. RILEY, C. V.—Continued. INSECT ENEMIES OF THE HONEY-BEE—Continued. The bee-killer, Promachus apivorus. It is an Asilid fly, 168—Mr. Thompson's account of the fly, 168—How it captures and kills bees, 168—No remedy known, 168.	168
BENEFICIAL INSECTS.	
THE REAR-HORSE, alias CAMEL-CRICKET, alias DEVIL'S RIDING-HORSE, Phasmomantis carolina Its food, 169—How it grasps its prey, 169—Difference in the sexes, 170—The larva, 170—The egg-mass, 170—The mode of egg-laying, 170—Voracious disposition of the Mantis, 171—Its beneficial influence, 171—Trachina parasite of the Mantis, 171.	169
INNOXIOUS INSECTS.	
The Solidago Gall-Moth, Gelechia gallesolidaginis Gall caused by Trypeta solidaginis, 173—Gall produced by the Solidago gall-moth, 173—Its natural history, 173—Provision of the larva for its protection within the gall, 174—Previous account of the gall, 174—Gall caused by Cochylis hilarana on Artemisia campostris, 175—Description of the Solidago gall-moth as larva, chrysalis, and imago, 175—Parasites attacking it, 175; the inflating chalcis, 176; Eurytoma bolteri, n. sp., 176; Hemiteles (?) eressonii, n. sp., 177; Microgaster gelechiæ, n. sp., 177; other parasites, 178—Oberea larvæ intruding in the gall, 178. The Chickweed [=knotweed] Geometer, Hæmatopis grataria Its natural history, 179—Description of larva and chrysalis, 179. The thistle plume, Platyptilus carduidaetylus Work of its larva on thistle-heads, 180—Description of the larva, chrysalis, and imago, 180.	173 179 180
1060. RILEY, C. V. Eggs of the Mantis or rear-horse. < Mo	ore's
Rural New Yorker, 10 April, 1869, v. 20, p. 234, 2 figs. No. 3, p. 52. Answer to inquiry of T. C. Bartle; description of Mantis [=Phasmomo carolina; its food, habits, oviposition, and parasites; figures its egg-mand those of Phylloptera [Amblycorypha] oblongifolia.	antis]
1061. RILEY, C. V. Native bark-lice on apple-trees. < Prairie Far	

17 April, 1869, [v. 39], n. s., v. 23, p. 122, 1 fig. S.-b. No. 3, p. 57.

Answer to inquiry of T. B. Gardner; life history of and means against Diaspis harrisii [= Chionaspis furfurus]; figure of the same.

1062. RILEY, C. V. Curculio. < Prairie Farmer, 17 April, 1867, [v. 39], n. s., v. 23, p. 122. S.-b. No. 3, p. 57.

Answer to inquiry of James Weed; habits of and means against Conotrachelus nenuphar.

1063. RILEY, C. V. Cherry-tree borers. < Prairie Farmer, 17 April, 1869, [v. 39], n. s., v. 23, p. 122. S.-b. No. 3, p. 57.

Answer to inquiry of Onargo Horticultural Society; larvæ of Buprestis [=Dircera] divaricata and of Trochilium sp., injurious to cherry-trees; softmaple attacked by T. [= Egeria] acerni; means against these insects.

- 1064. RILEY, C. V. White-grub fungus. < Prairie Farmer, 15 May, 1869, [v. 39], n. s., v. 23, p. 154. S.-b. No. 3, p. 76.
 - Letter from D. W. Tindall, with answer; ravages of Lachnosterna quercina [=fusca] in Clinton County, Mo., in 1868; Torrubia growing on the larvæ in 1869.
- 1065. RILEY, C. V. The canker-worm, Anisopteryx vernata Peck. <Moore's Rural New Yorker, 29 May, 1869, v. 20, p. 345, figs. S.-b. No. 3, p. 73.
 - Description, figures, and natural history of canker-worms; the two species are confounded.
- 1066. RILEY, C. V. The seed corn maggot, Anthomyia zeas, Riley. Destroying the seed after it is planted. < Moore's Rural New Yorker, June, 1869, fig. S. b. No. 3, p. 81. Advance print, with changes: <1st Ann. Rept. State Ent. Mo., March, 1869, pp. 154–156, figs. 86–87.
 - See No. 1059 for synopsis of contents.
- Yorker, 10 July, 1869, v. 20, p. 443. S.-b. No. 3, p. 83.

 Answer to inquiry of G. J. Magee; means against Aphis [= Myzus] cerasi.
- 1068. RILEY, C. V. Gooseberry span-worms. < Moore's Rural New Yorker, 10 July, 1869, v. 20, p. 443. S.-b. No. 3, p. 83.
 - Letter from A. Yancey, with answer; occurrence of Eufitchia ribearia in Iowa; habits and means against it; habits of and influence of weather on Blissus leucopterus; scarcity of Doryphora 10-lineata in Iowa.
- 1069. RILEY, C. V. Apple-leaf crumpler mistaken for Curculio. < Prairie Farmer, 10 July, 1869, [v. 40], n. s., v. 24, p. 218, fig. S.-b. No. 3, p. 80.
 - Letter from B. T. Taylor; improper use of the word Curculio; figures larvacase and image of *Phycita nebulo* [= Acrobasis indiginella]; means against it; outline figure of Conotrachelus nenuphar.
- 1070. RILEY, C. V. Peach-tree borer. <Prairie Farmer, 10 July, 1869, [v. 40], n. s., v. 24, p. 218. S.-b. No. 3, p. 80.
 - Answer to inquiry of C. Allen; habits of and means against *Egeria* [=Sannina] exitiosa.
- 1071. RILEY, C. V. To protect plums from Curculio. Prairie Farmer, 10 July, 1869, [v. 40], n. s., v. 24, p. 218. S.-b. No. 3, p. 80.
 - Answer to inquiry of H. H. S.; recommends the collection of the image of Conotrachelus nenuphar by jarring.
- 1072. RILEY, C. V. White-grub; information wanted. < Prairie Farmer, 28 August, 1869, [v. 40], n. s., v. 24, p. 274, 4 figs. S.-b. No. 3, p. 95.
 - Letter from A. T. V., with answer; habits of and means against *Lachnosterna* quercina [=fusca]; figures of larva and imago.
- 1073. RILEY, C. V. Unknown corn pest. < Prairie Farmer, 28 August, 1869, [v. 40], n. s., v. 24, p. 274. S.-b. No. 3, p. 95.
 - Letter from E. B. Hickey, with answer; ravages of cut-worms and of an unknown pest on maize; brief description and habits of the latter.

- 1074. RILEY, C. V. Larva of the grape-vine flea-beetle. <Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, pp. 100, 103.
 - Answer to inquiry of D. D. Vosburgh; habits, ravages, and means against Graptodera [= Haltica] chalybea.
- 1075. RILEY, C. V. Rose bug. < Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, pp. 100, 103.
 - Answer to inquiry of a subscriber; description of imago, transformations of, and means against *Macrodactylus subspinosus*.
- 1076. RILEY, C. V. Large green caterpillar on the apple. <Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100.
 - Answer to inquiry of S. E. A. Palmer; descriptions of larva and imago of Attacus cecropia; its habits and seasons.
- 1077. RILEY, C. V. Conical galls on leaves of wild grape-vine. (Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100.
 - Answer to inquiry of D. McClaine; description of galls of Cecidomyia vitisviticola; characters of the larvæ of Cecidomyia.
- 1078. RILEY, C. V. A strange bug. < Moore's Rural New Yorker, 28
 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100; No. 4, p. 3.
 The insects described in North Carolina's "A strange bug," were probably

 Psocus venosus; habits of the genus Psocus.
- 1079. RILEY, C. V. Currant-worms and black-currants. < Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100; No. 4, p. 3.
 - Critical review of Addi on currant-worms; in North America three species of larvæ feed on the leaves of currants, and two species of borers live within the stem.
- 1080. RILEY, C. V. Curculio. < Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100; No. 4, p. 3. Commendation of an editorial criticism of Northwest's article on the Curculio.
- 1081. [RILEY, C. V.] Insects injurious to the grape-vine. [No. 1.] <Amer. Ent., August, 1869, v. 1, pp. 231–234, figs. 169–173. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 87–91, figs. 60–63.
 - Treats of *Prionus laticollis* and *P. imbricornis*. See No. 1127 for synopsis of contents.
- 1082. [RILEY, C. V.] Insects infesting the sweet-potato. <Amer. Ent., August, 1869, v. 1, pp. 234-238, figs. 174-181. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 56-64, figs. 26-38.
 - Treats of tortoise-beetles, Cassidida. See No. 1127 for synopsis of contents.
- 1083. RILEY, C. V. The borers. <Western Rural, September, 1869. S.-b. No. 4, p. 2.
 - Means against Ægeria cucurbitæ [= Melittia ceto].

- 1084. RILEY, C. V. Supposed bark-lice eggs in Missouri. < Prairie Farmer, 4 September, 1869, [v. 40], n. s., v. 24, p. 282. S.-b. No. 3, p. 95.
 - Letter from J. Reed, with answer; eggs of a moth mistaken for those of Mytilaspis pomicorticis [= pomorum]; the latter unknown in Missouri.
- 1085. RILEY, C. V. New York weevil on apple-trees. < Prairie Farmer, 4 September, 1869, [v. 40], n. s., v. 24, p. 282, 3 figs. S. b. No. 3, p. 95.
 - Letter from R. M. Guy, with answer; figures of larva and imago, description of imago, geographical distribution, methods of oviposition, food-plants, and means against *Ithycerus noveboracensis*.
- 1086. [RILEY, C. V.] Insects injurious to the grape-vine. No. 2. < Amer. Ent., September-October, 1869, v. 2, pp. 22-24, figs. 12-16. Reprint, with slight changes. < 2d Ann. Rept. State Ent. Mo., March, 1870, pp. 71-73, figs. 44-48.
 - Treats of Charocampa pampinatrix [= Ampelophaga myron]; see No. 1127 for synopsis of contents.
- 1087. RILEY, C. V. Larva of the imperial moth. < Prairie Farmer, 9
 October, 1869, [v. 40], n. s., v. 24, p. 322. S.-b. No. 3, p. 92.
 - Letter of C. H. Thayer, with answer; description of larva, pupa, and imago of Ceratocampa [= Eacles] imperialis; food-plants of the same and of C. [= Citheronia] regalis.
- 1088. RILEY, C. V. Apple snout-beetle or four-humped Curculio. < Prairie Farmer, 9 October, 1869, [v. 40], n. s., v. 24, p. 322. S.-b. No. 3, p. 92.
 - *Letter from R. W. Gandy, with answer; habits and description of Anthonomus quadrigibbus; recommends jarring.
- New Yorker, 20 November, 1869, v. 20, p. —. S.-b. No. 3, pp. 107; 112.
 - Critical review of several recent articles upon tomato-worms; structure and harmlessness of larvæ of Sphingidæ; geographical distribution of Sphinz [=Protoparce] carolina and S. quinquemaculata [=P. celeus].
- 1090. [RILEY, C. V.] The bag-worm, alias basket-worm, alias dropworm, Thyridopteryx ephemeræformis, Haw. <Amer. Ent., November, 1869, v. 2, pp. 35–38, fig. 24.
 - Geographical distribution, food-plants, seasons, transformations, parasites, copulation, and oviposition of and means against *Thyridopteryx ephemeræ-formis*; figures and descriptions of larva, larva-cases, pupa, and imago; description of eggs; abodes constructed by insects.
- 1091. [RILEY, C. V.] Insects injurious to the grape-vine. No. 3. < Amer. Ent., November, 1869, v. 2, pp. 54–55, figs. 33–35. Reprint, with slight changes. <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 74–76, figs. 49–51.
 - Treats of Philampelus achemon; see No. 1127 for synopsis of contents.
- 1092. RILEY, C. V. The saddle-back caterpillar. < Moore's Rural New Yorker, 4 December, 1869, v. 20, p. —, fig. S.-b. No. 3, p. 103.
 - Letters from A. W. Baker and G. T. Cost, with answer; figures and characterization of the larva of *Empretia stimulea*; food-plants, urticating properties, and transformations; description of the imago.

- - Answer to communication of W. J. B., concerning Cicada [= Tibicen] septendecim.
- - Occurrence of Goniocotes hologaster in great numbers on a hen; means against the same; figure of it.
- - Habits and place of oviposition of Agrotis inermis [=saucia]; description of its eggs and larve.
- 1097. RILEY, C. V. That glow-worm. < Cultivator and Country Gentleman, 6 January, 1870, v. 35, p. 5, fig. S.-b. No. 4, p. 23. Reprint: < Amer. Ent., October, 1880, v. 3, n. s., v. 1, p. 254, fig.
 - Figures larva and image of *Photuris pennsylvanica*; *Photinus pyralis* also luminous in larval and adult stages; comparison with *Lampyris noctiluca*.
- 1098. [RILEY, C. V.] In memoriam. <Amer. Ent., December, 1869– January, 1870, v. 2, pp. 65–68.

 Biographical and obituary notice of B. D. Walsh.
- 1099. [RILEY, C. V.] The harlequin cabbage-bug, Strachia histrionica, Hahn. <Amer. Ent., December, 1869–January, 1870, v. 2, pp. 79–80, fig. 56.
 - Enumeration of the enemies of the cabbage in the United States; methods of their injury; importation and spread of some species; figures of the imago; description of eggs, habits, geographical distribution, seasons, and odors of Strachia [= Murgantia] histrionica; extract from G. Lincecum's "Texan cabbage-bug."
- 1100. [RILEY, C. V.] An entomologist caught napping. <Amer. Ent., December, 1869-January, 1870, v. 2, p. 84.
 - Criticism of the view that trees, grasses, or any other particular forms of vegetation are the natural coverings of the earth; this criticism applied especially to John Curtis.
- 1101. [RILEY, C. V.] Poisonous qualities of the Colorado potato-bug. Amer. Ent., December, 1869–January, 1870, v. 2, pp. 85–86. Extract from "Winona Republican," with remarks upon the poisonous nature of the blood of *Doryphora* 10-lineata.
- 1102. [RILEY, C. V.] Insects injurious to the grape-vine. No. 4. <Amer. Ent, December, 1869-January, 1870, v. 2, pp. 89-90, figs. 58-59. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 76-78, figs. 52-53.
 - Treats of Philampelus satellitia [=pandorus]; see No. 1127 for synopsis of contents.

- 1103. [RILEY, C. V.] Toads vs. bugs. < Amer. Ent., December, 1869–January, 1870, v. 2, p. 91.
 - Translation of extracts from Fogt's "Noxious and beneficial animals;" existence of a commerce in toads between France and England; usefulness, tameness, and gratitude of toads.
- 1104. [RILEY, C. V.] The tomato-worm again. <Amer. Ent., December, 1869-January, 1870, v. 2, pp. 91-92.
 - Reprint, with comments, of article entitled "The tomato-worm," from Syracuse Standard; absurd nature of newspaper accounts of insects.
- 1105. [RILEY, C. V.] Mr. Walsh's successor. < Amer. Ent., December, 1869–January, 1870, v. 2, pp. 92–93.
 - Condition in which B. D. Walsh left the office of State Entomologist of Illinois at his death; plans of his intended second report; provisions for the preparation of the same and the filling of the office.
- 1106. [RILEY, C. V.] To our subscribers. <Amer. Ent., December, 1860-January, 1870, v. 2, p. 93.
 - Proposed change in the character of the American Entomologist.
- 1107. [RILEY, C. V.] The Walsh entomological collection. <Amer. Ent., December, 1869-January, 1870, v. 2, pp. 93-94.
 - Statement of the conditions under which the Walsh collection of insects is to be sold; expression of preferences in regard to its disposition; extent and method of preservation of the collection.
- 1108. [RILEY, C. V.] A State entomologist for Minnesota. < Amer. Ent., December, 1869–January, 1870, v. 2, p. 94.
 - Commendation of resolutions passed by the Minnesota State Horticultural Society recommending the appointment of a state entomologist; promotion of entomological studies by appropriations from the several States.
- 1109. RILEY, C. V. [Field for the entomologist in the South.] < Amer. Ent., v. 2, December, 1869-January, 1870, p. 94; February, 1870, p. 121.
 - Extent and novelty of entomological work in the southern United States;
 J. P. Stelle at work in this field.
- 1110. [RILEY, C. V.] On our table. < Amer. Ent., December, 1869– January, 1870, v. 2, pp. 95, 96.
 - Notices of J. T. C. Ratzeburg's works on "Forest trees" and "Weeds of Germany and Switzerland."
- 1111. [RILEY, C. V.] Information wanted. <Amer. Ent., December, 1869–January, 1870, v. 2, p. 96.
 - Answer to inquiry of M. A. Kendall; habits of Xylocopa carolina [= virginica], Rhyssa [= Thalessa] lunator, and Spectrum [= Diapheromera] femorata.
- 1112. [RILEY, C. V.] The Cecropia moth, Attacus cecropia, Linn. <Amer. Ent., February, 1870, v. 2, pp. 97–102, fig. 59 [bis]–67.
 - Descriptions and figures of larva, cocoon and image of Attacus cecropia; figure of pupa; descriptions of egg and of young larva at its several stages; nomenclature, food-plants, and parasites; its value as a silk-worm; figures and descriptions of Ophion macrurum, Exorista cecropiæ n. sp., and Chalcis [= Spilochalcis] mariæ n. sp.; figure of larva of Ophion macrurum and of cocoons of Cryptus nurcius; habits of these parasites; Exorista cecropiæ considered a variety of E. militaris [= Nemoræa leucaniæ].

- 1113. RILEY, C. V. Report of the committee on entomology. Read ... before the Missouri State Horticultural Society. < Amer. Ent., February, 1870, v. 2, pp. 106-109. Reprint: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 5-8, 13-15. See No. 1127 for synopsis of contents.
- 1114. [RILEY, C. V.] Silk-worm eggs. < Amer. Ent., February, 1870, v. 2, p. 109.
 - Seventy-eight packages of silk-worm eggs, valued at \$800 per package and weighing two tons, shipped from Yokohama to France via California and the Pacific Railroad.
- 1115. RILEY, C. V. Imported insects and native American insects. <Amer. Ent., February, 1870, v. 2, pp. 110-112, fig. 72-75. Reprint: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 8-12.
 - See No. 1127 for synopsis of contents.
- 1116. [RILEY, C. V.] The trumpet grape-gall, Vitis viticola O. S. < Amer. Ent., February, 1870, v. 2, pp. 113-114, fig. 76.
 - Figure of the gall of Cecidomyia vitis-viticola. C. v.-lituus a synonym; occurrence of similar gall in England on Tilia probably caused by mites.
- 1117. [RILEY, C. V.] The goat-weed butterfly, Paphia glycerium Doubleday. < Amer. Ent., February, 1870, v. 2, pp. 121-123, figs. 81-83. Reprint, with slight changes. <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 124-128, figs. 94-96.
 - See No. 1127 for synopsis of contents.
- 1118. [RILEY, C. V.] Insects injurious to the grape-vine. No. 5. < Amer. Ent., February, 1870, v. 2, pp. 123-124, fig. 84. Reprint, with slight changes. <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 78-79, fig. 54.
 - Treats of Thyreus abbotii; see No. 1127 for synopsis of contents.
- 1119. [RILEY, C. V.] Scorpion in Kansas. < Amer. Ent., February, 1870, v. 2, p. 126.
 - Answer to inquiry of K. Kelsey; occurrence of Buthus carolinianus in Texas, Missouri, and Kansas.
- 1120. [RILEY, C. V.] The grain Bruchus of Europe just imported. <Amer. Ent., February, 1870, v. 2, pp. 126-127, fig. 85.
 - Answer to inquiry of A. S. Fnller; notice of the recent importation into New York of the European Bruchus granarius; danger of and remedy for such importation; habits and ravages of and means against the beetle in Europe as described in J. Curtis's, "Farm Insects;" figure of the imago and of infested beans; figures of larva, pupa, and imago of B. pisi and of infested pea.
- 1121. [RILEY, C. V.] Locust borer. < Amer. Ent., February, 1870, v. 2, pp. 127–128, figs. 86–89.
 - Answer to inquiry of J. M. Shaffer; habits, food-plants, seasons, and ravages of and means against Xyleutes [= Cossus] robiniæ; figures larva, pupa, and male and female imagos.

- 1122. [RILEY, C. V.] Eggs of snowy tree-cricket on raspberry canes. <Amer. Ent., February, 1870, v. 2, p. 128.
 - Answer to inquiry of J. B. Root; oviposition of and means against *Œcanthus* niveus.
- 1123. [RILEY, C. V.] Spotted rove-beetle. <Amer. Ent., February, 1870, v. 2, p. 128.
 - Answer to inquiry of J. Huggins; habits of Staphylinus maculosus and of rovebeetles in general.
- 1124. [RILEY, C. V.] Raspberry gouty gall. < Amer. Ent., February, 1870, v. 2, p. 128, fig. 90.
 - Answer to inquiry of C. Carpenter; ravages of and means against *Agrilus* ruficollis; description and figure of larva.
- 1125. [RILEY, C. V.] Parasitic cocoons. < Amer. Ent., February, 1870, v. 2, p. 128, fig. 91.
 - Answer to inquiry of G. C. Brackett; figures cocoons and conjectures as to the habits of [Microplitis ceratomiæ var. actuosus].
- 1126. [RILEY, C. V.] The pigeon Tremex in apple. <Amer. Ent., February, 1870, v. 2, p. 128.
 - Answer to inquiry of J. Huggins; habits and food-plants of Tremex.columba; description of imago; Rhyssa [= Thalessa] lunator destroys the larva.
- and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <5th Ann. Rept. State Board of Agric. for 1869, March, 1870, pp. 136 + 6, 99 figs. Separate: <Jefferson City, Mo., March, 1870, pp. 136+6, 99 figs.

CONTENTS.

5

8

NOXIOUS INSECTS.

REPORT OF THE COMMITTEE ON ENTOMOLOGY OF THE STATE HORTI-CULTURAL SOCIETY

Noxious insects less injurious in Missouri in 1869 than usual, 5—The army-worm and the grain plant-louse considerably injurious in Missouri in 1869, 5—The chinch-bug and the codling-moth less injurious, 6—A species of Thrips destroying great numbers of the Curculio, 6—Eggs of the apple-tree plant-louse destroyed by insect foes and birds, 6—According to Dr. Hull the "scab" in apples is caused by the apple-tree plant-louse, 7—The pickle-worm doing great damage during 1869, 7—Importance of preventing the introduction of injurious insects, 7—Cultivation causes insects to multiply unduly, 8—More attention paid in Europe to injurious insects than in this country, 8.

Imported insects and native American insects.

The imported currant-worm much more injurious than the native, 8—Other instances showing the greater destructiveness of imported insect enemies than of their native representatives, 9—Almost all our worst insect pests and pernicious weeds have been introduced from Europe, 10—Few American insects and plants have become naturalized in Europe, 11—The American fauna and

1127. RILEY, C. V.—Continued.

flora not so highly improved and developed as in Europe, 12—Australian fauna still more "old-fashioned" than America, 12—The parasites of injurious insects are not imported with the insects themselves, 13—Government aid should be solicited to exterminate recently imported injurious insects, 13—But little attention given so far by our Government to assist the study of economic entomology, 14—Danger of introducing the oyster-shell bark-louse into Missouri, 15—Immunity of the Pacific States from many of our fruit insects, 15.

THE CHINCH-BUG,* Blissus leucopterus

15

1 37

37

It is the most injurious of all insects infesting grain, 16-Its past history, 17; it was known in South Carolina in Revolutionary times, 17; it was injurious in Missouri as early as 1854, 17; noticed in Illinois in 1840, 17; it was very injurious in Missouri in 1868, 17; but hardly noticed in 1869, 17—Probable reason why it was not noticed in Missouri in former times, 18-Why it is not injurious in Massachusetts and New York, 18-Its natural history, 18—The pupa state in the different insect orders, 18—Time required for different insects to complete the cycle of development, 19-The chinch-bug is two-brooded in Missouri, 19-Its winter quarters, 20—Its rapid multiplication, 20—Dr. Shimer's account of its nuptial flights, 21-It deposits the eggs underground on the roots of the plant, 21—The egg, 22—Dimorphous forms of the ehinch-bug, 22—Its destructive powers, 22—Account of its appearance in immense numbers, 23-Heading off the marching bugs by a barrier of pine boards, 23—Heavy rains destructive to the chinch-bug, 24-Moistnre injurious to the egg, 24—The chinch-bug is always worse in a dry season than in a wet one, 24-Dr. Shimer's theory on epidemic disease affecting the chinch-bug, 25—Cannibal foes of the chinch-bug, 25; several species of ladybirds, 25; the weeping lacewing, 26; how the lacewing larva seizes its prey, 26; the insidious flower-bug, 27; the common quail, 28-Amount of damage done by the chinchbug, 28-Remedics, 28-Burning in winter the old corn-stalks and other dead stuff on and near the fields, 29-Mixing winter rye among spring wheat, 29-Intercepting the marching bugs by fence-boards, 29—Sowing gas-lime, 30—Other remedies, 30— Bogus chineh-bngs, 31-Several species of Heteroptera confounded with the true chinch-bug, 31—The smell emitted by the half-winged bugs, 32-The insidions flower-bug, 32-The ashgray leaf-bug and its injury to grape-vines, 33—The flea-like negro-bug, 33; injury caused by it to raspberry, strawberry, and garden flowers, 34-Two other species of negro-bug, 35-Recapitnlation of the natural history of the chinch-bug, 36.

THE ARMY-WORM, Leucania unipuncta.

Four distinct caterpillars designated as army-worms in this country, 37.

^{*} Extract in < Rept. State Board Agric., Kansas for 1873-'74, pp. 129-131.

127.	RILEY, C. V.—Continued.	
	The Southern grass-worm, Laphygma frugiperda	41
	It resembles in habits the veritable army-worm, 41.	
	The true army-worm	41
	Its past history, 41—Rev. Powers's account of its invasion in the New	
	England States in 1770, 42—Accounts of later invasions previous	
	to 1861, 43—Years of its appearance in Illinois, 43—The invasion	
	of the year 1861, 44—Its appearance in Missouri in 1869, 44—Its	
	sudden appearance and disappearance, 45—Reason for the ap-	
	parently sudden appearance, 45—Army-worm years are wet with the preceding year dry, 46—Reason for the increase and decrease	
	of the number of worms, 46—Its natural history, 47—Previous ac-	
	counts of its natural history, 47—When the eggs are laid, 47—	
	Where they are laid, 48—Misdirected instinct in insects and	
	birds, 48—Exceptions to the normal habit of the worm, 48—Color	
	of the worm, 49—The chrysalis and imago, 49—Parasites, 50;	
	the red-tailed Tachina-fly, 50; its beneficial work, 50; it in-	,
	fests also other insects, 50; Walsh's description of the fly, 51; it	
	has been redescribed as Exorista osten sackenii, 51; the yellow-	
	tailed Tachina-fly, 51; description of the fly, 51; the glassy Me-	
	sochorus, 52; the diminished Pezomachus, 52; the military	
	Microgaster, 52; the purged Ophion, 53; the army-worm Ichneumon-fly, 53—Habits of the army-worm and suggestions for	-
	its destruction, 53—Burning grass meadows in winter or early	
	spring, 54—Plowing late in the fall, 54—The marching of the	
	worms, 54—Plants they prefer, 54—They become beneficial by	
	devouring the chess in the fields, 55—Ditching, 55—Description	
	of the insect as larva and imago, 56.	
	Insects infesting the sweet-potato	56
	Tortoise-beetles	57
	The clubbed tortoise-beetle affects the Irish potato, 56—Its gen-	
	eral appearance, 57—Characteristics of tortoise-beetles, 57—	
	Merdigerous habits of tortoise-beetles and others of the same family, 58—General appearance of the larvæ, 58—Their dung	
	parasol, 59—Larval molts, 59—Egg of tortoise-beetles, 60—The	
	chrysalis, 60—Habits of and injury done by the beetles, 60—Rem-	
	edies, 61.	
	The two-striped sweet-potato beetle, Cassida bivittata	61
	It seems to be confined to that plant, 61—The larva and the use of	
	its fork, 61—Its pupa and imago, 61.	
	The golden tortoise-beetle, Coptocycla aurichalcea	62
	Food-plants and characteristics of the larva, 62—Brilliant color of	
	the beetle, 62.	·co
	The pale-thighed tortoise-beetle, Coptocycla aurichalcea.	62
	It is hardly distinguished from the foregoing species, 62. The mottled tortoise-beetle, Coptocycla guttata	63
	Characteristics of the beetle, 63; of the larva, 63.	00
	The black-legged tortoise-beetle, Cassida nigripes	63
	Characteristics of the imago and larva, 63.	
	THE PICKLE-WORM, Eudioptis nitidalis	64
	Other insects infesting cucurbitaceous yines	64
	The squash-borer, 64—It seems to be confined to the Eastern States,	
	64—The striped cucumber-beetle, 64—Injury done by the beetle,	
	64; by the larva, 65—The larva and pupa, 65—Number of annual	
	generations, 65—Remedies, 66—Extent of the injury caused by	
	it, 66—The 12-spotted Diabrotica, 66.	

1127.	RILEY, C. V.—Continued.	
	The pickle-worm	67
	Characteristics and description of the worm, 67—Its habits, 67—	
	Characteristics of the moth, 68—Accounts of injury done by the	
	worm in Missouri and Illinois, 69; in other portions of the coun-	
	try, 70—It was not known before as injurious, 70—Remedy, 70.	
	INSECTS INJURIOUS TO THE GRAPE-VINE	71
	The hog-caterpillar of the vine, Ampelophaga myron	71
	The egg, 71—Characteristics of the larva, 71—Its habits when	
	about to transform, 72—The chrysalis and imago, 72—It is one-	
	brooded North and two-brooded farther South, 72—It is very in-	
	jurious, 73—The Microgaster parasite and its development, 73—	
	Habits of caterpillar infested with the parasite, 73.	~4
	The achemon Sphinx, Philampelus achemon	74
	Appearance and habits of the larva, 74—The chrysalis, 75—The	
	insect is single-brooded, 75—The moth and its issuing from the	
	pupa-shell, 75—No parasites known, 76.	76
	The satellite Sphinx, Philampelus pandorus	70
	—Development of the larva, 76—Variations in color of the larva,	
	77—Its position when at rest, 77—The moth, 78.	
	The Abbot Sphinx, Thyreus abbotii	78
	Its distribution, 78—The larva varies much in color, 78—The chrys-	• •
	alis and imago, 79.	
	The blue caterpillars of the vine	79
	The eight-spotted forester, Alypia octomaculata, 80—Larva pre-	•
	viously mistaken for it, 80—Habits and characteristics of the	
	larva, 80—Harris's description of the larva, 81—The moth, 81—	
	Mr. Andrews's account of its ravages, 81—Remedies, 82.	
	The beautiful wood nymph, Eudryas grata, 83—Characteristics of	
	the moth, 83—Close resemblance between the larva of this and	
	the foregoing species, 83—The differences pointed out, 83—De-	
	velopment of the insect, 83.	
	The pearl wood nymph, Eudryas unio, 83—It greatly resembles the	
	beautiful wood nymph, 83—Its probable larva, 84—Practical im-	
	portance of distinguishing these closely allied species, 84.	
	The American Procris, Harrisina americana	85
	Work of its larva, 85—Description of full-grown larva, 86—The	
	moth, 86—It is not very destructive, 86—Two annual broods of	
	the insects, 86—Parasite of the American Procris, 87.	_
	The new grape-root borer, Prionus laticollis	87
	Correction of opinion formerly expressed, 87.	
-	The broad-necked Prionus, 87—Duration of the larva state, 87—	
	Its transformation, 88—It bores also in apple roots, 88—Great	
	damage done by the borer, 88—No good remedy known, 88.	
	The tile-horned Prionus, Prionus imbricornis, 89—How it differs	
	from the foregoing, 89—Its occurrence on prairie land, 90—Small	
	dimorphous male form, 90—The larva subsists also upon the	
	roots of herbaceous plants, 90—Practical considerations, 91.	92
	The grape seed magget, Isosoma vitis	04
	The grape-seed Curculio larva of the first report is that of a hymen-	
	opterous insect, 92—The perfect insect is closely allied to the joint-worm fly, 92—Mr. Saunders' account and description of the	•
	joint-worm ny, 92—Mr. Saunders' account and description of the	

1127. RILEY, C. V.—Continued.	
The canker-worm, Paleacrita vernata; Anisopteryx pometaria The eggs, 94—The larva and larval changes, 95—Importance for	94
the orchardists to recognize the true canker-worm, 95—The imported elm leaf-beetle mistaken for it, 95—Description of the larva, 96—Its food-plants, 96—The chrysalis, 96—Only one annual brood in the latitude of St. Louis, 97—The moth and its	,
varieties, 97—It is less injurious in Missouri than in the Eastern States, 97—Remedies, 98—Classification of remedies proposed, 98—The trough and bandage systems, 99—Muriate of lime as remedy, 100—Jarring the tree, 101—Late fall plowing, 101—	
Summer plowing, 102—Efficiency of hogs, 102—Enemies, 102; birds, mites, and parasites, 102; ground-beetles, 103; the fraternal potter-wasp, 103.	
Cabbage-worms	104
The Southern cabbage butterfly, Pieris protodice. Its geographical range, 104—Injury caused by it in Missouri, 104	104
—Description of the larva, 105—The chrysalis and imago, 105—Habits and other food-plants, 105.	
The potherb butterfly, Pieris olearacea, 105—It is a Northern species, 105—It will very likely never occur in Missouri, 106—Geo-	
graphical range of insects principally influenced by temperature, 106—Isentomic lines, 106—Southern insects found near St. Louis, 106.	
The imported cabbage butterfly, Pieris rapa, 106—Amount of damage caused by it in Canada, 107—Its spread westward, 107—It will undoubtedly spread to St. Louis, 107—The insect in Eu-	
rope, 107—History of its introduction, 107—The insect in its different stages, 108—Its food, 108—Remedies, 109—Parasites, 109. The cabbage Plusia, Plusia brassica	110
Characteristics and habits of the larva, 110—Its transformations, 111—Remedics, 111—Description of larva, chrysalis, and imago, 111—A similar worm occurring on thistles, 112.	110
The zebra caterpillar, Mamestra picta Habits and characteristics of the larva, 112—The chrysalis and the moth, 113—Two annual broods, 113—Food-plants, 113.	112
THE TARNISHED PLANT-BUG, Lygus pratensis Injury caused by it to various trees and plants, 114—It is a very variable species, 114—Its development, 114—No effective remedy	113
known, 115—Preventive measures, 115.	110
THE PHILENOR SWALLOW-TAIL, Papilio philenor Its food-plant, 116—Damage done by it, 116—Characteristics and development of the larva, 116—Description of the larva, 117—The pupa, 117—The imago, 117—Prevention, 118.	116
THE COTTONWOOD DAGGER, Acronycta lepusculina General appearance of the larva, 119—Two annual broods, 119— Chrysalis and moth, 119—Larvæ of other species belonging to the genus Acronycta, 119—Parasites, 120—Description of larva and imago, 120—Characters and habits of other species of the	119
same genus, 121.	
The Missouri Bee-Killer, Proctacanthus milbertii The true scientific name of the Nebraska bee-killer, 121—Wing-veins of the genera Asilus, Promachus, and Erax, 122—Description of the Missouri bee-killer, 122—How to destroy the flies, 123—Habits and life-history of Asilus-flies, 123—Description of larva and pupa	121

of Erax bastardi (?), 124-Synonymical notes on the imago, 124.

INNOXIOUS INSECTS.

THE GOAT-WEED BUTTERFLY, Paphia glycerium	125
Its geographical distribution and position in classification, 125—Its	
food-plant, 125—Habits of the larva, 126—Larval changes, 126—	
Conformity in the color of the larva with that of the leaves,	
127—Description of the full-grown larva, 127—Transformation	
of the larva to chrysalis, 127—The two sexes of the imago, 127—	
Hibernation, 128.	
THE BLACK BREEZE-FLY, Tabanus atratus	128
Breeze-flies beneficial in the larva state, 128—Tormenting power	
of breeze-flies, 128—Their mode of flight, 129—Our knowledge	
of their larval character and habits, 129—General characters of	
the larva of the black breeze-fly, 129—It is semi-aquatic, 129—	
Walsh's description of the larva, 130—Habits and food of the	
larva, 130—Its transformations, 131—Description of the pupa,	
131—Probable habits of breeze-fly larvæ on the Western prai-	
ries, 132.	
GALLS MADE BY MOTHS.	132
THE FALSE INDIGO GALL-MOTH, walshia amorphella	132
The gall and its structure, 132—General appearance of the larva	
and the moth, 133—Description of larva and imago, 133.	
THE MISNAMED GALL-MOTH, Euryptychia saligneana	134
Is it a true gall-maker or an inquiline?, 134—Walsh's description	
of the larva, 134—Description of the imago, 134—Generic char-	
geters 134_Reasons why the insect is an intruder and not a goll-	
maker, 134—Enumeration of the known gall-making moths, 135—	
How the gall is formed, 135.	

1128. [RILEY, C. V.] Mr. Walsh's portrait. <Amer. Ent., March, 1870, v. 2, p. 129.

Remarks accompanying the portrait of B. D. Walsh; resolutions passed on the death of Walsh by the London branch of the Eutomological Society of Ontario; by the American Entomological Society, the Illinois State Horticultural Society, and by the Kausas State Horticultural Society.

1129. RILEY, C. V. The plum Curculio, Conotrachelus nenuphar Herbst. <Amer. Ent., March, 1870, v. 2, pp. 130-137, fig. 92.

Paper read before the Illinois State Horticultural Society at its 14th annual meeting; summary of established facts and discussion of mooted points in the life-history of Conotrachelus nenuphar; seasons, habits, transformations, food-plants, enemies of and means against the same; descriptions and figures of larva and imago; figure of pupa; hibernation and the effects of climate on the prolongation of the life of insects.

1130. RILEY, C. V. Insects injurious to the grape-vine. No. 6. <Amer. Ent., March, 1870, v. 2, pp. 150-153, figs. 100-102. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 80-82, fig. 55.

Treats of Alypia octomaculata; see No. 1127 for synopsis of contents.

1131. [RILEY, C. V.] Pithy blackberry gall. <Amer. Ent., March, 1870, v. 2, pp. 159-160, fig. 103.

Answer to inquiry of C. W. ["S. C."] Spaulding; descriptions and figures of larva and gall of Diastrophus nebulosus; figure of pupa; seasons, foodplant, guest-fly [Periclistus sylvestris], and parasite [Eurytoma diastrophi] and means against this species; the genus Diastrophus confined to Rosaceæ, Cynips to Cupuliferæ and Antistrophus to Compositæ.

1132. [RILEY, C. V.] Clover-worms. < Amer. Ent., March, 1870, v. 2, p. 160.

Answer to inquiry of G. Pauls; geographical distribution, food-habits, vernacular names and synonymy of and means against Asopia costalis.

1133. [RILEY, C. V.] Seed-ticks under bark of apple-trees. < Amer. Ent., March, 1870, v. 2, p. 160.

Answer to inquiry of O. B. Galusha; character of insects as regards the number of legs; occurrence of *Ixodes unipunctata* under outer bark of apple-trees at Morris, Ill.

1134. [RILEY, C. V.] Parasitic cocoons. <Amer. Ent., March, 1870, v. 2, p. 160.

Answer to inquiry of S. W. Beckworth; occurrence of a mass of cocoons of [Microplitis ceratomiæ var. actuosus] under red-oak at South Pass, Ill.

1135. [RILEY, C. V.] Is any knowledge useless? < Amer. Ent. and Bot., April, 1870, v. 2, pp. 164–166.

keprint of article from Manufacturer and Builder, November, 1869; minute investigations in science may be of great practical importance; cases in which a knowledge of life-history of Galeruca calmariensis [=xanthomelæna], Conotrachelus nenuphar, and Lymexylon navale was or might have been of great value.

1136. [RILEY, C. V.] Tomato fruit-worm. < Amer. Ent. and Bot., April, 1870, v. 2, p. 172.

Notice of statement by J. J. Weir that *Heliothis armigera* was bred from larvæ which fed on the fruit of tomato in England; food-plants of this insect.

1137. RILEY, C. V. Insects injurious to the grape-vine. No. 7. < Amer. Ent. and Bot., April, 1870, v. 2, pp. 173–174, figs. 107–108. Reprinted, with slight changes, from < 2d Ann. Rept. State Ent. Mo., March, 1870, pp. 85–87, figs. 58–59.

Treats of Procris [=Harrisina] americana; see No. 1127 for synopsis of contents.

1138. [RILEY, C. V.] The death web of young trout. <Amer. Ent. and Bot., April, 1870, v. 2, p. 174.

Reprint, with review, of Seth Green's "An enemy to young trout;" young trout and young white-fish killed by the web of an unknown aquatic larva, probably one of the caddis-flies; habits of the larvæ of *Phryganeidæ*. See No. 1160.

1139. [RILEY, C. V.] "Scab" in apple v. apple-tree plant-lice. < Amer. Ent. and Bot., April, 1870, v. 2, p. 178.

Notes the freedom of apple-trees in 1870 from the eggs of Aphis mali, and the opportunity to test the question of the connection of these insects with the "scab."

- 1140. [RILEY, C. V.] Insects named. <Amer. Ent. and Bot., April, 1870, v. 2, p. 179.
 - Answer to inquiry of M. Hobart; identification of several insects; food-plants of Grapta comma, Geometra [=Zerene] catenaria, Serica vespertina, and Tetraopes 5-maculatus; geographical distribution of Grapta comma; description of larva of Ecpantheria scribonia.
- 1141. [RILEY, C. V.] Supposed trout enemy. < Amer. Ent. and Bot., April, 1870, v. 2, pp. 179–180.
 - Answer to inquiry of F. Mather; identification of several insects; habits and occurrence of *Capnia minima*; occurrence of *Piophila casci* and *Ptinus brunneus* at Honeoye Falls, N. Y.
- 1142. [RILEY, C. V.] Food for trout. < Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of Seth Green and Collins; with knowledge of the character of the worms which forms a desirable food for young trout, it may be possible to suggest some method of propagating the worms artificially.
- 1143. [RILEY, C. V.] Hair-snakes. < Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of E. W. M.; parasitic habits, abodes and prolificacy, and classificatory relations of *Gordius varius* and *G. aquaticus*.
- 1144. [RILEY, C. V.] Egg-sack of some unknown spider. <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.

 Answer to inquiry of A. Engelmann; figure of egg-sack of *Epeira* sp.?
- 1145. [RILEY, C. V.] Do worker bees sting the drones to death? <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of M. W. V.; there is no good reason to donbt that worker bees sting the drones to death when the mission of the latter is ended.
- 1146. [RILEY, C. V.] Red spider. < Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of R. H. Warder; Trombidium [= Tetranychus] telarius imported from Europe; size, color, and abodes of and means against the same.
- 1147. [RILEY, C. V.] Insect named. April, 1870, v. 2, p. 180.
 - Answer to inquiry of M. Barrett; food-habits of Psocus venosus and other Psocidæ.
- 1148. [RILEY, C. V.] To destroy plant-lice. <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of B. F. Lazear; means against Aphidida on house plants.
- 1149. [RILEY, C. V.] Raspberry root-gall. <Amer. Ent. and Bot., April, 1870, v. 2, p. 181, fig. 110.
 - Answer to inquiry of K. Parsons; description and figure of gall of *Rhodites* radicum occurring on roots of Rosaccæ; genera of parasites raised from it; interest of the question of the manner and extent of parasitization of this gall.
- 1150. [RILEY, C. V.] Spined slug-worm. <Amer. Ent. and Bot., April, 1870, v. 2, p. 181.
 - Answer to inquiry of L. G. Saffer and A. R. Bodley; brief description of the larva of *Limacodes* sp.

- 1151. [RILEY, C. V.] Apple-tree insects. < Amer. Ent. and Bot., April, 1870, v. 2, p. 181.
 - Answer to inquiry of L. Camfield; identification of specimens and directions for the destruction of Orgyia leucostigma and Phycita nebulo [= Acrobasis indiginella].
- 1152. [RILEY, C. V.] Native apple-tree bark-lice. <Amer. Ent. and Bot., April, 1870, v. 2, p. 181, fig. 111.
 - Answer to inquiry of A. C. Hammond; food-plants of and means against Diaspis harrisii [= Chionaspis furfurus]; figure of infested twig of apple; the presence of enemies and parasites prevent this species becoming a formidable pest.
- 1153. [RILEY, C. V.] The hedge-hog caterpillar. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182, fig. 112.
 - Answer to inquiry of H. Burt; descriptions and figures of larva and imago of Arctia [= Pyrrharctia] isabella; figure of pupa and cocoon; habits and hibernation; its larva and that of Ecpantheria scribonia called "feverworm," and ignorantly supposed to cause malarial fevers; food-habits of Horinus [= Merinus] lavis.
- 1154. [RILEY, C. V.] Chick-weed Geometer. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of J. Huggins; identification of specimens; reference to account of *Hamatopis grataria*; *Cermatia forceps* common in houses in the latitude of St. Louis, Mo.; the sulphur remedy for canker-worms absurd.
- 1155. [RILEY, C. V.] Bean-weevil. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of G. W. Copley; specimens identified; occurrence and ravages of *Bruchus obsoletus* in Illinois.
- 1156. [RILEY, C. V.] Bag-worm at South Pass, Ill. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of G. H. Baker; occurrence of Thyridopteryx ephemeræformis at South Pass, Ill.; importance of its destruction.
- 1157. [RILEY, C. V.] Eggs of oblong-winged katy-did. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of E. D. Ladd; differences between the eggs of *Phylloptera* [= Amblycorypha] oblongifolia and those of *Platyphyllum* [= Cyrtophyllus] concavus; the former occur on current and various trees.
- 1158. [RILEY, C. V.] Insects injurious to the grape-vine. No. 8. <Amer. Ent. and Bot., May, 1870, v. 2, pp. 208-209, fig. 127. Reprint, with slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 61-63, fig. 24.
 - Treats of Desmia maculalis; see No. 1301 for synopsis of contents.
- 1159. [RILEY, C. V.] The periodical Cicada alias the 17-year and 13-year locust. <Amer. Ent. and Bot., May, 1870, p. 211.
 - Quotes, from the 1st Ann. Rept. State Ent. Mo., the localities in which *Cicada* [= Tibicen] septendecim and C. [= T.] tredecim will appear in 1870, with request for reports of the occurrence of these insects.
- 1160. [RILEY, C. V.] The death-web of young trout. < Amer. Ent. and Bot., May, 1870, v. 2, p. 211.
 - Supplementary to No. 1138; the larvæ mentioned belong to the genus Simulium.

- 1161. [RILEY, C. V.] Worms under mulch hay. <Amer. Ent. and Bot., May, 1870, v. 2, p. 212.
 - Answer to inquiry of J. F. Flagg; description, season, habits, and ravages of and means against the larva of *Tipula* sp.
- 1162. [RILEY, C. V.] A new pear-tree insect. <Amer. Ent. and Bot., May, 1870, v. 2, p. 212, fig. 129.
 - Answer to inquiry of E. J. Ayres; food-habits and ravages of and means against *Platycerus quercus*; figure of imago.
- 1163. [RILEY, C. V.] Apple-twig borer. < Amer. Ent. and Bot., May, 1870, v. 2, p. 212.
 - Answer to inquiry of J. B. Myers; Bostrichus [= Amphicerus] bicaudatus bores into the axil of limbs of pear-trees.
- 1164. [RILEY, C. V.] Cocoons of polyphemus moth. < Amer. Ent. and Bot., May, 1870, v. 2, 212.
 - Answer to inquiry of H. J. Dunlap; cocoon of Attacus [= Telea] polyphemus found on Morello cherry-tree.
- 1165. [RILEY, C. V.] Galls on supposed dock. <Amer. Ent. and Bot., May, 1870, v. 2, p. 212.
 - Answer to inquiry of S. V. Summers; Gelechia gallæ-solidaginis forms galls on stems of Solidago; Gastrophysa [= Gastroidea] cyanea breeds on Rumex.
- 1166. [RILEY, C. V.] Mossy rose-gall. <Amer. Ent. and Bot., May, 1870, v. 2, p. 213, fig. 130.
 - Answer to inquiry of W. M. Locke; description of gall, larva, and imago of *Rhodites rosæ*; figure of the gall; description of a parasitic larva.
- 1167. [RILEY, C. V.] Punctures on the rose-twig. <Amer. Ent. and Bot., May, 1870, v. 2, p. 213, fig. 131.
 - Answer to inquiry of G. W. Copley; punctures in rose-stem; eggs, probably of a cricket, imbedded in pith of the same; description of the egg and figure of injured stem.
- 1168. [RILEY, C. V.] Snout-beetle. < Amer. Ent. and Bot., May, 1870, v. 2, p. 213.
 - Answer to inquiry of Mary Treat; Hylobius confusus a timber borer and usually in pinc.
- 1169. [RILEY, C. V.] The oyster-shell bark-louse in Missouri. <Amer. Ent. and Bot., May, 1870, v. 2, pp. 213–214, fig. 132.
 - Answer to inquiry of B. P. Hanan; occurrence of Aspidiotus conchiformis [=Mytilaspis pomorum] in Missouri; importance of the thorough extirpation of the insect; figure of a piece of bark infested by the same.
- 1170. [RILEY, C. V.] The pod-like willow-gall. <Amer. Ent. and Bot., May, 1870, v. 2, p. 214, fig. 133.
 - Answer to inquiry of J. R. Muhleman; description and figure of the gall of *Cecidomyia salicis-siliqua*; figure of larva; food-plants, synonymy, and description of the pupa of the same.
- 1171. [RILEY, C. V.] Bee-nest. < Amer. Ent. and Bot., 1870, v. 2; May, p. 214, fig. 134; September, p. 307.
 - Answer to inquiry of J. R. Muhleman; description and figure of larva of *Prosopis affinis* in hot, low currant-stem; probability that *Ceratina dupla* breeds twice a year.

- 1172. [RILEY, C. V.] Beetles named. < Amer. Ent. and Bot., May, 1870, v. 2, p. 214.
 - Answer to inquiry of S. V. Summers; specimens identified; difference between Canthon lavis and C. chalcites.
- 1173. [RILEY, C. V.] Great discovery: Curculio extermination possible! <Amer. Ent. and Bot., June, 1870, v. 2, pp. 225–227.

 Notice: <Cultivator and Country Gentleman, 9 June, 1870, v. 35, p. 361.
 - Reprint of articles by J. E. Chamberlain and Mrs. H. Wier on the destruction of the Curculio; letter from W. B. Ransom; criticism of the same; means against Conotrachelus nenuphar; distinctions between C. nenuphar and Anthonomus quadrigibbus.
- 1174. [RILEY, C. V.] The death-web of young trout. < Amer. Ent. and Bot., June, 1870, v. 2, pp. 227–228, figs. 143–144.
 - Value and progress of fish-culture; habits of Simulium piscicidium, with the observations of Seth Green and Sara J. McBride; figures larva and pupa of S. piscicidium and the imago of S. molestum.
- 1175. [RILEY, C. V.] Insects injurious to the grape-vine. No. 9. < Amer. Ent. and Bot., June, 1870, v. 2, pp. 234–235, fig. 148. Reprint, with slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 65–68, fig. 27.
 - Treats of Pterophorus [= Oxyptilus] periscelidactylus; see No. 1301 for synopsis of contents.
- 1176. [RILEY, C. V.] The apple Curculio. <Amer. Ent. and Bot., June, 1870, v. 2, p. 243, fig. 152. Figures of the image of Anthonomus quadrigibbus.
- 1177. [RILEY, C. V.] The new Curculio remedy. <Amer. Ent. and Bot., June, 1870, v. 2, p. 243.
 - Results of experiments in the use of Ransom chip-trap for Conotrachelus nenuphar.
- 1178. [RILEY, C. V.] Tarantula of Texas. < Amer. Ent. and Bot., June, 1870, v. 2, p. 244.
 - Answer to inquiry of L. J. Stroop; the figure of Mygale hentzii given in volume one is somewhat incorrect.
- 1179. [RILEY, C. V.] Ailanthus silk-worm naturalized. < Amer. Ent. and Bot., June, 1870, v. 2, p. 244.
 - Answer to inquiry of A. S. Fuller; introduction and naturalization of Attacus cynthia.
- 1180. [RILEY, C. V.] Cypress-gall. <Amer. Ent. and Bot., June, 1870, v. 2, p. 244, fig. 153.
 - Answer to inquiry of J. P. Stelle; description and figures of gall and description of gall and image of *Cecidomyia cupressi-ananassa* n. sp. on cypress; figures breast-bone of the larva.
- 1181. [RILEY, C. V.] Tent-caterpillar of the forest. <Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to the inquiries of A. M. Brown and J. H. Evans; habits, food-plants, and ravages of Clisiocampa sylvatica [=disstria] and C. americana.

- 1182. [RILEY, C. V.] Worm-boring into peach. <Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiries of W. C. Flagg, A. C. Hammond, and M. M. Hooton; food-plants and description of larva of Xylina cinerea [=Lithophane antennata].
- 1183. [RILEY, C. V.] Insects named. <Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiry of A. Engelman; descriptions of Smilia auriculata and Membracis ampelopsidis; both species common on grape-vines.
- 1184. [RILEY, C. V.] Lice on "snow-balls. < Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiry of Mrs. C. L. Seymour; means against Aphidida.
- 1185. [RILEY, C. V.] Twig-borer. < Amer. Ent. and Bot., June, 1870, v. 2 p. 245, fig. 154.
 - Answer to inquiries of S. H. Kriedelbaugh and G. F. Merriam; figures of male and female Bostrichus [=Amphicerus] bicaudatus, which bore into the axils of grape-buds.
- 1186. [RILEY, C. V.] Bee enemy. < Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiry of F. Brewer; Staphylinus maculosus, found eating a bee, is rather a scavenger than an insect of prey.
- 1187. [RILEY, C. V.] Knots on apple-tree roots caused by root-lice. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of B. N. McKinstry; ravages of and means against Schizoneura lanigera on roots of young apple-trees.
- 1188. [RILEY, C. V.] Beetles named. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of W. W. Daniells; Ithycerus noveboracensis injures appleleaves.
- 1189. [RILEY, C. V.] Bag-worm. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of M. M. Hooten; occurrence of young larva of *Thyridopteryx ephemeraformis* on peach-trees; manner in which they carry their cases.
- 1190. [RILEY, C. V.] The larder-beetle. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of S. H. Kriedelbaugh; food-habits and description of the larva and image of *Dermestes lardarius*.
- 1191. [RILEY, C. V.] Water-bug. Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of W. H. Harrington; habits and description of Ranatra fusca.
- 1192. [RILEY, C. V.] Gregarious worms on horse-chestnut. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of W. R. Howard; oviposition, food-plant, and parasite of *Tortrix* [= Cacoecia] rileyana.
- 1193. [RILEY, C. V.] Pupa of the disippus butterfly. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246, fig. 155.
 - Answer to inquiry of T. Montgomery; figure and colors of pupa of Limenitis disippus; food-plants and hibernations of larva.

- 1194. [RILEY, C. V.] Prickly-rose gall. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of J. Cochrane and J. P. Stelle; brief description of the gall of Rhodites bicolor on wild rose.
- 1195. [Riley, C. V.] Insects feeding on sap of black-walnut. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.

 Answer to inquiry of M. Barrett; food-habits of Psocus venosus.
- 1196. [RILEY, C. V.] Locust-borer. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.

Answer to inquiry of "Arbor;" means against Arhopalus [= Cyllene] robinia.

- 1197. [RILEY, C. V.] To exterminate cockroaches. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.

 Answer to inquiry of R. F. Weitbree; means against cockroaches.
- 1198. [RILEY, C. V.] The white-lined morning Sphinx. (Deilephila lineata, Fabr.) < Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 257-258, figs. 162-164. Reprint, with additions and slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 140-142, figs. 60-62.

Treats of Deilephila lineata; see No. 1301 for synopsis of contents.

- 1199. [RILEY, C. V.] Descriptive entomology. <Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 258-261. Abstract: <Cultivator and Country Gentleman, 6 April, 1871, v. 36, p. 218.
 - Critical review of a remark by J. A. Lintner; calculation of the cost, labor, and extent of a work containing the description and figure of every existing species of insect.
- 1200. [RILEY, C. V.] The tent-caterpillar of the forest. (Clisiocampa sylvatica, Harr.) < Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 261–266, figs. 165–168. Reprint, with additions and slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 121–129, figs. 52–54.

Treats of Clisiocampa sylvatica [= disstria]; see No. 1301 for synopsis of contents.

- 1201. [RILEY, C. V.] The Ransom Curculio remedy. <Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 268–271.
 - Discussion of the efficacy of the Ransom chip-trap against *Conotrachelus* nenuphar; extracts from and criticism of articles by E. S. Hull, W. B. Ransom, and others on this subject.
- 1202. [Riley, C. V.] Insects injurious to the grape-vine. No. 10. <Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 272-273, fig. 170. Reprint, with additions and slight changes. <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 68-69, fig. 28. Treats of Spilosoma virginica; see No. 1301 for synopsis of contents.
- 1203. [RILEY, C. V.] The Walsh entomological cabinet. < Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.

Remarks on the purchase and disposal of the entomological collection of B. D. Walsh.

- 1204. [RILEY, C. V.] The currant-worm! < Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.
 - Comments upon the confusing way in which some horticultural publications treat of the current-worm, without discrimination of species.
- 1205. [RILEY, C. V.] Water larva. < Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.
 - Answer to inquiry of F. Mather; habits of the larvæ of Ephemeridæ.
- 1206. [RILEY, C. V.] Large black potato-beetles. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.
 - Answer to inquiry of R. S. Elliott; Epicauta corvina? injuring potato-vines in Kansas.
- 1207. [RILEY, C. V.] Destroying cherry plant-lice. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.

 Answer to inquiry of G. C. Brackett; means against Myzus cerasi.
- 1208. [RILEY, C. V.] Caterpillars on grape-vines. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.
 - Answer to inquiry of G. A. Watson; larvæ of Alypia octomaculata and Acronycta oblinita found on grape-vines; food-plants of the latter species.
- 1209. [RILEY, C. V.] Ash-gray blister-beetle. <Amer. Ent. and Bot.,
 July-August, 1870, v. 2, p. 275.
 - Answer to inquiry of P. H. Foster; Lytta cinerea [= Macrobasis unicolor] feeding on the three-thorned Acacia.
- 1210. [RILEY, C. V.] Specimens lost. < Amer. Ent. and Bot., July-August, 1870; v. 2, p. 276.
 - Answer to inquiry of C. H. Roberts; food-habits and description of larva of Gortyna nitela and of an undetermined moth on peach-trees.
- 1211. [RILEY, C. V.] White willow worm. < Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer to inquiry of S. H. K.; food-plants and description of larva of and means against Nematus ventralis.
- 1212. [RILEY, C. V.] Bark-lice on grape-vine and raspberry saw-fly. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Auswer to inquiry of S. Thompson; Lecanium [= Pulvinaria] vitis common on grape-vines in Europe and North America; its oviposition and occurrence in Illinois; description and means against the larva of Selandria [= Monophadnus] rubi.
- 1213. [RILEY, C. V.] Apple-tree borer; variations in the two-striped Saperda. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer to inquiry of D. B. Wier; colorational variations in Saperda bivittata [=candida]; abundance and ravages of Capsus oblineatus [=Lygus pratensis].
- 1214. [RILEY, C. V.] The plum Curculio breeds in apple. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer inquiry of E. Leming; Conotrachelus nenuphar breeds in the fruit of apple.

- 1215. [RILEY, C. V.] Cecropia worm. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer to inquiry of J. F. Thompson; occurrence and ravages of the larva of Attacus cecropia on apple-trees.
- 1216. [RILEY, C. V.] Gigantic rhinoceros beetle. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.

Answer to inquiry of L. G. Saffer; variations in coloration of *Dynastes tityus*.

- 1217. [Riley, C. V.] Roman-nosed pupa. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer to inquiry of E. D. Van Winkle; food-habits of *Limenitis ursula* and L. disippus; the pupe of the two species are alike.
- 1218. [RILEY, C. V.] The onward march of the Colorado potato-beetle. A word to our Canadian neighbors. <Amer. Ent. and Bot., September, 1870, v. 2, pp. 289-291, fig. 181.
 - Remarks on the spread of *Doryphora decemlineata* into Ontario, and means of checking it; efficacy and proper use of Paris green; other remedies; *Lebia grandis* a natural enemy of the larvæ.
- 1219. [RILEY, C. V.] The tarnished plant-bug. (Capsus oblineatus, Say). <Amer. Ent. and Bot., September, 1870, v. 2, pp. 291–293, fig. 182. Reprinted, with additions and slight changes, from <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 113–115, fig. 83.

Treats of Capsus oblineatus [=Lygus pratensis]; see No. 1127 for synopsis of contents.

- 1220. [RILEY, C. V.] Osage orange for the mulberry silk-worm. Amer. Ent. and Bot., September, 1870, v. 2, p. 293.
 - Reprint and critical review of S. Cornaby's article on the above; important' articles should not be published anonymously.
- 1221. [RILEY, C. V.] Insects injurious to the grape-vine. No. 11. <Amer. Ent. and Bot., September, 1870, v. 2, p. 295, fig. 185. Reprint with slight changes. <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 77–79, fig. 34.

Treats of Pelidnota punetata; see No. 1301 for synopsis of contents.

- 1222. [RILEY, C. V.] The slug on pear and cherry trees. < Amer. Ent. and Bot., September, 1870, v. 2, p. 296.
 - Reprint and critical review of "Addi" on the above; ravages of and means against Selandria [= Eriocampa] cerasi.
- 1223. [RILEY, C. V.] Appendix to joint-worm article published in vol. 1, No. 8 < Amer. Ent. and Bot., September, 1870, v. 2, pp. 296–297.
 - Introduction to and explanation of the preparation and publication of Walsh's Eurytomides. See No. 384.
- 1224. [RILEY, C. V.] Entomology indeed run mad! <Amer. Ent. and Bot., September, 1870, v. 2, p. 305.
 - Critical review of Mark Miller's article on the currant-worm; Nematus ventrieosus [=ribesii] confounded with Eufitchia ribearia.
- 1225. [RILEY, C. V.] Red spider. < Amer. Ent. and Bot., September, 1870, v. 2, p. 305.
 - Ravages and transformations of Tetranychus telarius.

- 1226. [RILEY, C. V.] Insects named. < Amer. Ent. and Bot., September, 1870, v. 2, p. 306.
 - Answer to inquiry of J. K. Kidd; food-habits of Silpha peltata [= americana] and allied forms, of Calosoma scrutator and of Laphria [= Dasyllis] thoracica; mimicry between Laphria and Bombus and its use.
- 1227. [RILEY, C. V.] Caterpillar of white-marked tussock moth. <Amer. Ent. and Bot., September, 1870, v. 2, p. 306, fig. 186.
 - Answer to inquiry of G. C. Brackett; figure of larva and description of image of Orgyia leucostigma; posture of male in repose; habits of female; Saperda bivittata [= candida] usually perishes if it has not changed to a pupa before the death of the tree; Chrysobothris femorata lives for weeks on dead wood.
 - 1228. [RILEY, C. V.] Does the apple Curculio go underground to transform? <Amer. Ent. and Bot., September, 1870, v. 2, p. 306.
 - Answer to inquiry of W. Muir; Anthonomus quadrigibbus transforms within the fruit; it does not attack stone fruit.
 - 1229. [RILEY, C. V.] Walnut caterpillars. <Amer. Ent. and Bot., September, 1870, v. 2, p. 306.
 - Answer to inquiry of G. M. Levette; seasons, habits, and food-plant of and means against *Datana ministra*; description of larva.
 - 1230. [RILEY, C. V.] Striped blister-beetle. <Amer. Ent. and Bot., September, 1870, v. 2, p. 306, fig. 187.
 - Answer to inquiry of A. Galt; means against Epicauta vittata and other Meloida on potato vines; figure of Epicauta vittata.
 - 1231. [RILEY, C. V.] Grape-vine Fidia. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307, fig. 188.
 - Answer to inquiry of J. Hetzel; habits, ravages, and food-plants of and means against Fidia viticida; figure of this species.
 - 1232. [RILEY, C. V.] Some interesting insects. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of A. S. Fuller; ravages of Bruchus obsoletus in Lima beans; of Butalis cerealella in flint corn in stems and of Lobesia [=Endemis] botrana in blossoms of blackberry; description of the larva case of Phycita nebulo [=Acrobasis indiginella] and of galls of Cccidomyia tubicola on Carya.
 - 1233. [RILEY, C. V.] The green hag-moth. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of S. B. Shaw; food-plants, and synonymy of Callochlora viridis [=Parasa chloris]; description of larva and imago; the larva described by Reakirt does not belong to this species.
 - 1234. [RILEY, C. V.] The antiopa butterfly. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of A. S. Moss; description of larva and image of *Vanessa antiopa*; the larva feeds on willow; vernacular name and comparative abundance.
 - 1235. [RILEY, C. V.] Rose-gall and pupa of archippus butterfly. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307, fig. 189.
 - Answer to inquiry of L. B. Custar; figure of pupa of Danais archippus; descriptions of three undetermined galls on rose-leaf formed by Rhodites sp.?

 10 ENT

- 1236. [RILEY, C. V.] White grubs in strawberry beds. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of J. B. Miller; food-habits of larva and description of imago of Cyclocephala immaculata.
- 1237. [RILEY, C. V.] Larva of the thoas swallow-tail. < Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of E. H. Sprague; food-plant and description of larva of Papilio thoas [=cresphontes]; the larva rare in Missouri.
- 1238. [RILEY, C. V.] Larva of clubbed tortoise-beetle. < Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of A. R. Bodley; Cassida, Coptocycla and Deloyala feed on Convolvulacew with the exception of Deloyala [=Coptocycla] elavata which feeds on Solanacew.
- 1239. [RILEY, C. V.] The banded Ips in calyx of pear. <Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of G. C. Brackett; food-habits and description of *Ips* fasciatus.
- 1240. [RILEY, C. V.] The larder-beetle. <Amer. Ent. and Bot., September, 1870, v. 2, p. 308, fig. 191.
 - Answer to inquiry of F. S. Sleeper; figures of larva and image and of magnified hair of larva of *Dermestes lardarius*; ravages in collections of preserved animals.
- 1241. [RILEY, C. V.] Moth named. < Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of E. M. Hale; description of Ctenucha latreillana [=vir-ginica]; its abundance in 1870 in the vicinity of St. Louis, Mo.
- 1242. [RILEY, C. V.] The little Cicada. <Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of G. O. Hardeman; occurrence of Cicada [=Melampsalta] parvula in Missouri.
- 1243. [RILEY, C. V.] The brown mantispian. <Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of G. C. Bracket; Mantispa brunnea common, predaceous, and beneficial.
- 1244. [RILEY, C. V.] Small reddish snont-beetle on apple. <Amer. Ent. and Bot., September, 1870, v. 2, p. 30.
 - Answer to inquiry of J. Weed; food-habits and description of Anthonomus cratægi.
- 1245. [RILEY, C. V.] Prickly rose-gall. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309, fig. 192.
 - Answer to inquiry of "Subscriber;" description and figure of galls and description of Rhodites bicolor.
- 1246. [RILEY, C. V.] Questions answered. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiries of K. Parsons; formation and use of portable cases by *Tincina*; habits and character of the young of *Mytilaspis pomicorticis* [=pomorum]; food-habits of ants.

- 1247. [RILEY, C. V.] Hog-caterpillar of the vine infested with parasites. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of J. M. Wilson; occurrence of larva of Darapsa [=Am-pelophaga] myron parasitized by Apanteles congregatus.
- 1248. [RILEY, C. V.] Larva of Abbot Sphinx. < Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of S. E. Todd; occurrence of larva of *Thyreus abbotii* on grape-vine; means against the same.
- **1249.** [RILEY, C. V.] Cecropia worm. < Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of E. G. Hofman; occurrence of Attacus cecropia on plum.
- 1250. [RILEY, C. V.] Flat-headed borer in soft inaples. < Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of L. R. Elliott; ravages of and means against Chryso-bothris femorata on soft maple; ravages of Arhopalus [= Cyllene] robinia on black-locust.
- 1251. [RILEY, C. V.] Cherry plant-lice and their foes. < Amer. Ent. and Bot., September, 1870, v. 2, p. 309, fig. 193.
 - Answer to inquiry of C. H. Roberts; occurrence on cherry-trees of Aphis [= Myzus] cerasi, and of its enemies, the larva of Hippodamia convergens and Syrphus sp.; figure of the larva, pupa, and imago of the former; positions in which both species transform.
- 1252. [RILEY, C. V.] Grape-vine flea-beetle. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of C. H. Roberts; ravages of Haltica chalybea; means against this species and Selandria vitis [= Blennocampa pygmæa] on grapevines.
- 1253. [RILEY, C. V.] Blood-sucker and pear-slug. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of G. A. Watson; food-habits of *Pirates* [= *Melanolestes*] picipes and of Hemiptera in general; means against *Selandria* [= *Eriocampa*] cerasi.
- 1254. [RILEY, C. V.] The codling-moth. (Carpocapsa pomonella, Liunaeus.) < Amer. Ent. and Bot., October, 1870, v. 2, pp. 321-322. Reprint: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 101-104.
 - Treats of Carpocapsa pomonella; see No 1301 for synopsis of contents.
- 1255. [RILEY, C. V.] Insects injurious to the grape-vine. No. 12. <Amer. Ent. and Bot., October, 1870, v. 2, pp. 327-328, figs. 204-205. Reprint, with slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 79-81, figs. 35, 36; Cultivator and Country Gentleman, 1 June, 1871, v. 36, p. 343.
 - Treats of Graptodera [= Haltica] chalybea; see No. 1301 for synopsis of contents.
- 1256. [RILEY, C. V.] The fall army-worm. < Amer. Ent. and Bot., October, 1870, v. 2, pp. 328-329, figs. 206-207.
 - Seasons, food-plants, and ravages of Laphygma frugiperda; figures larvæ of Leucania unipuncta and Laphygma frugiperda; comparison between the latter and Heliothis armigera.

- 1257. [RILEY, C. V.] The rape butterfly; our new cabbage pest. <Amer. Ent. and Bot., October, 1870, v. 2, p. 338.

 Spread of and means against *Pieris rapa*.
- 1258. [RILEY, C. V.] Paris-green for the Curculio. <Amer. Ent. and Bot., October, 1870, v. 2, p. 338.

 Inefficacy of the use of Paris-green against Conotrachelus nenuphar.
- 1259. [RILEY, C. V.] Beetles working in wheat, oats, and rye; the grain Silvanus. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339, fig. 208.
 - Answer to inquiry of M. H. Boye; figure, description, and ravages of *Silvanus* surinamensis; means against the same and against *Calandra granaria* in grain; habitat and probable origin of the *Silvanus*.
- 1260. [RILEY, C. V.] Beetles in dried English currants. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of T. V. Munson; Silvanus surinamensis breeding abundantly in dried English currants.
- 1261. [RILEY, C. V.] The same in flouring mills. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.

 Answer to inquiry of S. Blanchard; abundance of Silvanus surinamensis in flouring-mills.
- 1262. [RILEY, C. V.] Carolina Sphinx. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.

 Answer to inquiry of W. R. Howard; the larva of Macrosila [= Protoparce] carolina feeds on tobacco.
- 1263. [RILEY, C. V.] Insects clustered on apple-trees. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.

 Answer to inquiry of R. L. Ham; habits and food of *Psocus venosus*.
- 1264. [RILEY, U. V.] Larvæ named. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of T. W. Gordon; stinging powers of the larvæ of *Empretia stimulea* and *Hyperchiria varia* [=io]; *Microgaster* sp. parasitic upon *Macrosila quinquemaculata* [= *Protoparce celeus*].
- 1265. [RILEY, C. V.] Mite-gall on sugar-maple. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of A. Furnas; description of gall of Acarus aceris-crumena n. sp. on leaves of sugar-maple; similar mite-galls occur on plum and cherry.
- 1266. [RILEY, C. V.] Cheese-fly and blow-fly. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of B.; skippers in cheese are larvæ of *Piophila casei*; those found in bacon of *Calliphora vomitoria*.
- 1267. [RILEY, C. V.] Fall army-worm. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiries of K. Kelsey and G. Pauls; seasons, ravages, and foodplants of *Laphygma frugiperda*; seasons and food-plants of *Leucania unipuneta*.

- 1268. [RILEY; C. V.] Larva of imperial moth; Thoas swallow-tail. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of G. M. Dodge; larva of Ceratocampa [= Eacles] imperialis feeds on maple.
- 1269. [RILEY, C. V.] Large Asilus fly. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of L. G. Saffer; food-habits of Promachus rertebratus, P. [= Erax] bastardii, and Asilus missouriensis [= Proctacauthus milbertii]; occurrence of undetermined galls under white-oak trees.
- 1270. [RILEY, C. V.] Mole cricket. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of V. K. Deyo; habits of Gryllotalpa borealis.
- 1271. [RILEY, C. V.] A rare capture in Illinois. <Amer. Eut. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of H. S. Bontell; geographical distribution of Callidryas philea; occurrence of Thysania zenobia in Iowa.
- 1272. [RILEY, C. V.] Hag-moth larva. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340, fig. 209.
 - Answer to inquiries of D. M. Hunter and G. Pauls; figure of larva and description of imago of Limacodes [= Phobetrum] pithecium; larva found on apple-tree; formation of its eoeoon.
- 1273. [RILEY, C. V.] Insects named. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of Mrs. E. U. B.; larva of *Papilio asterias* feeds on parsnip; that of *Alaria* [= *Rhodophora*] *florida* on evening primrose; habits of the image of the latter.
- 1274. [RILEY, C. V.] Some friends and foes. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of C. W. Spaulding; food-habits of Harpactor [=Milyas] cinctus, Mysia [=Anatis] 15-punctata, and Calosoma calidum; Prionus imbricornis bred from grape-vine roots.
- 1275. [RILEY, C. V.] The royal-horned caterpillar. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of J. T. Hodgen; larva of Ceratocampa [=Citheronia] regalis found on persimmon.
- 1276. [RILEY, C. V.] Caterpillars named. <Amer. Ent. and Bot., October, 1870, v. 2, p. 341, fig. 210.
 - Answer to inquiry of E. H. King; food-plants and descriptions of larvæ of Papilio asterias and of Aeronyeta oblinita; function of the osmaterium of the former; vernaeular name and figure of larva, eoeoon, and imago of the latter.
- 1277. [RILEY, C. V.] The Abbot Sphinx; parasites on its larva. <Amer. Ent. and Bot., October, 1870, v. 2, p. 341.
 - Answer to inquiry of T. W. Gordon; description of Thyreus abbotii; life-history of Microgaster sp. parasitic on larva of Sphingida.
- 1278. [RILEY, C. V.] Crane-flies; rose-bugs; ants. < Amer. Ent. and Bot., October, 1870, v. 2, p. 341.
 - Answer to inquiry of J. W. Potts; food-habits of Tipula sp. and of Macrodactylus subspinosus; structure and habits of the several sexes of Formicidæ.

- 1279. [RILEY, C. V.] Cabbage-worms. < Amer. Ent. and Bot., October, 1870, v. 2, p. 341.
 - Answer to inquiry of B. H. Foster; ravages of and means against Pieris rapæ.
- 1280. [RILEY, C. V.] The unicorn prominent. <Amer. Ent. and Bot., October, 1870, v. 2, p. 341.
 - Answer to inquiry of E. Payne; description of larva and image of *Notodonta* [= Cwlodasys] unicornis; food-plants and mimicry of the larva.
- 1281. [RILEY, C. V.] Insects injurious to the grape-vine. No. 13. <Amer. Ent. and Bot., December, 1870, v. 2, pp. 353-359, figs. 218-219. Reprint with changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 84-96, figs. 37-40. See: <Bull. de la Soc. Cent. d'Agric., 1870.
 - Treats of Phylloxera vastatrix; see No. 1301 for synopsis of contents.
- 1282. [RILEY, C. V.] The fall army-worm. < Amer. Ent. and Bot., December, 1870, v. 2, pp. 363–365, figs. 221–223.
 - Descriptions of egg, larva, pupa, and imago of *Prodenia autumnalis* n. sp. [= Laphygma frugiperda]; variation of the imagos; seasons, habits, and ravages of and means against the same; figures of larva and imago of *Prodenia commelinæ* and of imago of *Leucania unipuncta*.
- 1283. [RILEY, C. V.] The so-called web-worm of young trout. < Amer. Ent. and Bot., December, 1870, v. 2, pp. 366-367.
 - Editorial remarks appended to S. J. McBride's communication; description of Simulium piscicidium n. sp.
- 1284. [RILEY, C. V.] Hybrid between a grape-vine and a hickory. <Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Correction of error in mistaking a gall of Cecidomyia vitis-pomum for a hybrid fruit.
- 1285. [RILEY, C. V.] Death of noted entomologists. <Amer. Ent. and Bot., December, 1870, v. 2, p. 373.

 Notice of death of Julius Lederer and J. T. Lacordaire.
- 1286. [RILEY, C. V.] Osage orange for the mulberry silk-worm. <Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Explanation of diff rences in experience in feeding silk-worms on osage orange.
- 1287. [RILEY, C. V.] Insects named. < Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of Mrs. M. Chappellsmith; Astoma [= Trombidium] locustarum parasitic on grasshoppers; similar mites on other insects; food-habits of Nemobius vittatus and Orocharis saltator.
- 1288. [RILEY, C. V.] Locust-borer. < Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of W. R. Howard; seasons and method of oviposition of Arhopalus [= Cyllens] robiniæ in black-locust; description of eggs and imago of the same.
- 1289. [RILEY, C. V.] The northern lady-bird; its larvæ. < Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of C. E. Billings; food-habits and description of larva of Epilachna borealis.

- 1290. [RILEY, C. V.] Not eggs, but parasitic cocoons. <Amer. Eut. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of R. Couch; occurrence and true nature of cocoons of *Microgaster* sp. on larva of *Darapsa* [= Ampelophaga] myron.
- 1291. [RILEY, C. V.] The cabbage Plutella. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of C. E. Bessey; ravages of Plutella cruciferarum on cabbages.
- 1292. [RILEY, C. V.] Gigantic rhinoceros beetle. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374, fig. 224.
 - Answer to inquiry of "Subscriber;" figure of male and description of the male and female of *Dynastes tityus*; distribution; food-habits of larva.
- 1293. [RILEY, C. V.] Bee-bread devoured by worms. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of L. C. Francis; food-habits of Ephestia zew [=interpunctella] and Galleria cereana; ravages of the former in old beehives.
- 1294. [RILEY, C. V.] Rape butterfly. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.

 Answer to inquiry of J. E. Cowden; ravages of *Pieris rapæ* on cabbages.
- 1295. [RILEY, C. V.] Grape-leaf gall. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of H. C. Beardslee; occurrence of *Phylloxera vastatrix* at Painesville, Ohio.
- 1296. [RILEY, C. V.] Bean weevil. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of H. Kleinhaus; ravages and distribution of Bruchus obsoletus.
- 1297. [RILEY, C. V.] Cabbage insects. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of W. R. Howard; ravages, distribution, and synonyms of *Plutella cruciferarum*; first occurrence of *Strachia* [= *Murgantia*] *histrionica* in Missouri.
- 1298. [RILEY, C. V.] Two-striped walking stick. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of C. R. Edwards; occurrence of Spectrum bivittatum in Kentucky.
- 1299. [RILEY, C. V.] Ladder spider. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of L. G. Saffer; explanation of the vernacular name of Epeira [=Argiope] riparia.
- 1300. [RILEY, C. V.] Sugaring for moths; preserving larvæ. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of H. S. Boutwell; occurrence of *Papilio thoas* [=cresphontes] in Illinois; directions for the preparation of a fluid for preserving caterpillars.

PREFACE

1301. RILEY, C. V. Third annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <6th Ann. Rept. State Board of Agric. for 1870, April, 1871, pp. 176+7, 73 figs. Separate: Jefferson City, Mo., April, 1871, pp. 176+7, 73 figs.

CONTENTS.

3

29

NOXIOUS INSECTS.	
SNOUT-BEETLES, Rhynchophora	F
The whole vegetable kingdom and every part of each plant serve	
as food for insects, 5—Enumeration of insects affecting the dif-	
ferent parts of the apple-tree, 5—Other food-habits of insects, 7—	
Vast extent of the science of entomology, 8—Beauty and sim-	
plicity of classification in entomology, 8—Each family distin-	
guished by its general appearance, 8—Unity of habits in each	
family, 9—Distinguishing characters of snout-beetles, 9—Their	
larvæ, 10—They are among the most injurious beetles, 10—In-	
jurious snout-beetles in Europe, 11.	
<u> </u>	13
The common plum Curculio, Conotrachelus nenuphar	1.
It is single-brooded, 11—Experiments to prove this fact, 12—It hibernates as beetle, 13—Form of the egg, 13—Feeding habits of	
, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
the beetle, 13—Creaking noise produced by it, 14—Stridulation	
in other insects, 14—It is nocturnal rather than diurnal, 14—	
Habits of the beetle at night, 14—Remedies, 15—The Ransom	
chip-trap process, 15; explanation of the process, 15; it is not so	
successful as anticipated, 15; it is not a new discovery, 16; num-	
ber of Curculios caught by it, 16; its success dependent on the	
character of the soil, 17; directions for using the process, 17;	
more experiments needed, 17—Offering premiums for collecting	
specimens, 17—Absurdity of the application of Paris green for	
the Curculio, 18—Jarring by machinery, 18—The Hull Curculio-	
catcher defective in several respects, 18—The Ward Curculio-	
catcher, 20; how it could be improved, 20; rules for using the	
machine, 21—Curculio-catcher invented by Claxton & Stevens,	
22—Hooten's Curculio-catcher, 22; its advantages and mode of	
operation, 24—Two true parasites of the Plum Curculio, 24—The	
Sigalphus Curculio parasite, 24; Fitch's account of it, 25; Walsh's	
doubt about its being parasitic on the Curculio, 25; experiments	
and observations proving that it is a parasite of the Curculio,	
25; its development and frequent occurrence around St. Louis,	
25; it attacks also other soft-bodied larvæ, 26; points in its nat-	
ural history, 26; its position in the system, 27; description of	
the imago, larva, pupa, cocoon, and of the var. rufus, 27—The Pori-	
zon Curculio parasite, 28; how it differs from the foregoing, 28;	

The apple Curculio, Anthonomus quadrigibbus.

How it differs from the plum Curculio, 30—Its natural history, 30—
Food-plants, 30—The punctures it drills into the fruit, 31—The egg, 31—The larva and its habits, 31—The larva transforms within the fruit, 32—Amount of damage it does, 33—How its

tion of parasites, 29.

description of the imago, 28—Importance of the work of these parasites of the Curculio, 29—Artificial propagation and distribu-

BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	153
1301. RILEY, C. V.—Continued.	
work in the fruit can be distinguished from that of other insects, 33—It is very injurious in Southern Illinois and parts of Missouri, 33—It is less injurious to apples than the plum Cureulio, 33—Injury done by it to pears, 33—The rot in apples principally	
produced by it and by the plum Curculio, 34—Season of its appearance, 34—Remedies and preventive measures, 34—Descrip-	
tion of larva and pupa, 35.	35
The quince Cureulio, Conotrachelus cratægi	00
36—Dr. Trimble's account of the damage done by it, 36—How	
the beetle differs from the two preceding Curculios, 36—Its trans-	
formations and habits, 37—Its puneture on the fruit, 37—It hiber-	
nates in the larva state, 37—Its larva mistaken by Dr. Fitch for	
that of the plum Cureulio, 38—The imago state lasts only two	
months, 38—It does not attack the apple, 38—Remedies, 38—De-	
scription of the larva and pupa, 39. The plum-gouger, Coccotorus scutellaris	39
How it differs from the preceding species, 39—Injury caused by it,	00
39—Food-plants, 40—It is often mistaken for the plum Cureulio,	
40—Season of its appearance, 40—Holes bored by it in the fruit,	3
41—The larva lives within the kernel of the fruit, 41—Remcdies, 41	
The strawberry erown-borer, Tyloderma fragaria	42
Distribution of the insect, 42—Injury done by it, 42—Habits and characters of the larva, 43—Habits of the beetle, 43—Remedies,	
43—Parasite, 44—How the larva differs from that of the grape-	
vine Colaspis, 44—Description of the image and larva, 44.	
The pea-weevil, Bruchus pisi	44
Insect enemies of the garden-pea, 44—Characters of the Bruchida,	
45—Habits of other species of Bruchidee, 45—Frequent occurrence	
of its larva in green peas, 45—Charaeteristics of the beetle, 46— It is in all probability an indigenous North American insect, 46—	
The beetle does not sting the peas, 46—The eggs are fastened by	
the female beetle on the outside of the pod, 47—The larva and	
its habits, 47—Its transformations, 47—Remedies and prevent-	
ives, 48—Examination of peas intended for secd, 48—Concerted	
action necessary to exterminate the insect, 48—Mr. Saunders's ac-	1
eount of the occurrence of the pea-weevil in Canada, 49—Other preventive measures, 49—Birds destroying the insect, 50.	
The grain Bruchus, Bruchus granarius	50
Its introduction from Europe, 50—How it differs from the pea-	
weevil, 50—Curtis's account of its habits in Europe, 51.	
The American bean-weevil, Bruchus fabæ	52
Food-plants, 52—Its geographical distribution, 52—Accounts of	
damage done by it in New York and Penusylvania, 52—It has only lately become injurious, 53—Habits of the larva and bectle,	
54—The proper nomenclature of the species, 54—Description of	
the imago, 55—Its differences from allied species, 55—Note on	
descriptions based upon individual variations, 56.	
The New York weevil, Ithycerus novcboracensis	57
Damage caused by it to fruit trees, 57—Former descriptions of the insect, 57—Its breeding habits, 57—The pear blight not eaused	
by the beetle, 58.	
The imbrieated snont-beetle, Epicarus imbricatus	58

Injury done by it to vegetation, 58—Its natural history still un-

known, 58-General appearance of the beetle, 58.

1301. RILEY, C. V.—Continued.	
The corn Sphenophorus, Sphenophorus sculptilis	59
Damage caused by it to corn plants, 59—Characteristics of the	
beetle, 59—Its larval history still unknown, 59—Probable habits	
of the larva, 59—Walsh's description of the imago, 59.	
The cocklebur Sphenophorus, Rhodobanus 13-punctatus	60
It is not injurious, 60—Coloration of the beetle and its probable	
identity with S. 13-punctatus, 60—The larva, 60—Enumeration of	
other injurious snout-beetles, 60.	
INSECTS INJURIOUS TO THE GRAPE-VINE	61
The grape leaf-folder, Desmia maculalis	61
Its geographical distribution, 61—Generic characters, 61—Charac	
ters of the moth, 61—Sexual differences, 62—Habits of the larva,	
62—Remedy and prevention, 62—Natural enemies, 62—Descrip-	
tion of the larva, 62.	00
The grape-vine epimenis, Psychomorpha epimenis.	63
Its larva formerly mistaken for that of the pearl wood nymph, 63—	
Characters of the moth and sexual differences, 63—Habits of the	
larva when about to transform, 64—Its grape-vine feeding hab-	
its formerly unknown, 64—Description of the larva, 64; of the chrysalis, 65.	
The grape-vine plume, Oxyptilus periscelidaetylus	65
Work of its larva and of that of the grape-vine Epimenis, 65—Both	00
larvæ rather beneficial when not too numerous, 65—Characters	
of the larva, 66—Peculiar form of the pupa, 66—Protective mim-	
icry of the pupa, 67—Habits and appearance of the moth, 67—Is	
it single or double-brooded, 67.	
The common yellow bear, Spilosoma virginica	68
How the young larvæ differ from the mature larva of the grape-	
vine plume, 68—Food-plants, 68—Color variations in the larvæ,	
69—The chrysalis, 69—The imago, 69—Parasite, 69—Remedy, 69.	
The smeared dagger, Acronycta oblinita	70
The larva is polyphagous, 70—Characters of the larva, pupa, and	
imago, 70—Remedy, 70—Parasites, 71—Description of imago and	
larva, 71; of the pupa, 72.	
The pyramidal grape-vine worm, Pyrophila pyramidoides	72
Distinguishing characters of the worm, 72—Its food-plants, 72—	
Its transformations, 72—It is single or double-brooded according	
to latitude, 73—Its closely allied congener in Europe, 73—Rem-	
edies, 73—Description of the larva, 73; of the pupa and imago, 74—How it differs from Amphipyra pyramidea, 74—Description of	
the larva and image of the spattered copper under-wing, 75.	
The grape-root borer, Sciapteron polistiformis	75
Its distribution, 75—Distinguishing characters of the larva, 76—	10
Its transformation, 76—Characters of the moth, 76—Description	
of the image and its sexual differences, 76—Work of the larva on	
grape-roots, 77—Remedies, 77.	
The spotted Pelidnota, Pelidnota punctata	77
It is usually not injurious, 77—The larva and its habits, 78—The	
beetle, 78—Description of the larva, 78.	
The grape-vine flea-beetle, Haltica chalybea	79
It is well known to the grape-grower in Missouri, 79—Its distri-	
bution and food-plants, 80—Hibernation, 80—Damage done by	
the beetle in spring, 80—The eggs, 80—Damage caused by the	
larvæ, 80: their transformation, 80—Remedies, 80—Description	

of the larva and pupa, 81.

81

84

79

101

1301. RILEY, C. V.—Continued.

The grape-vine Colaspis, Colaspis flavida.

Fitch's account of the insect, 82—Varieties of the beetle, 82—The larva in all probability attacks sometimes that of the leaf-folder, 82—It lives on the roots of the strawberry, 82—Larva of the European Colaspis barbara, 82—Difference in habits of larvæ of allied species, 83—Singular characters of the larva of the grape-vine Colaspis, 83—Work of the larva on strawberry roots, 83—Remedy, 84—Description of the larva, 84.

The grape-leaf gall-louse, Phylloxera vastatrix..... Its life-history not yet fully studied, 84—Previous accounts of the insect by Fitch, Shimer, and Walsh, 85-The root disease in France, 85—The Phylloxera vastatrix recognized as the cause of this disease, 85-Identity of the gall-louse with the root-iuhabiting insect, 86—The American and Europeau insects are identical, 86—Remedies tried in France, 86—The disease directly eaused by the Phylloxera, 87—Injury done by the Phylloxera in Missouri, 87-Forming of the gall, 87-Propagation of the lice and multiplication of the galls, 88-The gall-liee descend in the latter part of the season to the roots, 88—Change of the insect after passing from the leaves to the root, 88-Questions still to be settled in the life-history of the Phylloxera, 88-Rare occurrence of the winged form, 89-The insect can be transported from one place to another on roots, 89—It hibernates on the roots, 89—Grape-vines that should be planted, 89-Number of indigenous species of the grape-vine, 90—Grape-vines which are most seriously infested with the grape-leaf louse, 90—Danger in planting the Clinton among other grapes, 91-Insects acquiring different food-habits as illustrated in the apple-magget and the pine-leaf seale, 92— The different forms of the grape-leaf louse, 93—Discussion on the proper place of the insect in the classification, 93—On Dr. Shimer's proposed new families Dactylosphæridæ and Lepidosaphidæ, 93—Objections to Fitch's specific name vitifolia, 95—Identity of the European with the American insect, 95-The apple-root louse is identical with the woolly Aphis, 95—The gall-inhabiting form of the Phylloxera identical with the root-inhabiting type, 96— Characters of the genus Phylloxera and its place in the system, 96.

The codling-moth again, Carpocapsa pomonella...

Hay-bands around the trunk of the tree more effectual than rags placed in the fork, 102—The codling-moth is single-broaded in the more northern countries, but double-broaded in the latitude of St. Lonis, 102—Sexual differences of the moth, 103—Sexual characters in the genera Argynnis and Grapta, 103—The codlingmoth also infests peaches, 103.

1301.	RILEY, C. V.—Continued.	
	The corn-worm alias boll-worm, Heliothis armigera. Its geographical range, 104—Injury done by it to corn, 104—It attacks tomatocs and other plants, 105—Food-plants of the stalk-borer, 105—The egg of the corn-worm, 105—Mr. Glover's account of the habits of the boll-worm, 106—The larva is very variable in color, 107—Its transformations, 107—Number of annual broods, 107—Amount of damage done by it, 107—Remedies, 108—Attracting the moth by sweets, 108—Heard's moth-trap, 109.	104
	Reports of its appearance in 1870, and how it was generally mistaken for the true army-worm, 109—It was also mistaken for the boll-worm, 111—Injury caused by it, 111—How it differs from the true army-worm, 112—It is a very variable species in the imago state, 113—The spiderwort owlet-moth, and how it differs from the fall army-worm moth, 113—Number of annual broods and time of appearance of the fall army-worm, 114—The eggs and how they are deposited, 114—Preventive measures, 115—It is never injurious during two consecutive years, 115—Parasitic checks, 116—Description of the imago, 116; of the varieties and the earlier states, 117.	109
	The APPLE-TREE TENT-CATERPILLAR, OR AMERICAN LACKEY-MOTH, Clisiocampa americana The web-nests of the caterpillar and importance of their destruc- tion, 118—The egg-mass, 118—The caterpillar and its habits, 119— Transformations of the insect, 119—The imago very variable in color, 119—Food-plants of the caterpillar, 120—Remedies, 120— Parasites and enemies, 120.	117
	The tent caterpillar of the forest, Clisiocampa disstria The egg-mass and how the eggs are deposited by the female moth, 121—Development of the larva, 122—Fitch's description of the full-grown larva, 123—Confusion arising from want of uniform rule in describing larvæ, 123—The cocoon, 124—The chrysalis and the moth, 124—The web spnn by the caterpillar, 124—Mr. Ferris's observations on differences in habits, appearance, and food-plants of the caterpillar, 125—Phytophagic varieties or species, 127—Food-plants of the caterpillar, 127—Its destructive powers, 127—Remedies, 128—Natural enemies and parasites, 128—Snmmary,	121
	The fall web-worm, Hyphantria cunea It is often mistaken for the tent-caterpillars, 130—It feeds upon almost every kind of tree and shrub, 130—The wcb spun by the worm, 130—General appearance of the worm, 130—The chrysalis and image, 131—Number of annual broods, 131—Plants it prefers, 131—How it differs from the tent-caterpillar, 132—Remedics, 132—Description of the larva, 132.	1 30
	THE BLUE-SPANGLED PEACH-WORM, Callimorpha lecontei Winter retreat of the larva, 132—General appearance of the larva, 133—Chrysalis and imago, 133—Callimorpha vestalis Packard synonymous with C. fulvicosta, 133—Food-plants, 134—Descrip-	132
	tion of the larva, 134. THE ASH-GRAY PINION, Lithophane cinerea	134

BENEFICIAL INSECTS.	
The Glassy-winged soldier-bug, Campyloneura vitripennis It preys upon different species of leaf-hoppers, 137—Its larva and pupa, 138—How it seizes its prey, 138—Coloration of the insect, 138—It was never observed before to attack the leaf-hoppers of the grape-vine, 139—Habits of the Phytocoride, 139.	137
INNOXIOUS INSECTS.	
The white-lined morning Sphinx, Deilephila lineata. Resemblance of the moth to a humming bird, 140Interest attached to the study of lepidopterous larvæ, 140Larval variations in the same species, 141-Food-plants of the larva, 141- The light and dark colored forms of the larva, 142Its geographical range, 142Parasite, 142.	140
Two of our common butterflies	142
The archippus butterfly, Danais archippus Synonymy of its specific name, 143—Its geographical distribution, 143—Characters of the Danaidæ, 143—Sexual differences, 143— Food-plants, 144—Hibernation, 144—Two annual broods, 144— Description of the egg, 144—The larva and its larval changes, 145—How the horns of the larva become longer at each molt, 145—The full-grown larva, 146—Interest attached to the metamorphoses of insects, 146—How the larva becomes a chrysalis, 147—The hardened chrysalis, 147—Duration of the chrysalis state, 147—The issuing of the butterfly, 148—Protective colors of insects, 148—Nauscous odor of the archippus butterfly in all its stages, 149—The Tachina-parasite of the archippus larva, 149— Action of the parasitized larva, 149—How the Tachina larva and other insects prepare the lid of their puparia, 149—Characters of this Tachina-fly, 150—Difficulties of the study of the Tachinariæ, 150—Two forms occurring in Tachina archippivora, 150— The butterfly often congregating in immense swarms, 151—Probable reasons of this assembling in swarms, 152.	143
The disippus butterfly, Limenitis disippus	153
Distinguishing characters of the Nymphalidæ, 153—Food-plants and geographical range, 153—The egg, 153—Description of the egg, 154—Development of the larva, 154—Description of the mature and young larva, 154—The insect hibernates as young larva, 155—Case prepared by the larva for its winter quarters, 155—Modifications of the case, 156—Peculiar habit of the autumnal larvæ, 156—Parasites, 157—Tachina-fly, 157—The Disippus eggparasite, 157—The disippus Microgaster, 158.	100
Mimicry as illustrated by these two butterflies, with some remarks on the theory of natural selection Conformity of color between animals and their surroundings, 159— Definition of the term "mimicry," 160—Pungent odor possessed by the Danaidæ, 160—Their mode of flight, 160—Protection they derive from their peculiar odor, 161—Pieridæ and Danaidæ in the valley of the Amazon, 161—Mimetic forms of Pieridæ, 161—Explanation of the origin of mimetic forms, 162—Mimicry between the archippus and disippus butterflies, 163—The ursula butterfly, 163—Its caterpillar and chrysalis undistinguishable from those of the Disippus, 163—Its imago has no protective color,	159

producing mimetic forms, 165—Mr. Scudder's objections, 166—Discussion of Mr. Scudder's arguments, 167—Mr. Murray's ob-

- 1301. RILEY, C. V.—Continued.
 - jections to the connection between mimetic resemblances and natural selection, 170—Natural selection not the only power producing mimicry, 171—Reasons for discussing in this report the theory of natural selection, 172—Natural selection involves belief in the doctrine of evolution, 173—Darwinism is neither irreligious nor atheistic, 174.
- 1302. RILEY, C. V. Snout-beetles injurious to fruits. <Trans. Ill. Hortic. Soc. for 1870, 1871, n. s., v. 4, pp. 89–124, figs. 1–11. Reprint, with additions and omissions: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 5–44, figs. 1, 3, 4, 7–14. See No. 1301 for synopsis of contents.
- 1303. RILEY, C. V. Bark-lice on rose bushes. <Moore's Rural New Yorker, 24 June, 1871, v. 23, p. 393. S.-b. No. 4, p. 3.

 Answer to inquiry of W. A. French; occurrence of and means against Lecanium rosæ [= oleæ] and Diaspis rosæ on rose bushes.
- 1304. RILEY, C. V. Canker-worms; not army-worms. < Moore's Rural New Yorker, 24 June, 1871, v. 23, p. 393.
- 1305. RILEY, C. V. How to distinguish between *Limenitis disippus* Godt., and *L. ursula* Fabr., in their preparatory states. <Ca. Ent., July, 1871, v. 3, pp. 52–53, fig. 24.
 - Description and figures of structural characters distinguishing the larva and pupa of *Limenitis disippus* from those of *L. ursula*; see No. 1306.
- 1306. RILEY, C. V. Friendly notes. <Ca. Ent., September, 1871, v. 3, pp. 117-119.
 - Distinctive structural characters of pupe of Limenitis disippus and L. ursula; criticism of W. Couper's articles; description, food, and habits of Gelechia [=Holcocera] glandulella n. sp.; review of statements concerning the dimorphism of Grapta interrogationis.
- 1307. RILEY, C. V. Friendly criticism. < Gardener's Mo. and Hortic., November, 1871, v. 13, p. 341.
 - Inefficacy of the roller remedy for potato-beetles and grasshoppers; identifies "Melolontha philophaga" as Lachnosterna quercina [=fusca].
- 1308, RILEY, C. V. Parasites on fowls. The chicken mite. < Moore's Rural New Yorker, 16 December, 1871. S.-b. No. 5, p. 31.
 - Answer to inquiry of H. Hales; injuries to fowls caused by *Dermanyssus galline?* [=avium].
- 1309. RILEY, C. V. The American Entomologist. <Cultivator and Country Gentl., 21 December, 1871, v. 36, p. 809. Reprint: <Gardener's Mo. and Hortic., January, 1872, v. 14, p. 23. <Ca. Ent., January, 1872, v. 4, p. 19.
 - Announcement of the continued suspension of publication of the American Entomologist.
- 1309a. [RILEY, C. V.] (Lxodes bovis.) < Rept. Comm. of Agric. on the diseases of cattle. Washington, 1871, p. 118, fig. Description and figure of Lxodes bovis n. sp.
- 1310. RILEY, C. V. The acorn moth. *Holcocera glandulella*, n. sp. Ca. Ent., January, 1872, v. 4, pp. 18, 19.
 - Description of larva and imago of Holeocera glandulella n. sp.; habits of larva; food-plant and description of larva of Balaninus rectus.

1311. RILEY, C. V. Fourth annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <7th Ann. Rept. State Board of Agric. for 1871, April, 1872, pp. 146+6, 66 figs. Separate: <Jefferson City, Mo., April, 1872, pp. 146+6, 66 figs. Notice: <Horticulturist, August, 1872, v. 27, p. 251. <Ent. Mo. Mag., July, 1872, v. 9, p. 47.

CONTENTS.

Preface	3
NOXIOUS INSECTS.	
NOTES OF THE YEAR	5
	5
Its injuries in 1871, 5—Its appearance in great numbers in early spring, 5—Exorbitant price of Paris green, 6—Natural enemies of the beetle very abundant, 6—Diminution in numbers of the beetle later in the season, 7—Causes of such diminution, 7—Damage caused by the potato-beetle in Missouri, 7—It invaded Canada in 1870, 8—The three-lined potato-beetle mistaken for it in New York and Massachusetts, 8—Its further spread eastward irresistible, 8—Slow spread of the insect in the South, 9—Its present extent northward, 9—It spreads but does not leave the districts already invaded, 9—It is not injurions to potatoes in Colorado at a certain altitude, 10—New food-plants, 10—It feeds upon cabbage, 10—Its hibernation, 11—Objections raised against the use of Paris green, 11—Paris green is an efficient remedy and now in general use, 12—Box for dusting Paris green, 12—Mixing the poison with diluents, 12—No serious cases of poisoning have come to knowledge, 13—Antidote for Paris green, 13—Other applications, 13—Messrs. Saunder's and Reed's experiments with various substances, 14—Experiments with decoctions of various	
plants, 15—Air-slacked lime as a remedy, 15—Mechanical means, 15—Squire's brushing machiue, 15—Creighton's improved patent insect destroyer, 15—Disadvantage of all mechanical means, 16—A simple and effective way of brushing off and killing the bugs, 16—Natural enemies increasing, 16—Chickens acquiring a taste for eggs and larve of the beetle, 16—Spiders are among its enemies, 17—The 15-spotted lady-bird and its larva, 18—The icy lady-bird, 18—The ring-banded soldier-bug, 19—The dotted-legged plant-bug, 19—The spined soldier-bug in its earlier states, 20—The Nebraska bee-killer, 21—The Kansas bombardier-beetle, 21—Rove-beetles of the genera Philonthus and Quedius, 21. The codling-moth again, Carpocapsa pomonella. Time of year that the first moths appear, 22—Time required for development, 22—Proper time to apply the bandages around the tree, 22—It attacks peaches, 22—Best kind of bandages, 23—Wier's apple-worm trap, 23—Advantages and disadvantages of the trap, 24—Overestimating the value of Wier's trap, 25—Jarring, 25—Mr. Chapin's method of knocking down the wormy apples, 26—When this operation should begin, 26—Fires, lights, and bottles of liquid not to be recommended as a remedy, 27—Worth-	22

lessness of Todd's book, "The apple culturist," 28-Natural ene-

^{*} Extract in < Sei. Amer., 25 May, 1872, v. 25, p. 351.

1311.	RILEY, C. V.—Continued.	
	mies, 28—The Pennsylvania soldier-beetle and its larva, 28—The	
	two-lined soldier-beetle and its larva, 29—Description of the	
	larva of Telephorus bilineatus, 30—Summary, 30.	
	The periodical Cicada, Tibicen septendecim.	30
	Reproduction, with corrections and additions, of the chronolog-	00
	ical table of Broods I to VI from Report I, 31.	
	The grape-vine Colaspis again, Colaspis flavida	34
	Redescription of the larva from well-preserved specimens, 34.	•
	THE HARLEQUIN CABBAGE-BUG, Murgantia histrionica	35
	It was not known in Missouri prior to 1870, 35—Its geographical	00
	range and color variations, 35—Insect enemies of the cabbage-	
	plant, 35-Dr. Lyncecum's account of its habits and injury	
	caused by it in Texas, 36—Its appearance in Missouri in 1870,	
	36—The egg, 37—The larva and pupa, 37—Several annual broods,	
	37—The mature bug, 37—Injury caused by it, 37—Its congener	
	in Europe, 38.	
	THE RASCAL LEAF-CRUMPLER, Acrobasis indignella	38
	Its proper scientific name, 38—It is hardly noticed in summer time,	
	38—Injury caused by it, 39—It hibernates as larva, 39—Habits of	
	the larva, 39—The larval case, 39—Characteristics of the moth,	
	39—Food-plants, 39—Remedies, 40—Natural enemies, 40—Le	
	Baron's description of Tachina phycitæ, 40—Limneria fugitiva, 41—	
	Description of imago, larva, and chrysalis of the rascal leaf-	
	crumpler, 41—Description of the variety nebulella, 42.	
	THE WALNUT CASE-BEARER, Acrobasis juglandis	42
	Other case-bearers enumerated, 42—The case of the walnut case-	
	bearer, 42—Differences between the moth and that of the rascal	
	leaf-crumpler, 43—Natural enemies, 43—Description of Perilitus	
	indigator, 43.	
	THE APPLE-LEAF SKELETONIZER, Pempelia hammondi	44
	Its work on the leaves of apple-trees, 44—The worm and the chrys-	
	alis, 44—Appearance of the moth, 45—Mr. Hammoud's account	
	of the injury caused by the worm, 45—The European Acrobasis	
	consociella, 45—Remedies and parasites, 45—Description of the	
	imago, larva, and pupa, 46.	
	THE GREEN APPLE LEAF-TYER, Teras cinderella	46
	It occurs almost always in company with the foregoing, 46—	
	Characters and habits of the worm, 46—The chrysalis, 47—De-	
	scription of imago, larva, and chrysalis, 47.	
	THE LESSER APPLE LEAF-FOLDER, Teras malivorana	47
	Its larva and pupa closely resemble those of the foregoing species,	
	47—Mr. Wier's account of its habits, 48—Remedy, 49.	40
	THE APPLE-LEAF BUCCULATRIX, Bucculatrix pomifoliella	49
	It is not very injurious in Missouri, 49—Account of damage caused	
	by it in New York, 50—The worm and its habits, 50—Its transformations, 50. Season of the appearance of the moth, 50. Rem	
	formations, 50—Season of the appearance of the moth, 50—Remedies, 50—Habits of Bucculatrix thuiella, 51—Description of larva	
	and pupa of the apple-leaf Bucculatrix, 51.	
		51
	THE APPLE-TWIG BORER, Amphicerus bicaudatus	91
	52—The holes made by it in the twigs, 52—The holes are made	
	only for food and protection, 52—The insect breeds probably in	
	the sap-wood of forest trees, 52—The larva of Sinoxylon basilare	
	mistaken for that of the apple-twig borer, 52—Remedy, 53.	
	missaken for the apple-twig botter, oz-nemeny, so.	

72 72

	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	101
1311.	RILEY, C. V.—Continued.	
	INSECTS INJURIOUS TO THE GRAPE-VINE	53
	The red-shouldered Sinoxylon, Sinoxylon basilare	53
	Characteristics of the insect as imago, larva, and pupa, 54—Dam-	
	age done by it to grape-vines, fruit trees, and hickory, 54—De-	
	scription of the larva and pupa, 54.	
	GRAPE DISEASE	55
	The grape-leaf gall-louse, Phylloxera vastatrix	55 55
	Its scientific name, 55—The law of priority, 55—European grape-	JJ
	vines valueless in the eastern United States, 55—Deterioration	
	of some of our native vines, 56—Climatic reasons for the failure	
	of European vines, 56—The principal cause of this failure is the	
	Phylloxera, 56—Further proof of the identity of the American	,
	with the European insect, 57—Reasons for the identity of the	Ŷ
	gall-louse with the root-louse, 57—Further facts respecting	
	the habits of the root-louse, 58—Underground forms, 58—The	
	young lice and their habits, 59—Hibernation, 59—The pupa, 59—	
	The winged female, 59—Susceptibility of different vines to the	
	attacks of the louse, 60—Classification of the North American	
	grape-vines according to their practical importance, 60—Synopsis	
	of the true grape-vines of the United States, by Dr. George En-	
	gelmann, 60—Difficulty of separating the cultivated varieties of	
	grape-vine, 61—Importance of a proper classification of culti-	
	vated grape-vines, 62—Importation into Europe of resisting	
	American vines, 62—Enumeration of the cultivated varieties and	
	their susceptibility to the disease, 63—No variety is entirely ex-	
	empt from the attacks of the root-louse, 64—American vines	
	which resisted the root-louse in Europe, 64—Means of contagion	
	from one vine to another, 64—The spread of the root-lice from	
	one vine to another, 64—Transportation of the louse upon the	
	roots of seedlings and cuttings, 64—Spread of the disease in	
	France, 64—The winged female of the root-lonse and her func-	
	tion, 65—The vernal leaf-gall, 65—Preference of the winged	
	Phylloxera for the Clinton grape and its allies, 65—Power of	
	flight of <i>Phylloxera carywfoliw</i> , 66—Deposition of the eggs upon	
	the leaf, 66—Probable reasons why the injuries of the Phylloxera	
	are greater in Europe than in America, 66—Outward and more	
	visible effects of the root-disease, 67—Practical suggestions, 67—	
	No need to destroy the Clinton vines, 67—Influence of soil on	
	the intensity of the disease, 67—Remedies, 68—Destruction of	
	the gall-lice, 68—Destruction of the root-lice, 68—Results of ex-	
	periments in France with various substances, 68—Irrigation and	
	submersion, 69—Résumé of the insect's history, 69—No need of	
	unnecessary alarm, 70.	

BENEFICIAL INSECTS.

	•
SILKW	ORMS
Intr	oductory
Tł	ne Morus multicaulis fever and its reaction, 72—Increasing atten-
	tion lately given to silk-culture in America, 72-North America
	well adapted to the raising of silk, 73—General outlines of the
	natural history of the eight species of silkworms treated of in
	this report, 74.

The mulberry silkworm, Sericaria mori.....

75

Its past history, 75—Earliest silk-culture in China and India. 75— Its introduction into Europe, 76-Value of silk produced in France, 76-The "Silk Supply Association" in England, and its objects, 76-Countries exporting raw silk, 77-The name given to silk by different nations, 77-History of the mulberry silkworm in America, 77-Its introduction and failure of earlier efforts, 77—Renewal of silk-culture within the past decade, 77— Silk manufacture in the United States, 78—Favorable prospects for raising silk in this country, 78-Silk-growing in California, 79—Mr. Prevost's "California Silk-grower's Manual," 79—False statements and exaggerations in Prevost's book, 79—Sale of Californian silkworm eggs, 80-Disastrous effect of the Franco-Prussian war on the egg trade, 81—Success of silk-culture in California dependent on the ability to reel the silk, 81-Silk-enltnre in Kansas, 82-Mons. Boissière's silk establishment and its ehances of success, 82—Silk-growing in Missouri, 83—The fall season in Missonri eminently propitious for rearing silkworms, 84—Natural history of the silkworm, 84—Races of the silkworm produced by domestication, 84—Effects produced on the insect by domestication, 85—The egg, 86—Larva and larval changes, 86-Cocoon, chrysalis, and moth, 87-No insect parasite of the silkworm in Europe, 87-The "Uji" disease in China and Japan, 87—Diseases of the silkworm, 88; the Muscardine, its effect and cause, 88; the Pébrine disease, its symptoms and cause, 89; nature and origin of these diseases, 90—Other diseases of the silkworm, 91—Best varieties or races, 90—Different forms of eoeoons produced by different races, 92-How best to rear silkworms, 92—Rearing a very simple process, 92—Character of elimate of the Japanese silk districts, 93-Keeping the eggs dnring the winter, 93—Hatching of the eggs, 93—Room and building for the rearing of the worms, 93—The feeding net or fillet, 94— Importance of carrying all the worms simultaneously through their molts, 94—Regularity of feeding, 95—Regulating the temperature, 95—Cocoonery, 96—Choking the ehrysalis, 96—Egglaying, 97—Selecting and fastening the cocoons for breeding purposes, 97—Treatment of the female moths after copulation, 97— Preservation of the eggs, 97—Reeling, 98—Great skill required to accomplish the work properly, 98-Classification of raw silk, 98—Preparing the cocoons for reeling, 98—Objects of and manipulations in recling, 99—Best food for the worms, 100—Varieties of the mulberry, 100-Cultivation of the mulberry, 100-Osage orange as silkworm food, 100-Introduction of the osage orange into France, 100-Experiments in America with feeding silkworms on osage orange, 101-Advantages and disadvantages of the osage orange, 102.

The cecropia silkworm, Attacus cecropia.....

Changes made in its scientific generic name, 103—General appearance of the moth, 103—Fitch's explanation of the specific name, 104—Food-plants, 104—The econon, 104—Value of the cocoon as compared with that of the polyphemus moth, 105—How the moth issues from the cocoon, 105—The moth immediately after hatching, 106—The egg, 106—Description of the larval changes,

103

112

121

123

125

130

137

106—Strange habit of birds of using the empty cocoon as a store-honse, 107—The eecropia worm can not be classed as an injurious insect, 107—Samia columbia ought to be considered a variety of cecropia, 107—Parasites, 107—The long-tailed Ophion, 107; its mode of oviposition, 108; habits of its larva, 108—The eccropia Tachina-fly, 108; how it affects its victim, 108; its larva and imago, 109—The Mary Chalcis-fly, 109; how it escapes from the cocoon of the moth, 109; description of the imago, 110—The cecropia Cryptus, 110; its habits, 110; description of the two sexes, 111; how it differs from the allied species, 111.

The Ailanthus silkworm, Attacus cynthia.

Difference between the silkworms of the castor-bean and Ailanthus, 112—Introduction of the Ailanthus silkworm into Europe, 113—Ailanthiculture in Europe, 113—Its introduction into America, 114—Disadvantages of the Ailanthus silkworm, 115—Its acclimatization in America and Australia, 115—Value of the cocoon, 115—Mr. Forgemol's device for unwinding the cocoon, 116—Natural history of the Ailanthus silkworm, 117—The egg and mode of egg-laying, 117—Larval changes, 117—Pupation, 118—The image and its variations, 118—Food-plants, 118—Directions for raising the worms, 119—A good word for the Ailanthus tree, 120.

The luna silkworm, Actias luna.

The beautiful appearance of the moth, 123—The cocoon, 124—Foodplants, 124—Larval changes, 124—Season when the imago issues, 125—No parasites of the worm known, 125—Actias selene probably identical with luna, 125.

The yama-maï silkworm, Antheraa yama-mai...

Its native home and food-plant, 130—Its introduction into Europe and Australia, 130—Attempts at raising it in America, 130—The egg and how it should be kept over winter, 131—Climate most favorable for the worm, 131—The worm and its resemblance to the leaf, 132—Food-plants, 132—Larval changes, 132—The cocoon and the value of its silk, 133—The moth and its habits, 133—Difficulties of rearing the worm, 133—Mr. Adams's account of rearing the yama-maï worm in Japan, 134—The Uji parasite, 136.

The perny silkworm, Antherwa pernyi

Its native home, 137—How it differs from the preceding, 137—Larval changes, 137—The cocoon and its silk, 137—The moth, 137—Its culture in China, 138—The tusseh silkworm, 138.

1311.	RILEY, C. V.—Continued.	
	Summary	138
	Successful silk culture possible in this country, 138—Comparative value of the different species of silkworms, 138.	
	INNOXIOUS INSECTS.	
	The horned Passalus, Passalus cornutus. Its frequent occurrence in old logs, 139—The noise produced by the beetle, 139—It occurs only in decaying wood, 140—The larva and its exceptional character, 140—Description of the larva, 140—Previous description of Passalus larvæ, 141—Description of	139
	the pupa, 141. THE GREAT LEOPARD MOTH, Ecpantheria scribonia Characters of the larva, 141—Food-plants, 142—Hibernation and transformations, 142—The imago, 142—The larva able to resist extreme cold, 143—Description of the larva, 143.	141
	The Isabella tiger moth, Pyrrharctia isabella. Characters of the larva, 143—Food-plants, 143—Cocoon and chrysalis, 143—The moth, 144—The popular name "fever-worm" in the South, 144—No parasite known of this and the preceding species, 144.	143
	The acorn moth, Holcocera glandulella The egg laid in acorns that have been ruined by Balaninus rectus, 144—Habits of the larva, 144—Characters of larva and moth, 145—Description of imago, larva, and pupa, 145.	144
1312.	RILEY, C. V. Thomas Wier's apple-worm trap. <amer. 1="" 142–143,="" 1872,="" 31,="" ag="" and="" apple-worm="" april,="" description="" fig.="" figure="" of="" pp.="" td="" trap.<="" v.="" wier's=""><td>ric.,</td></amer.>	ric.,
1313.	[RILEY, C. V.] Remarkable parasitic fungus. <sci. 1872="" 26,="" 347.="" 40],="" [v.="" amer="" and="" description="" figure="" fungus="" fusca.<="" grub,="" infesting="" lachnosterna="" larva="" may,="" n.="" of="" p.="" s.,="" td="" the="" unidentified="" v="" v.=""><td></td></sci.>	
1314.	RILEY, C. V. Cut-worm lion. < Colman's Rural World, 15 July 1872, fig. Sb. No. 5, p. 47. History of the larva and image of Calesoma calidum, an enemy to cut-week.	
1315.	RILEY, C. V. Cut-worms. < Cultivator and Country Gentler 20 June, 1872, v. 37, p. 392. Dandelions and other weeds furnish food for young ent-worms which he in the fall; land should be kept clear of weeds at that season.	·
1316.	RILEY, C. V. Flat-headed apple tree-borer in horse chest Colman's Rural World, 22 June, 1872, fig. Sb. No. 5, p Auswer to inquiry of S. S. R.; natural history of larva and image of Chebothris femorata.	. 47.
1317.	RILEY, C. V. A new insect. < Western Planter, 29 June, 1 Sb. No. 5, p. 49. Answer to inquiry of F. Halsinger; Nysius n. sp., very destructive to tatoes.	
1318.	RILEY, C. V. Codling-moth; jarring down infested for Cultivator and Country Gentleman, 4 July, 1872, v. 37, p.	

Excrement outside of the fruit indicates the exit of the worm, with some varieties; with others the worm may be often found after the excrements

are visible.

- 1319. RILEY, C.-V. Food for silkworms. < Colman's Rural World, 13
 July, 1872. S.-b. No. 5, p. 40.
 Osage orange superior to red-mulberry.
- 1320. RILEY, C. V. The apple maggot-fly, Trypeta pomonella Walsh. <Amer. Agric., July, 1872, v. 31, pp. 263–264, 2 figs.
 - Answer to letter of J. H. Spatter; comparative descriptions and figures of larvæ, pupæ, and imagos of Trypeta pomonella and Carpocapsa pomonella; figures showing injuries of both species; distribution and means against Trypeta pomonella.
- 1321. RILEY, C. V. Worms on Dutchman's pipe. <Colman's Rural World, 3 August, 1872, fig: S.-b. No. 5, p. 136.

 Answer to inquiry of J. T. C.; life-history of Papilio philenor.
- 1322. RILEY, C. V. Apple-leaf worm. The apple-leaf skeletonizer. <Colman's Rural World, 10 August, 1872, fig. S.-b. No. 5, p. 138.
 - Answer to inquiry of L. R. Bryant; habits and times of appearance of Pempelia hammondi.
- 1323. RILEY, C. V. Eggs in grape-canes and apple-twigs. <Amer. Agric., August, 1872, v. 31, p. 302, figs. 1-7.
 - Figures of eggs and of twigs punctured by Orocharis saltator, Œeanthus niveus, and Ceresa bubalus; figures images of the same; habits, descriptions of eggs, and images of the three species.
- 1324. RILEY, C. V. Borers in evergreens. < Gardener's Mo. and Hortic., December, 1872, v. 14, p. 373.
 - Larvæ of Monohammus titillator the worst enemy of the white-pine; another species of Cerambycidæ injurious to red-cedars; a host of bark-borers affect all kinds of evergreens.
- 1325. RILEY, C. V. Einige unserer schädlicherer Insekten. <St. Louis, Mo., 1872, p. 35, il. S.-b. No. 19, pp. 174–190.
 - Causes of the decadence of viticulture in the United States; habits, ravages, specific identity, and figures of several stages of the leaf-inhabiting and root-inhabiting forms of Phylloxera vastatrix; identity of the European and American grape Phylloxera; susceptibility of the different species of grape to the attacks of the same; principal manifestations of attack; preventive and remedial measures; description and figures of Conotrachelus nenuphar; habits, ravages, parasites, enemies, and means against it; description of Ransom's trap, Hull's, Ward's, and Hooten's machines for capturing the imagos; figures of the machines. Natural history, food-plants, distribution, enemies, parasites, and means against Doryphora decemlineata; figures of all stages of the same; use of Paris green and other powders; mechanical devices for collecting the insects.
- 1326. RILEY, C. V. Harvest mites. '<Amer. Nat., January, 1873, v. 7, pp. 16–19, fig. 5. Extract: <Colman's Rural World, 21 June, 1876, 2 figs. S.-b. No. 10, pp. 170, 171.
 - List of insect parasites on man; description, figures, habits of and means against Leptus [= Tetranychus] americanus n. sp. and L. [= T.] irritans n. sp.
- 1327. RILEY, C. V. Vanessa antiopa. <Ent. Mo. Mag., January, 1873, v. 9, p. 195.

Habits and food-plants of Vanessa antiopa.

1328. RILEY, C. V. Entomological correction. < Cultivator and Country Gentleman, 6 March, 1873, v. 38, p. 149.

Lema trilineata does not occur in sonthwestern Missouri, as stated by W. R. Howard.

1329. RILEY, C. V. Fifth annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <8th Ann. Rept. State Board of Agric. for 1872, 18 April, 1873, pp. 160+8, 75 figs. Separate: <Jefferson City, Mo., 18 April, 1873, pp. 160+8, 75 figs. Review by A. S. Packard, jr.: <Amer. Nat., August, 1873, v. 7, pp. 471-477, figs. 115-130. Reply to Packard and rejoinder by Packard: <Amer. Nat., March, 1874, v. 8, pp. 181-188.

CONTENTS.

3 5

AEFACE
Entomology, its relations to agriculture and its advancement*
Definition of entomology, 5—The term "insect," 5—The four sub-
kingdoms of the animal kingdom, 6—Characteristics of insects,
7—Classification of insects, 8—Hymenoptera, 9—Coleoptera,
10-Lepidoptera, 11-Hemiptera, 12-Diptera, 13-Orthoptera,
14-Neuroptera, 14-Osculant and aberrant groups, 15-Import-
ance of entomology as a study, 17—Economic importance of en-
tomology, 18—Relation of insects to agriculture, 18—Damage
inflicted by insects to agriculture and horticulture, 19—Progress
of economic entomology, 19—Economic entomology in Europe,
19—Insects in America much more injurious than in Europe, 21—
State entomologists and entomologists in public positions in the
United States, 21—How to counterwork noxious insects, 23—
Prevention, 24—Checking the spread of injurious insects, 24—
Cure, 25—Modification of Dr. Hull's Curculio-catcher, 25—En-
couragement of the natural enemics of injurious insects, 26—
Duties of a State entomologist, 27—Annual report, 27—Corre-
spondence and other duties, 28—How to collect, preserve, and
study insects, 29-Appliances for collecting, 29-Use of the um-
brella, 29—The knapsack, 30—The hand-net, 30—The frame of
the net, 30—The bag of the net, 31—Use of the sieve, 31—Sugar-
ing, 32—Attracting moths by light, 32—How to kill insects, 32—
The cyanide bottle, 32-Use of chloroform, 33-Entomotaxy,
34-Insect-pins, 34-How to pin insects, 34-How to mount small
insects, 34—Spreading-board, 35—Drying-box, 36—Insect boxes
and cabinet, 37—Mr. Lintner's boxes for Lepidoptera, 38—Sub-
stances for lining insect-boxes, 40—Preserving insects in the
collection, 41—Relaxing specimens, 41—Breeding insects, 41—
Breeding-cage, 42-Note-book of the collector, 43-How to trans-
mit insects, 44—Text-books, 44.

NOVIOUS INSECTS

NOXIOUS INSECTS.	
Notes of the Year	46
The codling-moth, Carpocapsa pomonella	46
Experiments with Wier's apple-worm trap, 46-Value of different	
materials for bandages, 47—Jarring, 48—Occurrence of the apple-	

^{*} Pages 17-27 were reprinted under title Entomology in < Rept. State Board Agric. Kansas for 1873, 1874, pp. 266-274.

4 000		
1329	D. RILEY, C. V.—Continued.	
	worm in California, 49—Enemies and parasites, 49—The ring-	
	legged Pimpla, 49—The delicate long-sting, 50—Other enemies,	
	51—Efficiency of the spined soldier-bug, 51—False doctrines for	
	exterminating the codling-moth, 51.	
	The Colorado potato-beetle, Doryphora 10-lineata	52
	Its comparative harmlessness in 1872, 52—New food-plants, 52—	
	Its progress eastward up to 1872, 52—Experiments with Paris	
	green, 53—New enemies, 53—The rust-red social wasp, 54—The	
	rose-breasted grosbeck, 54.	
	The apple-twig borer, Amphicerus bicaudatus	54
	It attacks also pear twigs, 54—It has been bred from grape canes	
	by Dr. Shimer, 54.	
	Egg of the horned Passalus, Passalus cornutus	55
	The egg, 55—The newly hatched larva, 55—Rapid development of	
	the insect, 55.	
	Egg of the common May beetle, Lachnosterna fusca	55
	Description of the eggs and how they are deposited, 55.	
	Egg of the broad-necked Prionus, Prionus laticollis	56
	Characteristics of the eggs and where they are deposited, 56.	
	Eggs of American tent-caterpillar, Clisiocampa americana	56
	Correction of the figure given in Report III of the egg-belt, 56.	
	Counterworking the tobacco-worm, Protoparce celeus	56
	Mr. White's method of planting the Jamestown weed among the	
	potatoes and poisoning the blossoms thereof, 56.	
	THE GRAPE PHYLLOXERA, Phylloxera vastatrix:	57
	Its popular name, 57—Accounts of the unusual mortality among	
	grape-vines in the spring of 1872, 57—Causes of this mortality	
	given in the accounts, 59—Excessive drought and overbearing,	
	60—The Phylloxera is the true cause, 60—Actual proof of the	
	Phylloxera having caused the mortality among grape-vines, 61—	
	Influences that favored the increase of the lice in 1872, 61—Im-	
	portance of a full understanding and management of the Phyl-	
	loxera, 62—Its range in North America, 62—Its spread in Europe,	
	63—Inconstancy in the habits of the gall-lice, 63—The leaves of	
	the Clinton vine no longer affected by the gall-lice since 1871,	
	63—Method of formation of the leaf-gall, 64—Relative immunity	
	of American vines in Europe, 64—Propagating American varie-	
	ties from cuttings, 65Appreciation of American vines in Eu-	
	rope, 66—Careless statement published as to the immunity of	
	Labrusca vine, 66—Grafting the grape-vine, 67—New theories,	
	67—The Phylloxera is the true cause and not the effect of the	
	disease, 67—Mr. Laliman's theory that the insect has always ex-	
	isted in Europe, 68—Oidium tuckeri of Europe and America iden-	
	tical, 69—Mr. Saunders's account of the presence of Oidium tuckeri	
	in America, 70—Means of contagion of the disease from one vine to	
	another, 69—Flying capacity of the winged Phylloxera, 70—The	
	male louse, 71—Remedies, 71—Efficacy of carbolic acid and soot,	
	71—Value of submersion or irrigation, 72—Sprinkling with quick-	
	lime, ashes, etc., 72—Mr. Lichtenstein's experiments to allure	
	the lice, 72—Experiments with carbolic acid, 73.	73
	THE OYSTER-SHELL BARK-LOUSE OF THE APPLE, Mytilaspis pomorum	13
	Its occurrence in Missouri, 74—Its appearance in Luray County,	
	Missouri, 74—Mr. Hanan's account of its spread, 75—Its occur-	
	rence in Southern Missouri, 76—Its occurrence in Mississippi and	

Georgia, 78—Its appearance in Kansas, 79—It is double-brooded

in the South, 79-The waxy secretion of Homoptera, 80-The newly hatched louse, 80—The larval scale, 81—Development of the female scale, 81—Growth of the male scale, 82—The male louse, 83-Rare occurrence of the winged male, 84-Agamic multiplication, 85-Mode of spreading, 85-Food-plants, 86-Varieties of the apple-tree preferred by it, 86-Enemies and parasites, 87; mites, 87; Aphelinus mytilaspidis and Dr. Le Baron's account of its habits, 88-Easy transportation and introduction of the Aphelinus, 90—Remedies, 90—Application of oily substances, 90—Bibliographical and descriptive, 91—The generic name, 91— Signoret's classification of the Coccide, 92-Specific name, 92-Characteristics and habits of three allied species, 93—A new name necessary for our apple-tree species, 94—Description of the eggs and the winged male, 95; of the male and female scale and of the female louse, 96. THE PINE-LEAF SCALE-INSECT, Chionaspis pinifolii 97 Nature of the malady caused by it, 97—Natural history of the insect, 98—The male, 99—There are two annual broods, 99—It is confined to the pines proper, 100-Natural encmies, 100; the twice-stabbed lady-bird, 100; the painted lady-bird and description of its larva, 101—Remedies, 101—Stripping the old leaves, 102—Application of powdered and liquid substances, 102. THE HICKORY BARK-BORER, Scolytus 4-spinosus 103 Accounts of the damage caused by it in Missouri, 103-Habits of the European Scolytus destructor, 104—Various kinds of hickory attacked by the hickory bark-borer, 105—Its natural history, 105—Natural enemies, 106—Description of the three-banded Spathius, 106—Mr. Cresson's description of Bracon scolytivorus, 106—Remedies, 107—Description of the imago, 107—Is it different from Scolytus 4-spinosus? 107. THE ROSE CHAFER, Macrodactylus subspinosus..... 108 Great damage caused by the beetle in 1872, 108—Remedy, 109— Harris's account of its natural history, 109.

Harris's account of its natural history, 109.

THE FALSE CHINCH-BUG, Nysius angustatus

It was not known as injurious before 1872, 111—Accounts of injury done by it, 111—How it differs from the true chinch-bug, 112—Its probable natural history, 112—Description of imago, larva,

done by it, 111—How it differs from the true chinch-bug, 112—
Its probable natural history, 112—Description of imago, larva, and pupa, 113—Variation in the imago, 113—Its great abundance in the fall of 1872, 114.

INSECTS INJURIOUS TO THE GRAPE-VINE

111

116

117

The grape-vine tomato-gall, Lasioptera vitis.

Its curious resemblance to a tomato, 117—Various shapes assumed by it, 118—The larva, 118—Enemies of the larva, 118.

^{*} Pages 119-125 were reprinted under title Insect punctures, in < Rept. State Board Agric. Kansas for 1873, 1874, pp. 274-279, figs. 47-59.

1329.	RILEY, C. V.—Continued.	
	Probable eggs of the jumping tree-cricket, Orocharis saltator, 119-	
	The egg-punctures on grape-cane, 119—Characters of the egg,	
	119—General appearance of the cricket, 120.	
	Eggs of the snowy trec-cricket, <i>Ecanthus niveus</i> , 120—Trees and	
	shrubs attacked by this cricket, 120—The egg, 120—Habits and	
	natural history, 120-Injury done by it, 121-The eggs mistaken	
	by Fitch for those of the Buffalo tree-hopper, 121.	
	Egg-punctures of the Buffalo tree-hopper, Ceresa bubalus, 121—De-	
	velopment of the larva, 121—Characters and habits of the per-	
	feet insect, 122—Egg-punctures of some unknown tree-hopper, 122.	
	Egg-punctures of the frosted lightening-hopper, Paciloptera prui-	
	nosa, 122—Development and habits of the insect, 122.	
	Egg-punctures probably of Orchelimum glaberimum, 123—The egg,	
	123—Characters and habits of the imago, 123.	
	Eggs of the oblong-winged katydid, Amblycorypha oblongifolia,	
	123-They have been mistaken by Harris for those of the broad-	
	winged catydid, 123—Ovipositors and modes of egg-laying in	*
	the three katydids occurring in Missouri, 124-Increase in thick-	
	ness of katydid eggs before hatching, 124.	
	STINGING LARVÆ	125
	The potato-worm falsely considered as a stinging caterpillar, 125—	
	General harmlessness to man of insect larvæ, 125—Stinging cat-	
	erpillars in Europe and North America, 126—List of such larvæ	
	in United States, 126—Description of larva and chrysalis of	
	Acronycta xylinoides, 126.	
	The buck moth or maia moth, Hemileuca maia	127
	Flight and characteristics of the moth, 127—The egg and mode of	
	egg-laying, 128—Appearance of the full-grown larva, 129—	
	Larval changes, 129—Color variations of the larva, 130—Habits	
	of the young larva, 130—The sting of the larva, 131—The pupa,	
	131—Issuing of the moth, 132—Irregularity and retardation in	
	development, 132; the species benefits thereby, 132-Food-plants,	
	132—Natural enemies, 132.	100
	The io moth, Hyperchiria io	133
	The male and female moth, 133—The egg, 134—The larva and its	
	urticating power, 134—Larval changes, 135—Pupation, 135—	
	Food-plants, 136—Parasites, 136.	137
	THE GREEN-STRIPED MAPLE-WORM, Dryocampa rubicunda	101
	done by it to soft maples, 138—The egg, 138—Larval changes,	
	138—The chrysalis, 139—The imago, 139—Natural enemies, 139—	
	Description of Belvoisia bifasciata, 140—Remedies, 141.	
	Description of Decouse of asciata, 140—Itemetics, 141.	
	INNOXIOUS INSECTS.	
	THE HELLGRAMMITE FLY, Corydalus cornutus	143
	The eggs, 143—Respiratory apparatus of the larva, 143—Habits of	
	the larva when about to transform, 143—The pupa, 144—Sexual	
	difference in the imago, 145.	
	THE GOAT-WEED BUTTERFLY, Paphia glycerium	145
	The egg, 146—Habits of the larva, 146—Larval changes, 146—	
	Transformation, 147—New food-plant, 147—Simulation of the	
	color of the food-plant by the larva, 147—Two annual broods,	
	148—Hibernation of the imago, 148—Natural enemies, 149.	

ON A NEW GENUS IN THE LEPIDOPTEROUS FAMILY TINEIDÆ WITH REMARKS ON THE FERTILIZATION OF YUCCA, Pronuba yuccasella......

Generic characters of Pronuba, 150—Description of Pronuba yuccasella, 151—Plants requiring the aid of insects for fertilization,
152—Fertilization of orchids, 152—Fructification of Yucca, 153—
Yuccas must rely on insects for fertilization, 153—Insects frequenting Yuccas, 154—Diurnal and nocturnal habits of Pronuba,
154—How the female moth fertilizes the plant, 154—Oviposition,
155—The larva within the young fruit, 155—Description of the
larva, 155—Only a small percentage of fruit not infested by the
larva, 156—The larva leaves the capsule and enters the ground
for hibernation, 156—Mutual adaptation of plant and insect,
156—The moth doubtless occurs wherever Yuccas grow wild,
157—Easy transportation of the cocoon, 158—Further facts re-

Note.—Pages 1-44 were also printed as a separate, with title-page cover.

Solution City, Mo., 1873.
S.-b. No. 19, pp. 150-173.

garding the fructification of Yucca filamentosa and gloriosa, 158—Yuccas seeding in Europe, 159—Range of the insect, 159—Mr. Stainton's opinion on the characters on the Yucca moth, 160.

1330. RILEY, C. V. New York without a State entomologist. < Moore's Rural New Yorker, 5 May, 1873. S.-b. No. 8, p. 103.

Needs of a State entomologist for New York; Salix humilis infested with larva of Plectrodera scalator.

1331. RILEY, C. V. Tent-caterpillar of the forest. <N. Y. Tribune, 23 May, 1873. S.-b. No. 8, pp. 64-65.

Answer to inquiry of S. T. Gilbert; means against eggs of Clisiocampa sylvatica [= disstria].

1332. RILEY, C. V. Apple-tree borer. <N. Y. Tribune, 23 May, 1873, S. b. No. 8, p. 64.

Answer to inquiry of J. Durbin; means against Saperda bivittata [= candida].

1333. RILEY, C. V. Punctured grape-canes. <N. Y. Tribune, 23 May, 1873. S.-b. No. 8, p. 64.

Answer to inquiry of E. Snyder; food-habits and means against *Œcanthus* niveus.

1334. RILEY, C. V. The codling moth. Weir's trap. Amer. Agric., May, 1873, v. 32, p. 184, figs.

Unsatisfactory results of experiments with the Weir trap; relative value of materials used; criticism of J. S. Parker's theory for the extermination of Carpocapsa pomonella; this species breeds in apples, wild crabs, pears, peaches, and plums; figures females and the abdomen of the males of Pimpla annulipes and Macrocentrus delicatus parasitic upon the apple-worm.

1335. RILEY, C. V. Influence of extreme cold on the Curculio. <Gardener's Mo. and Hortic., May, 1873, v. 15, pp. 137-139.

Critical review of T. T. Southwick's article of same title; extract from 3d Ann. Rept. State Ent. Mo.

150

- 1336. RILEY, C. V. On a new genus in the lepidopterous family Tineidæ, with remarks on the fertilization of Yucca. <Trans.
 Acad. Sci. St. Louis, June, 1873, v. 3, pp. 55-64, 2 figs. Reprinted, with omissions and slight changes, from <5th Ann.
 Rept. State Ent. Mo., 18 April, 1873, pp. 150-160, figs. 74-75.
 Review by P. C. Zeller: <Verh. k.-k. zool.-bot. Ges. Wien,
 1876, Jahrg. 1875, Bd. 25, pp. 340-342. Reply to Zeller:
 <Trans. Acad. Sci. St. Louis, January-March, 1876, v. 3, pp.
 325-326.
 - Description and figures of *Pronuba* n. g. and of larva and image of *P. yuccasella* n. sp. See No. 1329 for synopsis of contents.
- 1337. RILEY, C. V. Supplementary notes on *Pronuba yuccasella*. <a href="mailto: Trans. Acad. Sci. St. Louis, June, 1873, v. 3, pp. 178–180, fig. 3. Extract: –135, fig. 3.
 - Descriptions and figures of male and female pupa of *Pronuba yuccasella*; hibernation, imagination, seasons, and distribution of the same; species of *Yucca* pollinated by it.
- 1338. RILEY, C. V. [To destroy the cotton-worm.] <Ill. Jour. of Agric., June, 1873. Reprint: <Colman's Rural World, 1873; <Rural Alabamian, July, 1873, v. 2, pp. 289–293; <Mobile Register, 1873; <Farmer's Advocate, 1873. Extract: <6th Ann. Rept. State Ent. Mo., 1874, pp. 17–18.
 - Ravages of Aletia argillacea [=xylina] in the cotton fields of Southern United States; inefficiency of measures hitherto employed against them; recommends the use of Paris green; directions for its use; natural history, description of egg, larva, and image of the Aletia; hibernation of the image.
- 1339. RILEY, C. V. [Imported plants and insects.] <Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 42-43 Proc.
 - Verbal communication; inequality of the exchange of plants and insects between Europe and North America; extent and causes of the same.
- 1340. RILEY C. V. [Mimicry and protective resemblances.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 44-45 Proc.
 - Verbal communication; mimicry of Danais archippus by Limenitis disippus and consequent greater abundance of the latter than of L. ursula.
- .1341. RILEY, C. V. [Silk-worms fed with osage orange.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 47 Proc.
 - Verbal communication; successful rearing of silk-worms on Maclura aurantiaca; exhibition of cocoons made by these worms.
- 1342. RILEY, C. V. [On the cause of deterioration in some of our native grape-vines.] <Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 51-52 Proc.
 - Verbal communication; abstract from 4th Ann. Rept. State Ent. Mo.; failure of grape-vines attributed largely to the ravages of *Phylloxera vastatrix*.
- 1343. RILEY, C. V. [Insects affecting the ailanthus.] <Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 53-54 Proc.

 Mentions Œta compta [= punctella] and Attacus cynthia.

- 1344. RILEY, C. V. [Posthumous papers by B. D. Walsh.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 77 Proc.
 - Verbal communication; disposition of sundry entomological manuscripts left by B. D. Walsh.
- 1345. RILEY, C. V. [Remarks on Simulium piscicidium.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 79 Proc.
 - Verbal communication; nature and supposed ravages of Simulium piscicidium.
- 1346. RILEY, C.V. [On Antherwa yama-mai as a silk-producer.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 84 Proc.
 - Verbal communication; superiority of Antherwa yama-mai to other silkworms tried as substitutes for Sericaria mori.
- 1347. RILEY, C. V. [On galls growing on wild sage.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 84 Proc.
 - Verbal communication; occurrence of three distinct undescribed galls on Artemisia tridentata in Utah.
- 1348. RILEY, C. V. [On a larva of *Scenopinus* sp. from the human lungs.] <Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 90
 Proc.
 - Verbal communication; a larva of *Scenopinus* sp. found in human expectoration; usual food-habits of the larvæ of this genus.
- 1349. RILEY, C.V. Agricultural editorial excursion. <Prairie Farmer, 1873, v. 44: 2 August, p. 241; 9 August, p. 248; 16 August, p. 256; 23 August, p. 265; 30 August, p. 273; 6 September, p. 281. Reprint: <Colman's Rural World, 1873, 2, 9, 16, 23, and 30 August, 6 and 13 September. See: <Colorado Mountaineer, 1 August, 1877. S. b. No. 14, p. 102.

 Brief notes on locusts.
- 1350. RILEY, C. V. Enemies of the elm. <N. Y. Tribune, 7 August, 1873. S.-b. No. 8, pp. 56-57. Reprint: <Gardener's Mo. and Hortic., August, 1876, v. 18, p. 246.
 - Answer to inquiry of T. S. Watson; natural history of and means against Galeruca calmaricus [= xanthomelæna].
- 1351. RILEY, C. V. Entomological information. <N. Y. Tribune, 16
 August, 1873. S.-b. No. 8, p. 58.
 - Answer to inquiry of J. W.; description of a number of traps for alluring and destroying insects.
- 1352. RILEY, C. V. "Controlling sex in butterflies." <Amer. Nat., September, 1873, v. 7, pp. 513–521. Separate: <Salem, Mass., August, 1873, pp. 9.
 - Critical review of Mary Treat's article of same title; females require more nourishment than males; sex determined in the egg; oviposition of Papilio and Anisota [= Dryocampa]; larvæ can not be forced to eat more than is natural to them; experiments on Thyridopteryx ephemeræformis, Orgyia leucostigma, Clisiocampa americana, Hyperchiria io, Hemileuca maia, and Anisota [= Dryocampa] rubicunda to determine the effect of the stinting of food upon the determination of sex; effect of this upon the number of molts and the longevity of Orgyia leucostigma and Megatoma serra; note on alternation of generations in Cynips [= Andricus] quercus-operator and C. [=A.] quercus-operatola; nature of parthenogenesis.

- 1353. RILEY, C. V. Cotton caterpillar.—Boll-worm. <Rural Alabamian, October, 1873. S.-b. No. 16, p. 121.
 - Criticism of recent articles on Anomis [= Aletia] xylina, Heliothis armigera, and Phylloxera vastatrix.
- 1354. RILEY, C. V. On the oviposition of the Yucca moth. <Amer. Nat., October, 1873, v. 7, pp. 619-623. Abstract: <Trans. Acad. Sci. St. Louis, 10 December, 1873—25 April, 1874, v. 3, pp. 208-210. Reprint, with slight changes and omissions. <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 131-135, fig. 38.

See No. 1363 for synopsis of contents.

- 1355. RILEY, C. V. Phylloxera: correction. < Gardener's Mo. and Hortic., November, 1873, v. 15, p. 342.
 - Critical review of report of remarks before Academy of Natural Sciences of Philadelphia.
- *1356. RILEY, C. V. Hackberry butterflies. Descriptions of the early stages of Apatura lycaon, Fabr., and Apatura herse, Fabr.; with remarks on their synonymy. <Trans. Acad. Sci. St. Louis, 10 December, 1873, v. 3, pp. 193–208, figs. 3–6. Reprint, with slight changes. <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 136–148, figs. 39–42.

Treats of A. lycaon [=celtis] and A.herse [=clyton]. See No. 1363 for synopsis of contents.

1357. RILEY, V. C. Economic entomology. <Trans. Kansas State Board Agric. for 1872, 1873, pp. 292–325, 18 figs. Extract: <Sci. Amer.

Characterization of entomology; importance of and means against noxious insects; list of imported noxious insects and plants; chapters on Anisota [=Dryocampa rubicunda], Eriosoma pyri[=Schizoneura lanigera], Paleacrita vernata, Galleria cereana, Œstrus ovis, Bruchus pisi, Macrodactylus subspinosus, Conotrachelus nenuphar, and Nysius destructor [=angustatus]. In general, descriptions and figures of the several stages and accounts of the habits and food-plants of and means against these insects, with some accounts and figures of their enemies and parasites, are given. Describes especially larva of Anisota [=D.] rubicunda and of Paleacrita vernata and of all active stages of Nysius destructor [=angustatus]. Figures larva, puparium, and imago of Pipiza radicum, imagos of Nothrus ovivorus, Microgaster [=Apanteles] militaris, Calosoma scrutator, C. calidum, and Blissus leucopterus and imago and nests of Eumenes fraternus.

Some of the chapters are based upon and some are reprinted from the 1st, 2d, and 3d Ann. Repts. State Ent. Mo. '

- 1358. RILEY, C. V. Curculios on pears. <Ill. Journ. Agric., 1873. S.-b. No. 16, pp. 109–110, figs. 1–2.
 - Habits, history, and means against Conotrachelus nenuphar, C. cratægi, and Anthonomus quadrigibbus; figures the two last-named species.
- 1359. RILEY, C. V. Length of thread of the silk-worm. < Popular Sci. Monthly, February, 1874, v. 4, p. 508.

Correction of statements in A. de Quatrefage's "Silk-worms and sericulture" as to the length and weight of the fiber in a single cocoon of Sericaria mori.

1360.	RILEY, C. V. A new (?) ægerian maple-borer. <amer. n<br="">February, 1874, v. 8, pp. 123-124.</amer.>	
	Critical review of P. Gennadius's "A new ægerian maple-borer." Trochi acericolum is the well-known Ægeria acerni, injurious to maple-trees.	lium
1361.	RILEY, C. V. Entomology in Missouri. < Amer. Nat., Mar. 1874, v. 8, pp. 181–188.	·
	Reply to criticisms of A. S. Packard; discussion on the number of segm in the head of an insect, on classification, and on <i>Mytilaspis pomicor</i> [=pomorum].	
1362.	RILEY, C. V. Economic entomology. <amer. 18="" 189-190.<="" 8,="" march,="" nat.,="" pp.="" td="" v.=""><td></td></amer.>	
	Demand for more attention to and better representation of the agricult interests of the country. Inadequacy of measures propounded for wholesale destruction of noxious insects.	
1363.	RILEY, C. V. Sixth annual report on the noxious, beneficial, other insects of the State of Missouri, made to the State Bo of Agriculture, pursuant to an appropriation for this purp	ard
	from the legislature of the State. <9th Ann. Rept. State Bo of Agric. for 1873, March, 1874, pp. 169+12, 55 figs. Separate	ard
t	<jefferson 169+12,="" 1874,="" 55="" city,="" figs.<="" march,="" mo.,="" p="" pp.=""> Table of contents</jefferson>	3
	Preface	6
	NOXIOUS INSECTS.	
	Notes of the Year The codling-moth, Carpocapsa pomonella Failure of the apple crop in 1873, 9—Dr. Le Baron's observations on the habits of the worm, 9—Proportion of worms leaving the fruit before it falls, 10—How it affects pears, 10—Time and method of using bandages, 10—Westward spread of the insect, 10.	. 9
	The Colorado potato-beetle, Doryphora 10-lineata. How it has affected the price of potatoes, 11—New food-plants, 11—Its progress eastward during 1873, 12—Improved methods of using Paris green, 13—Device for jarring off the bugs, 14—European publications on the insect, 15—Danger of its introduction into Europe, 15—Precautionary measures to be taken in Europe, 16.	
	The cotton-worm, Aletia xylina Paris green suggested as remedy, 17—Address before the National Agricultural Congress, 17—Mr. Glover's summary on experience with Paris green, 19—Experiments with the poison, 19—Johnson's sprinkling machine, 20—Patents on Paris green, 20—The Royal mixture, 21—Hibernation of the insect, 22—Natural enemies, 23—Geographical range, 23—Position of the moth when	17
	alighting, 24. The canker-worm, Paleacrita vernata; Anisopteryx pometaria Dr. Le Baron's summary of remedies, 24—Mr. Milliken's experience with the rope and tin trap, 25—The Paris green remedy, 26—A new trough, 26—Birds which destroy the worm, 27—Mr. Mann's observations on the insect, 28—Two species have hitherto been confounded, 28—The English sparrow and the increase of the	24

white-marked tussock-moth, 29.

The term "Phylloxera," 30—Bibliographical history, 30—Characters of the genus and its position in the system, 33-Biological history, 33-Different forms which the insect assumes, 33-The gall-inhabiting form, 34—The root-inhabiting form, 38—Balbiani's discovery of the true sexual individuals and the winter egg of Phylloxera quercus, 41—The winter egg not essential to the winter life of the grape Phylloxera, 42-Polymorphism in Phylloxera, 43—Conclusive proof of the identity of the root- and galllice, 44—Practical considerations, 44—The more manifest and cxternal effects of the disease, 44-Mode of spreading, 45-Swarming of winged specimens of Phylloxera caryafolia, 45-Power of flight in the grape Phylloxera, 45-Spread of the disease through the winged females, 46-Susceptibility of different vines to the disease, 46—Practical importance of knowing the more resisting and more susceptible vines, 46-List of the cultivated species and varieties of vines indicating their relative susceptibility to Phylloxera, 47—Prophylactic means of coping with the disease, 48 Grafting the more susceptible on to the roots of resisting vines, 48—Other preventive measures, 50—Natural encmies, 50— Insect enemies of the gall-louse, 50—Enemies of the root-louse, 52—The Phylloxera mite, 52—Different forms assumed by mites in their development, 52-Hoplophora arctata, 53-Peculiarities of the Oribatidæ, 54-Direct remedies, 55-Submersion a perfect remedy, 55-Application of pure insecticides without satisfactory results, 56—Range of the insect in America, 57—Injury caused by it in America, 58-Reasons why the insect is more injurious in Europe, 59-False theories, 60-The Phylloxera is the cause and not the effect of the disease, 60-It is a native American insect aud has been imported iuto Europe, 62-It is peculiar to the grape-vine, 63-The American oak Phylloxera and its natural history, 64-Conclusion, 65.

Appendix and notes to the article on grape Phylloxera.....

Diagnosis of Phylloxera vastatrix, 66—Influence of the insect's puncture, 67—The supposed male of the gall-louse, 67—The noncultivation of the Clinton grape, 67-Transient nature of the leaf-galls, 68—Constancy of the differences between the forms assumed by the insect, 68-Supposed sexual individuals, 68-Number of generations annually produced, 69—Number of molts, 69—Transplanting root-lice on to the leaves, 69—Nature of the swelling on the roots produced by Phylloxera, 70-The true grape-viues of the United States, by Dr. G. Engelmann, 70-Exceptional instances where the European vine has succeeded in America, 76—Grafting the more susceptible on to the roots of the resisting varieties, 78—Descriptions of Tyroglyphus phylloxera and Hoplophora arctata, 81—Efficacy of inundating the vineyards, 82—Facts showing that the disease of grape-vines in America is principally caused by Phylloxera, 82-Description of Phylloxera rileyi, 86—The true sexual individuals and the winter egg, 86.

characters of the larva, 87—Description of larva and chrysalis, 88.

2. The beautiful wood nymph, Eudryas grata, 88—Characters and food-plants of the larva, 88—The moth and the egg, 89—Description of the egg and larva, 89—Of the chrysalis, 90.

66

176	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	.*
1363	RILEY, C. V.—Continued.	
1000.	Insects injurious to the Grape-vine—Continued. 3. The pearl wood nymph, Eudryas unio, 90—Its larva almost undistinguishable rom that of the foregoing species, 90—Foodplant, 91—Distinguishing characters of the moth, 91—Mr. Lintner's description of the larva, 92—Mr. Lintner on the difference between the larva of Eudryas grata and E. unio, 93. 4. The eight-spotted forester, Alypia octomaculata, 94—Description of the larva, 94; chrysalis and imago, 95. Summary, 95—Comparison of the larvæ of these four species, 95—	
	Remedies 96.	
	The Red-Legged Ham-beetle, Necrobia rufipes Its popular name, 96—It has been the cause of an interesting lawsuit, 96; entomological information necessary to a just verdict, 98—Another case before a jury requiring botanical knowledge, 98—Injury caused by the beetle in St. Louis, 99—The eggs, and how they are deposited, 99—The larva and its habits, 99—The pupa, 99—Prevention, 100—Other species associated with it, 100—Habits of other species of Clerida, 101—How Necrobia ruficollis saved the life of Latreille, 101—Description of the larva of Necrobia rufipes, 101—Description of the pupa, 102.	96
	THE CLOVER-HAY WORM, Asopia costalis Its geographical distribution, 102—It has probably been imported from Europe, 101—Its past history and accounts of damage caused by it, 102—Its natural history, 105—Remedies, 105—Description of larva and chrysalis, 106; of the imago, 107—Its allied congener, Asopia olinalis, 107.	102
	The Legged Maple-Borer, Ægeria acerni The burrows of the larva, 108—Appearance of the moth, 108—Remarks on the nomenclature of the species, 108—Injury caused by it to maples, 109—It attacks only trees already injured by some cause or another, 109—Remedies, 110—Description of imago, larva, and pupa, 110.	107
	THE RASPBERRY ROOT-BORER, Bembecia marginata Work of Oberea perspicillata in canes of raspberry and blackberry, 111—Injury caused by the root-borer, 111—General appearance and habits of the worm, 112—Description of imago and larva, 113.	111
	The Northern Brenthian, Eupsalis minuta. Its occurrence and distribution, 113—Characteristics of the beetle, 114—Fighting habits of the males, 114—How the male assists the female in ovipositing, 115—How much time is required for the transformations of the insect, 115—Habits of the larva, 115—Description of the larva, 115; of the pupa, 116—Real position of the brenthians in the system, 116—The specific and generic	113

names, 116—How another larva has been mistaken for that of the Northern brenthian, 117—Description of this larva, which evidently belongs to the Tenebrionidæ, 118.

The Jumping sumach-beetle, Blepharida rhois

Sumach industry in Europe and America, 118—Species of sumach possessing economic value, 119—Appearance and habits of the beetle, 119—The egg-masses, 120—Development and habits of the larva, 120—Transformation, 120—The two annual broods,

118

120—Injury caused by it to the sumachs, 121—Geographical range, 121—Remedies, 121—Description of larva and pupa, 121—Variations of the beetle, 122.

BENEFICIAL INSECTS.

The unadorned Tiphia or white grub, 123—Cocoon and larva of Tiphia, 123—It is undoubtedly a parasite of the white-grub, 124—Characters of the genus Tiphia, 124—Habits of Tiphia femorata, 124—Habits of the Tiphia larva, 124—Characters of the Unadorned Tiphia, 125—Bectle parasitic upon it, 125—Description of larva and imago, 126—The species has been described under three different names, 126.	123
INNOXIOUS INSECTS.	
THE DOMINICAN CASE-BEARER, Coscinoptera dominicana. Characters of the beetle, 127—Mode of oviposition, 128—Oviposition of Chlamys plicata, 128—The young larva and its foodhabits, 129—The case of the larva, 129—Characters of the larva, 130—Time required for its development, 130.	127
THE YUCCA MOTH, Pronuba yuccasella Its natural history completed, 131—Description of the ehrysalis, 131—Hatching of the ehrysalis, 132—Method of oviposition, 133—Oviposition is followed by pollination, 134—The egg in the young fruit and the embryo larva, 134—It is the only insect that can well fertilize the Yuccas, 135.	131
HACKBERRY BUTTERFLIES. The European purple Emperor, 136—Insufficient former account of	136
their earlier states, 136—Species of Celtis in Missouri, 137.	4.00
The eyed Emperor, Apatura celtis	137
The tawny Emperor, Apatura elyton. How it differs from the eyed emperor, 140—The egg, 141—Larval changes and habits, 141—Egg parasite, 142. Bibliographical	140 142
The generic name, 142—The specific names of the two species, 143—Fabricius's original description of lycaon and herse, 144—Other species of the genus in the United States, 145—On the validity of alicia, 145—Descriptions of the earlier states of both species, 146—Their popular names, 148—The scientific specific names, 149.	192
KATYDIDS	150
General considerations, 150—Spring in Europe and America, 151—Stridulating noise produced by crickets and grasshoppers, 152—Sounds inaudible to man, 152—Grasshoppers, katydids, locusts, 153—Habits and general appearance of our katydids, 154—They oviposit above ground, 154—Their ovipositors not rudimental, 155.	
The angular-winged katydid, Microcentrum retinerve	155
It is the most common species in Missouri, 155—General character of the insect, 155—The eggs, 155—Erroneous statement concerning the egg, 156—How the female deposits the eggs, 156—Number of eggs laid by the female, 158—Hatching of the larva, 158—Food of the larva, 158—Changing from the pupa to the perfect 12 ENT	100
LA LIVI	

1363.	RILEY, C. V.—Continued.	
	state, 159—Its song, 159—It is capable of domestication to a cer-	
	tain degree, 160—Description of the immature states, 161—Nat-	
	ural enemies, 162—The back-rolling Wonder, an egg-parasite,	
	162—Curious habit of the female Antigaster, 162—Description of	
	Antigaster mirabilis, 163—Striking sexual differences, 163.	
	The narrow-winged Katydid, Scudderia curvicauda	164
	Distinguishing characters, 164—Method of oviposition, 165—The	
	egg, 165—Its song, 166—Description of larva and pupa, 166.	
	The broad-winged Katydid, Cyrtophyllus concavus	167
	It is the true katydid, 167—Distinguishing characters, 167—Mode	
	of oviposition as observed in confinement, 167—Mr. Jaeger's	
	erroneous statement regarding the oviposition, 168—Its song,	
	168.	
	The oblong-winged katydid, Amblycorypha oblongifolia	169
	It has not yet been found in Misscuri, 169—Distinguishing charac-	
	ters, 169—The eggs not yet known, 169—Abnormal female speci-	
	men, 169.	

1364. RILEY, C. V. The Colorado potato-beetle abroad. <N. Y. Tribune, 1 April, 1874. S.-b. No. 10, p. 37.

Criticism of some articles on *Doryphora decemlineata* published in Europe; danger of the importation of the insect into Europe, and suggestion of means to prevent such introduction.

1365. RILEY, C. V. Cabbage-lice. <N. Y. Tribune, 8 April, 1874. S.-b. No. 10, pp. 33-34.

Answer to letter from Mrs. M. Walker; means against Aphis brassica.

1366. RILEY, C. V. Peach-borers. <N. Y. Tribune, 8 April, 1874. S.-b. No. 10, p. 34.

Answer to letter from W. J. Clary; a peach-tree whose base was surrounded by ice in winter was free from borers; this observation of slight significance.

1367. RILEY, C. V. Apply soap. <N. Y. Tribune, 8 April, 1874. S.-b. No. 10, p. 34.

Answer to letter from T. H. Wakeley; means against Saperda bivittata [=can-dida] and Chrysobothris femorata.

1368. RILEY, C. V. Meadow enemy. <N. Y. Tribune, 8 April, 1874. S.-b. No. 10, p. 34.

Answer to letter from D. Freeman; ravages of a species of *Tipula* in grass lands in California; characters of the genus; habits of and means against the same.

1369. RILEY, C. V. A remedy for the cotton-worm. <N. Y. Tribune, 22 April, 1874. S.-b. No. 10, pp. 25–26. Reprint: <Vicksburg Herald, 1 May, 1874.

Experiments with Paris green; advocacy of its use against Aletia argillacea [=xylina]; directions for and successful results of this use.

1370. RILEY, C. V. Descriptions of two new subterranean mites. <Trans. Acad. Sci. St. Louis, 25 April, 1874, v. 3, pp. 215-216, figs. 8-9.

Description and figures of Tyroglyphus phylloxeræ n. sp. [p. 215] and of Hoplophora arctata n. sp. [p. 216]; food-habits of the same.

- 1371. RILEY, C. V. The habits of *Polistes* and *Pelopæus*. < Amer. Nat., April, 1874, v. 8, pp. 229-231.
 - Critical review of P. R. Uhler's article "On a remarkable wasp's nest found in a stump in Maryland;" probably the nests and specimens of Pelopaus lunatus [= cementarius] were mistaken for those of Polistes fuscatus [= metricus], in which case the observations criticised present no remarkable features.
- 1372. RILEY, C. V. The plum Curculio; natural history and how to catch him. <N. Y. Semi-weekly Tribune, 1 May, 1874. S.-b. No. 10, pp. 18–19; 38–40. Extract: <Cultivator and Country Gentleman, 14 May, 1874, v. 39, p. 310. <New England Farmer, 18 July, 1874, [v. 53,] n. s., v. 29, p. 1. See: <Indust. Record, 5 February, 1875. S.-b. No. 16, pp. 46–48.
 - Natural history of and means against Conotrachelus nenuphar; figures of the several stages of the insect and of machines for catching the same.
- 1373. RILEY, C. V. The apple-worm; natural history; remedies. <N. Y. Tribune, 20 May, 1874. S.-b. No. 10, pp. 8-9; 37-38.

 Description and figures of the several stages of Carpocapsa pomonella; habits of and means against the same.
- 1374. RILEY, C. V. The grape Phylloxera. < Popular Sci. Monthly, May, 1874, v. 5, pp. 1-16, 7 figs. Reprinted, with slight changes, from <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 30-46.

 See No. 1363 for synopsis of contents.
- 1375. RILEY, C. V. Rose chafers on grape-vines. <Colman's Rural World, 20 June, 1874, fig. S.-b. No. 17, pp. 103, 104.
 - Natural history and means against Macrodactylus subspinosus; figure of the same.
- 1376. RILEY, C. V. More about the grape vine pest. < Popular Sci. Monthly, June, 1874, v. 5, pp. 158-170, 10 figs. Reprinted, with changes and omissions, from <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 47-60, 64-65.
 - See No. 1363 for synopsis of contents.
- 1377. RILEY, C. V. Scale insects on magnolia. <N. Y. Tribune, 15 July, 1874. S.-b. No. 10, p. 24.
 - Answer to inquiry of R. H. Day; occurrence of an undescribed *Lecanium* sp.? on the leaves of *Magnolia fuscata*.
- 1378. RILEY, C. V. Confounding friend with foe. <N. Y. Tribune, 15 July, 1874, 2 figs. S.-b. No. 10, p. 24.
 - Answer to inquiry of E. J. Day; usefulness of Arma [= Podisus] spinosus figure of the same. Injury to plums by Anthonomus prunicida [= Coccotorus scutellaris]; figure of and means against the same.
- 1379. RILEY, C. V. The Colorado potato-beetle in New York. <N. Y. Tribune, 15 July, 1874. S.-b. No. 10, p. 24.
 - Answer to letter from J. H. P.; eastern limit of the range of Doryphora decemberata in 1873 and 1874.

- 1380. RILEY, C. V. Large willow-worm. <N. Y. Tribune, 15 July, 1874. S.-b. No. 10, p. 24.
 - Answer to letter of J. H. P.; food-plants, habits, and means against Cimber laportei [= americana].
- 1381. RILEY, C. V. Black blister beetles on potatoes. <N.Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to inquiry of A. Barr; ravages of and means against *Epicauta puncti-* collis.
- 1382. RILEY, C. V. Pear-tree slug. <N. Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to letter of C. S. B.; means against Selandria [= Eriocampa] cerasi.
- 1383. RILEY, C. V. The plug-ugly theory. <N. Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to letter of O. J. B.; insertion of sulphur, calomel, and similar powders in the trunks of trees has no effect upon the sap or upon the insects which injure the trees.
- 1384. RILEY, C. V. Cockscomb elm-gall. <N. Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to inquiry of J. S. Ruby; Thelaxes [= Colopha] ulmicola forms excrescences on the American elm.
- Courant, 15 August, 1874, v. 38, No. 195, p. 1. See: <N. Y. Tribune Extra: Lecture and Letter series No. 21, August, 1874, pp. 56–58, fig. Reprint: <Nature, 8 October, 1874, v. 10, pp. 463–465, figs. 1–2. Abstract: <Sci. Amer., 12 September, 1874 [v. 45], n. s., v. 31, p. 168. <Amer. Nat., November, 1874, v. 8, pp. 684–687. Reprint of abstract: <Hardwicke's Science Gossip, December, 1874, v. 10, pp. 272–275, figs. 179–182. Extract: <Ca. Ent., November, 1874, v. 6, pp. 207–214, figs. 25–26. Reprint of extract: <Proc. Amer. Assoc. Adv. Sci. for 1874, June, 1875, v. 23, pp. 18–25, 2 figs. Separate of reprint: <Salem, Mass., December, 1874, pp. 18–25, figs.
 - Insect-catching habits of Sarracenia variolaris; list of its victims; habits and figures of all stages of Xanthoptera [= Exyra] semicrocea and Sarcophaga sarracenia: capture of insects by other plants.
- 1386. RILEY, C. V. On the habits and transformations of Canthon hudsonias, Forst.; the common "tumbledung." <Hartford [Conn.] Daily Courant, 18 August, 1874, No. 11162, v. 38, No. 197, p. 2. Reprint: <N. Y. Tribune Extra: Lecture and Letter series, No. 21, August, 1874, pp. 75-76.
 - Breeding habits and oviposition of Canthon hudsonias [= lævis].
- 1387. RILEY, C. V. On the larval habits of the cantharid genera *Epicauta* and *Henous*. <Hartford [Conn.] Daily Courant, 18 August, 1874, No. 11162, v. 38, No. 197, p. 2. See: <N. Y. Tribune Extra: Lecture and Letter series, No. 21, August, 1874, p. 76.
 - Habits and hypermetamorphosis of Meloe angusticollis; probably parasitic food-habits of young larve of Epicauta and Henous; the images phytophagous,

- 1388. RILEY, C. V. Humming-bird moths caught by the tongue. <Moore's Rural New Yorker, 29 August, 1874, v. 30, p. 140.

 Method of capture of insects, particularly moths by the flowers of *Physian-thus albens*; capture of insects by other flowers.
- 1389. RILEY, C.V. The io moth (Saturnia io). <Illust. Journ. Agric., August, 1874, figs.
 - Answer to inquiry of G. Barter; natural history of Saturnia [= Hyperchiria] io; figures larva and \mathcal{J} and \mathcal{D} images.
- 1390. RILEY, C. V. Descriptions and natural history of two insects which brave the dangers of *Sarracenia variolaris*. <Trans. Acad. Sci. St. Louis, 1 September, 1874, v. 3, pp. 235–240, figs. 10–11.
 - Description of leaf of Sarracenia variolaris and its function of entrapping and digesting insects; description and figures of eggs, larva, chrysalis, and imago of Xanthoptera [=: Exyra] semi-crocea and larva, puparium and imago of Sarcophaga sarraceniæ n. sp.; habits of the two species; comparison of S. sarraceniæ with S. carnaria and of the genera Musca, Calliphora, and Sarcophaga; notice of A. S. Packard's "Transformation of the common housefly."
- 1391. [RILEY, C. V.] [Discussion on entomology.] <Trans. Ill. State Hortic. Soc. for 1873, 1874, n. s., v. 7, pp. 100-104.

 Description, habits, and means against Anisopteryx; effect of late plowing; enemies of Doryphora 10-lineata; habits of Gastrophilus equi.
- 1392. [RILEY, C. V.] Note on leaf-hopper. <Trans. Ill. State Hortic. Soc. for 1873, 1874, n. s., v. 7, p. 138.

 Means against Erythroneura [= Typhlocyba] vitis.
- 1393. [RILEY, C. V.] [Notes on the strawberry crown borer.] < Trans.

 Ill. State Hortic. Soc. for 1873, 1874, n. s., v. 7, p. 147.

 Description, habits, and means against Tyloderma fragariæ.
- 1394. RILEY, C. V. Lecture on entomology. <Trans. Ill. Hortic. Soc. for 1873, 1874, n. s., v. 7, pp. 172–178, figs. 1–3. Reprint: <3d Ann. Rept. Sec. State Pomol. Soc. Mich. for 1873, 1874, pp. 443–448. Abstract: <Rept. U. S. Commis. Agric. for 1873, 1874, [30 March, 1875], pp. 389–390.
 - Habits of and means against Carpocapsa pomonella; habits, transformations, prolificacy of and means against Phylloxera vastatrix; description and figures of leaf-galls; figures of root- and gall-forms; history and meaning of the word "Phylloxera;" statement by D. B. Wier concerning the invention and use of Wier's apple-worm trap.
- 1395. RILEY, C. V. "Walking-sticks or specters" becoming injurious. <N. Y. Weekly Tribune, 11 November, 1874, fig. S.-b. No. 23, p. 103.
 - Ravages in Yates County, N. Y., vernacular names, habits of, and means against Diapheromera femorata; description of its eggs; figure of imago.
- 1396. RILEY, C. V. Entomological notes. <Sci. Amer., 5 December, 1874 [v. 45], n. s., v. 31, p. 356.
 - Pemphigus imbricator found on beech; vesicatory potato-bectles, Meloidæ; cow manure and cow urine one of the earliest supposed remedies for Phylloxera.

- 1397. RILEY, C. V. The bark-louse. <Beach, A. E. The Science Record for 1874, N. Y., 1874, p. 356.

 Discovery of the male of Mytilaspis conchiformis [=pomorum].
- 1398. RILEY, C. V. Codling-moth heresies. <N. Y. Tribune, 2 January, 1875. S.-b. No. 13, p. 163.

 Criticism of paper of S. B. Peck: Carpocapsa pomonella confines its attacks

Criticism of paper of S. B. Peck; Carpocapsa pomonella confines its attacks to one apple and does not enter the ground.

1399. RILEY, C. V. Shall we scrape our trees? <N. Y. Tribune, 6
February, 1875. S.-b. No. 10, p. 10.

Careful scraping of trees early in spring is beneficial as a safeguard against the attacks of many injurious insects.

1400. RILEY, C. V. What are army-worms? <N. Y. semi-weekly Tribune, 6 February, 1875.

Differences between Laphygma frugiperda and Leucania unipuncta.

- 1401. RILEY, C. V. The hickory bark-borer, Scolytus caryæ. <Colman's Rural World, 6 February, 1875. S.-b. No. 17, p. 105.

 Natural history of Scolytus caryæ [= 4-spinosus].
- 1402. RILEY, C. V. Genuine vs. bogus chinch-bugs. <N. Y. Tribune, 10 February, 1875, figs. S.-b. No. 10, p. 10.

Answer to inquiry of C. H. Cushing; food-habits and figures of Nysius destructor [=angustatus] and Blissus lencopterus; larva of Deilephila lineata feeds on purslane.

- 1403. RILEY, C. V. Remedies for Phylloxera. <N. Y. Tribune, 10 February, 1875. S.-b. No. 10, p. 22.
 - At present no insecticide is effective against *Phylloxera vastatrix*; grafting of more susceptible varieties on the roots of the least susceptible advised.
- 1404. RILEY, C. V. Newest facts of grape Phylloxera. <N. Y. Tribune, 10 February, 1875. S.-b. No. 10, pp. 21–22.

Succession of different forms of individuals in the cycle of development of *Phylloxcra* described; seasons at which the different forms appear; places in which eggs are laid by the winged females.

- 1405. RILEY, C. V. Is the Colorado beetle poisonous? <N. Y. Weekly Tribune, 17 February, 1875. S.-b. No. 10, p. 9.
 - Insists upon the poisonous nature of the fumes from scalded or burning masses of *Doryphora decemlineata*, in opposition to statements by Prof. T. J. Burrill.
- 1406. RILEY, C. V. Notes of Phylloxera. <N. Y. Tribune, 4 March, 1875. S.-b. No. 10, pp. 20–21.
 - Critical review of A. S. Fuller's "Distribution of the grape-louse;" maintains the specific identity of the so-called gall-inhabiting and root-inhabiting forms of *Phylloxera*, and the North American origin of this insect.
- 1407. [RILEY, C. V.] The Colorado potato-beetle abroad. <N. Y. Weekly Tribune, 17 March, 1875. S.-b. No. 10, pp. 14–15.
 - Statement and criticism of measures adopted by several European Governments to prevent the introduction of *Doryphora decemlineata* into their countries.

- 1408. RILEY, C. V. Bud-eating insects. <Cultivator and Country Gentleman, 25 March, 1875, v. 40, p. 183.
 - Habits, food-plants, and means against Agrotis scandens; means against field cut-worms.
- 1409. RILEY, C. V. Description of a new species of Agrotis. < Proc. Bost. Soc. Nat. Hist., March, 1875, v. 17, pp. 286–288.

 Description of Agrotis morrisoniana n. sp. [p. 286].
- 1410. RILEY, C. V. On the summer dormancy of the larva of *Phyciodes nycteis*, Doubleday, with remarks on the natural history of the species. <Proc. Amer. Assoc. Adv. Sci. for 1874, 4 June, 1875, v. 23, pp. 108–112. Separate: <Salem, Mass., March, 1875, pp. 108–112. Abstract: <Hartford [Conn.] Daily Courant, 18 August, 1874, No. 11162, v. 38, No. 197, p. 2. See: <N. Y. Tribune Extra: Lecture and Letter series, No. 21, August, 1874, p. 75.
 - Observations on *Phyciodes nycteis* and *Argynnis bellona*; significance of the same; detailed description of the larva and pupa of *P. nycteis*.
- 1411. RILEY, C. V. Descriptions of two new moths. <Trans. Acad. Sci. St. Louis, 1 February [March], 1875, v. 3, pp. 240-242, figs. 12-13.
 - Descriptions and figures of Xanthoptera [= Exyra] ridingsii n. sp. and Cerura multiscripta n. sp.
- 1412. RILEY, C. V. The climate for Doryphora. <N. Y. Tribune, 2 April, 1875. S.-b. No. 10, p. 16.
 - Comments on letter from D. L. Garver; *Doryphora decemlineata* would be likely to thrive as well in the climates of Europe as in North America; its transportation across the ocean is not improbable.
- 1413. RILEY, C. V. [On an Acridium eaten out by ants.]. < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 102 Proc.
 - Verbal communication; inner soft parts of an Acridium americanum eaten out by Myrmica minuta.
- 1414. RILEY, C. V. [On the chrysalis of *Pronuba yuccasella*.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 106 Proc.
 - Verbal communication; adaptation of pupa of Pronuba yuccasella to its needs of prying its ways through the soil.
- 1415. RILEY, C. V. [On regulating sex in insects.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 108 Proc.
 - Verbal communication; critical review of Mrs. Treat's "Controlling sex in butterflies"; female insects need more nourishment than the males, but sex is determined in all animals at conception.
- 1416. RILEY, C.V. [On the peculiarities of Nephila plumipes.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 109 Proc.
 - Verbal communication; comparison of the structure of male and female Nephila plumipes; habits and silk-production of the female.
- 1417. RILEY, C. V. [On the peculiarities of the Mexican honey-ant.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 109 Proc.
 - · Verbal communication; structural and functional division of the neuters of Myrmecocystus mexicanus into two distinct kinds.

- 1418. RILEY, C. V. [On the peculiarities of *Physianthus albens.*] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 109 Proc.
 - Verbal communication; the flowers of *Physianthus albens* so constructed as to hold large *Sphingidæ* fast by the tongue.
- 1419. RILEY, C. V. [On the capture of moths by *Physianthus albens*.] <Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 115 Proc. Reprinted, with changes, from <Moore's Rural New Yorker, 29 August, 1874, v. 30, p. 140.
 - Verbal communication; method of capture of insects, particularly moths, by the flowers of *Physianthus albens*; records the capture of a number of *Noctuidæ* and of *Sphingidæ*, especially *Deilephila lineata*; *Nerium oleander* and *Enothera grandiflora* are said to capture Sphinx-moths in Europe.
- 1420. RILEY, C. V. [On the Yucca borer.] <Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 139 Proc.
 - Verbal communication; entomological interest attaching to the Yucca; abode and synonymy of Megathymus yucca.
- 1421. RILEY, C. V. [New biological facts regarding the grape Phylloxera.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, pp. 147–148 Proc.
 - Sequence of development of certain forms of *Phylloxera*; sexual forms of three species of *Phylloxera* obtained; alleged discovery by J. Lichtenstein of the winged form of *P. vastatrix* on *Quercus coccifera* in Europe, discredited, and this form considered by E. G. Balbiani as a new species, named *P. lichtensteinii*.
- 1422. RILEY, C. V. [On the connection of locust invasions with the occurrence of drought.] <Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 163 Proc.
 - Verbal communication; no connection between the occurrence of drought and of locust invasions.
- 1423. RILEY, C. V. Seventh annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture pursuant to an appropriation for this purpose from the legislature of the State. <10th Ann. Rept. State Board of Agric. for 1874, April, 1875, pp. 7 + 196 + 4, 40 figs. Separate: <Jefferson City, Mo., April, 1875, pp. 7 + 196 + 4, 40 figs.

PREFACE III
TABLE OF CONTENTS VII

1.

NOXIOUS INSECTS.

The Colorado Potato-beetle, Doryphora 10-lineata...

Its gradual spread eastwar 1, 1—It reached the Atlautic during the year 1874, 1—Injuries done during the year 1874, 2—Alarm about it in Europe, 3—Prohibiting the importation of American potatoes by European governments, 3—The insect probably introduced into Europe in the perfect form, 3—It would doubtless thrive in Europe if imported, 4—Its ravages exaggerated and underrated, 5—On the safety and advisability of the use of Paris green, 8—Past experience with the poison, 10—Influence of Paris green on the plant and on the soil, 11—Influence of the green on

THE COLORADO POTATO-BEETLE—Continued.

man indirectly through the soil or through the plant, 13-The beetle eats as well as the larva, 14—It passes the winter as imago, 14-New food-plants, 14-New means of destruction, 15-The Gray sprinkler, 15—The proper scientific name of the beetle, 16— Mr. Carrière's ridiculous statements, 17.

The Chinch-bug, Blissus leucopterus. Its disastrous work in 1874, 19—Circular distributed among farmers, 19—Appearance and transformations of the chinch-bug, 20— The short-winged form, 20-Description of the chinch-bug and its earlier states, 21-Its past history in America 22-Its past history in Missouri, 22-Destructive powers of the chinch-bug, 24—Its injuries in 1874, 24—Its injuries in Missouri in 1874, 25— Food-plants, 26—Time required for the complete development of various insects, 27-Number of annual broods, 27-Its rapid increase, 28-Where the eggs are laid, 28-Flight of the chinchbug, 29—Its migrations on foot, 30—Heavy rains destructive to it, 30—Direct remedies, 31—Irrigation, 31—Preventive measures, 32-Burning, 32-Rolling, 33-Manuring and early sowing, 34-Mixing seed, 34—Preventing the migration of the bugs from one fie'd to another, 35-Importance of winter work and combined action, 36-Other possible remedies, 37-Abstaining from the cultivation of grains, 38-Natural enemies, 38; lady-birds, 39; lace-wing fly and habits of its larva, 40; the insidious flower-bug and the many-banded robber, 41-Birds destructive to the chinchbug, 41—Discussion of other proposed remedies and preventive measures, 41—The chinch-bug injurious to stock, 43—Prognosticating, 44-Unnecessary tears, 44-Bogus chinch-bugs, 45-The false chinch-bug, 46—The insidious flower-bug and the ash-gray leaf-bug, 47—The flea-like negro-bug, 48—Recapitulation, 49.

Appendix to the article on the chinch-bug List of correspondents who replied to the circular, 51—Questions answered by correspondents, 52—Answers given by correspondents, 53.

THE FLAT-HEADED APPLE-TREE BORER, Chrysobothris femorata Extent of its ravages in fruit and shade trees, 71—Its natural history, 72-Natural enemies, 73-Chalcid larva, 74-The cherished Bracon, 75—The useful Labena, 75—Ants, 76—Remedies, 76— Keeping young trees vigorous and healthy, 77—Coating the trunks and larger branches with soap and other greasy substances, 77-Scraping, 78-Wrapping wire gauze around the trunk, 79-Cutting out the newly-hatched larva, 79.

Canker-worms, Paleacrita vernata; Anisopteryx pometaria..... Confusion regarding the two species of canker-worms, 80-Distinguishing characters of the spring canker-worm, 80-Description of egg, larva, and chrysalis of Anisopteryx vernata, 82-How the fall canker-worm differs from the preceding, 83—Description of egg, larva, and chrysalis of Anisopteryx pometaria, 84-Practical importance of distinguishing the two species, 85-Comparative description of the spring and fall canker-worms, 86-Conclusion, 88—Extracts from the original essay on the canker-worm by W. D. Peck, 89.

THE GRAPE PHYLLOXERA, Phylloxera vastatrix..... Completion of its natural history, 90—The true sexual individuals discovered, 91—Epitome of the life-history of the grape Phyllox19

51

71

80

THE GRAPE PHYLLOXERA—Continued.

era, 91—Different forms presented by the species, 93—Its power to change its habit, 93—Specific identity of the gall- and rootlouse, 94-Untrustworthy experiments made by the Department of Agriculture, 95-Proof of the identity of the two forms, 95-The gall-louse is but a transient form, 96—Where do the winged females lay their eggs? 96—The winged female lays the egg wherever she is carried by the wind, 97-Particular part of the vine chosen by the winged female for laying her eggs, 98—The trne sexual individuals, 98—Injury done by Phylloxera in America during the year 1874, 99—Range of the insect in America, 101— Does it occur in South Carolina and Georgia? 102—The Phylloxera in California, 103-Injury done during the year 1874 in France, 103—Its spread in Europe, 104—Direct remedies, 105—Natural enemies, 106—Susceptibility of different varieties of grape-vine, 106—Grafting as a means of counteracting the work of Phylloxera, 108—Underground grafting, 109—Methods of grafting above ground, 112—Roots to use as stock, 115—Varieties to graft, 116— American grape-vines abroad, 116.

THE ROCKY MOUNTAIN LOCUST, Caloptenus spretus..... Its natural history, 121—Method of egg-laying, 121—The egg, 122— The newly-hatched locust and its development, 122—Where the eggs are laid by preference, 123—The invading swarms are formed by a single species, 124—Difference between the Rocky Mountain and the red-legged locusts, 125-Mr. Thomas's description of the red-legged locust, 126—Variations, 126—Measurements of Caloptenus femur-rubrum, 127—Description of Caloptenus spretus and of its larva and pupa, 129—Measurements of Caloptenus spretus, 130— Summing up the difference between the two species, 132—Chronological history, 132-Locust invasions in the Old World, 132-Accounts of earlier locust invasions in America, 133—Chronological history of the Rocky Mountain locust, 134-Earlier invasions, 135—Data regarding the invasion of 1867, 137—The invasion of · 1873, 141—The invasion of 1874, 143—The invasion of 1874 in Missouri, 144—Questions addressed to correspondents in each county in Missouri regarding the locust, 144-Summary of the answers given by correspondents, 145—The invasion of 1874 in Kansas, 148—In Nebraska, 151—In Iowa and Minnesota, 153— In Colorado, 154—In Dakota and Manitoba, 155—Flight and ravages of a locust swarm, 156-Food-plants, 158-Time of appearance of invading swarms, 160—Eastern limits of locust invasions, 161—Native home of the species, 162—Explanation of the migratory instinct, 164—This locust can not thrive in the Mississippi Valley, 164—It is an subalpine insect, 165—What injury may be expected in Missonri in 1875, 166-Ravages of migratory locusts in the Atlantic States, 167—Description of the Atlantic migratory locust, 169—Differences between Caloptenus spretus, differentialis, and atlanis, 170-Injury from other, non-migratory, locusts, 171—The differential and two-striped locusts, 173—Encmics and parasites, 174—Birds destroying locusts or their eggs, 174—The 117

191

- 1423. RILEY, C. V.—Continued.
 - THE ROCKY MOUNTAIN LOCUST—Continued.

silky mite, 175—The locust mite and Dr. Le Baron's description thereof, 176—The anonymous Tachina-fly, 178—The common flesh-fly, 180—Remedics, 181—Classification of protective measures, 181—Natural agencies, 181—Destruction of the eggs, 181—Destruction of the young wingless locusts, 182—Driving off the winged locusts, 184—Further suggestions, 185—Locusts as food for man, 186—The popular and scientific names of the insect, 187—Prairie fires vs. the Rocky Mountain locust, 189.

1424. RILEY, C. V. The ways of bag-worms. <N. Y. Semi-weekly Tribune, 14 April, 1875, 3 figs. S.-b. No. 10, pp. 11, 36.

Figure of the larva-case of an undetermined species of *Psychidw* and of the several stages of *Thyridopteryx ephcmeræformis*; habits of and means against the latter.

- 1425. [RILEY, C. V.] [Poisonous qualities of the Colorado potatobeetle.] <N. Y. Tribune, 14 April, 1875. S.-b. No. 10, p. 17, 34. Mentions instances of poisoning from the fumes given out by scalded or crushed masses of *Doryphora decemlineata*.
- 1426. [RILEY, C. V.] Cure for canker-worm. <N. Y. Tribune, 21 April, 1875. S.-b. No. 10, p. 25.

Paris green recommended for destruction of canker-worms when other remedies are lacking. Paris green not to be used against all insects.

1427. [RILEY, C. V.] Paris green: Its effects on plants and soils, and through them on man. <N. Y. Tribune, 12 May, 1875. S.-b. No. 10, pp. 15–16; 34–35.

Paris green is not injurious to plants or the soil, nor dangerous to man when properly applied.

1428. [RILEY, C. V.] Swellings on roots of Ampelopsis. <Colman's Rural World, 5 June, 1875. S.-b. No. 10, p. 1.

Answer to letter of H. Hilker; swellings on roots of Ampelopsis possibly occasioned by root-lice, but not by Phylloxera.

1429. RILEY, C. V. Apple-tree plant-lice. <Colman's Rural World, 5 June, 1875.

Answer to letter of A. Moyer; occurrence of and means against Aphis mali.

- 1430. RILEY, C. V. [Torrubia elongata, the white-grub fungus.] < Colman's Rural World, 12 June, 1875, v—, p—.
 - Torrubia elongata proposed as the name for the white-grub fungus; figures of the same.
- 1431. RILEY, C. V. The Colorado potato-beetle, Doryphora decemlineata. <Garden [London], 24 July, 1875, v. 8, pp. 71-72, 5 figs. S.-b. No. 23, p. —.
 - Description and figures of all stages of Doryphora decemlineata and D. juncta; migrations, habits, prolificacy, enemies, and parasites of D. decemlineata; means against it; the use of Paris green; probability and probable method of its introduction into Europe and means of preventing such introduction; figures Lydella [= Exorista] doryphora, Harpactor [= Milyas] cinctus, Arma [= Podisus] spinosus, and Hippodamia convergens.

- 1432. RILEY, C. V. Locusts vs. chinch-bugs. <N. Y. Tribune, 4 August, 1875. S.-b. No. 10, p. 40.
 - Locust ravages are likely to prevent serious injury by *Blissus leucopterus*; the latter more noticeable than usual; excessive rains in 1875, also unfavorable to the development of *Blissus leucopterus*.
- 1433. RILEY, C. V. No locust injury in Kansas and Missouri this fall. <N. Y. Tribune, 1 September, 1875. S.-b. No. 10, p. 166.
 - Predicts immunity from attacks of Caloptenus spretus in the fall of 1875 in Kansas and Missouri.
- 1434. RILEY, C. V. Prof. Riley and the locusts. <St. Louis Daily Globe Democrat, 4 September, 1875, v. 1, No. 108, p. 3. S.-b. No. 10, pp. 158-159.
 - Reply to C. A. Davis's "Prof. Riley and the locusts;" food-plants, habits, parasites, and period of development of *Caloptenus spretus*; need of more extended investigation of this insect in its native haunts and breeding places.
- 1435. RILEY, C. V. The grape-leaf gall. < Cultivator and Country Gentleman, 9 September, 1875, v. 40, p. 567.

 Identity of the root and leaf-forms of Phylloxera vastatrix.
- 1436. RILEY, C. V. White-grub fungus. <N. Y. Tribune, 6 October, 1875. S.-b. No. 10, p. 26.
 - Answer to inquiry of A. C. G.; brief history of *Torrubia elongata*, a parasite of *Lachnosterna fusca*.
- 1437. RILEY, C. V. Flying locusts in Illinois. < Cultivator and Country Gentleman, 28 October, 1875, v. 40, p. 679.
 - Caloptenus spretus confounded with C. femur-rubrum by B. F. J.; C. spretus does not occur in Illinois.
- 1438. RILEY, C. V. Remarks on canker-worms and description of a new genus of *Phalænidæ*. <Trans. Acad. Sci. St. Louis, 5 November, 1875, v. 3, pp. 273–280, figs. 14–21. Separate: <St. Louis, Mo., 1875, pp. 8, figs. Reprint: <Sth Ann. Rept. State Ent. Mo., May, 1876, pp. 12–18, figs. 3–10. Notice: <Ca. Ent., November, 1875, v. 7, p. 219. <Proc. Bost. Soc. Nat. Hist., February, 1876, v. 18, p. 201.
 - Treats of Paleacrita [n. g.] vernata and Anisopteryx pometaria; see No. 1482 for synopsis of contents.
- 1439. RILEY, C. V. Notes on the natural history of the grape Phylloxera, *Phylloxera vastatrix*, Planchon. <Trans. Acad. Sci. St. Louis, 5 November, 1875, v. 3, pp. 281–287, fig. 22. Extract: <Amer. Nat., June, 1881, v. 15, pp. 483–484.
 - Discovery of the nidus in which eggs are deposited by the winged female of *Phylloxera vastatrix*; development, habits, description, and figures of the same; description of the male and of the impregnated egg; figure of the male of *P. carywcaulis*; summary of the natural history of *P. vastatrix*.
- 1440. [RILEY, C. V.] Grubs and guess-work. <N. Y. Semi-weekly Tribune, 12 November, 1875. S.-b. No. 10, pp. 17-18.
 - Characterizes some subdivisions of the Lamellicornia by their habits; contrasts the larve, imagos, and habits of Lachnosterna quercina [=fusca] and Lygyrus relictus; means against the former.

- 1441. R[ILEY], C. V. Entomological. Apple-tree borers; timber encourages them; new bag-worm. <Colman's Rural World, 13 November, 1875. S.-b. No. 10, p. 188.
 - Timber-trees near apple orchards increase the likelihood of the attacks of Saperda bivittata [=candida] upon the apple-trees; habits and food-plants of Psyche confederata.
- 1442. RILEY, C. V. The army-worm; an important point yet to ascertain in its history. How it comes and goes; its natural enemies; preventive measures. <N. Y. Tribune, 16 November, 1875, 8 figs. S.-b. No. 10, pp. 29-31; 31-33.
 - Various applications of the name "army-worm;" seasons, habits, and natural enemies of and means against Leucania unipuncta; supposed method of oviposition; figures of it in its several stages, of Exorista flavicauda, Microgaster [= Apanteles] militaris, Ophion purgatum, and Pezomachus minimus, and of cocoons of the last; habits of these parasites.
- 1443. RILEY, C. V. The flying locusts in Illinois. <Cultivator and Country Gentleman, 25 November, 1875, v. 40, p. 744.

 Extent of swarms flying south over Kansas and Nebraska on September (4th?).
- 1444. R[ILEY], C. V. Scabby potatoes. <N. Y. Tribune, 15 December, 1875. S.-b. No. 10, pp. 20, 29.

 The scab in potatoes is caused by imperfectly studied Acarina.
- 1445. [RILEY, C. V.] Not the Hessian-fly. <N. Y. Tribune, 15 December, 1875. S.-b. No. 10, pp. 20, 29.

 Answer to inquiry of J. H. K.; significance of the name and habits of Aphodius inquinatus.
- 1446. R[ILEY], C. V. How to destroy locusts. <Colman's Rural World, 23 December, 1875. S.-b. No. 10, p. 185.

 Answer to letter of C. Herschel; means against Acridide; recommends flooding and ditching.
- 1447. RILEY, C. V. Paris green as an insect destroyer. <N. Y. Tribune, 28 December, 1875. S.-b. No. 10, pp. 13-14; 28-29.

 Paris green as a means against *Doryphora decemlincata*; R. C. Kedzie's investigations show that Paris green is not deleterious to the soil or the crops.
- 1448. RILEY, C. V. Oak apple. <Amer. Cyclopædia, 1875, v. 12, pp. 558-559, 3 figs. S.-b. No. 14, pp. 47-48.

 Description of galls of Cycling terminalis. C. I.-. Amphibaline agreeus, spangification.
 - Description of galls of Cynips terminalis, C. [= Amphibolips] quercus-spongifica, and C. [=A.] q-inanis; manner of their formation; figures the latter two galls and a parent fly; nature of galls and problems involved in their study.
- 1449. [RILEY, C. V.] Phylloxera. <Amer. Cyclopædia, 1875, v. 13, pp. 477-480, figs. 1-8. S.-b. No. 10, pp. 84-87.
 - Habits, food-plants, and characters of the genus *Phylloxera*; natural history of *P. vastatrix*; figures its several forms and the galls formed by it; history of the ravages in the vineyards of France.

- 1450. [RILEY, C. V.] Potato-bug. <Amer. Cyclopædia, 1875, v. 13, pp. 768-771, fig. 1-7. S. b. No. 10, pp. 89-92.
 - Food-plants, original home, geographical distribution, natural history, enemies of, and means against *Doryphora decemlineata*; figures of it and of *Lydella* [= *Exorista*] *doryphora*, *Arma* [= *Podisus*] *spinosus*, *Harpactor* [= *Milyas*] *cinctus*, *Mysia* [= *Anatis*] 15-punctata, and *Doryphora juncta*.
- 1451. RILEY, C. V. Rocky Mountain locust. < Amer. Cyclopædia, 1875, v. 14, pp. 371-374, figs. 1-10.
 - Ravages, description, natural history, migrations, and enemies of and means against Caloptenus spretus; figures of it with details of its structure and habits; figures C. femur-rubrum, Astoma gryllarium [= Trombidium locustarum], Trombidium scriceum, and Sarcophaga carnaria.
- 1452. RILEY, C. V. The hateful or Rocky Mountain locust, Caloptenus spretus. <N. Y. Tribune, 1875. Reprint: <Trans. Kans. State Hortic. Soc. for 1874, 1875, v. 4, pp. 172–176.
 - Answer to inquiry of Z. F. Hopkins; probable limit of the ravages and range of Caloptenus sprctus in Kansas, Missouri, and neighboring States in 1875; means against and vernacular name of this species; vernacular names of Acridida and Locustida in general.
- 1453. RILEY, C. V. Prairie fires and hateful locusts: is there any connection between them? <N. Y. Tribune, 1875. Reprint: <Trans. Kans. State Hortic. Soc. for 1874-775, v. 4, pp. 176-180.
 - Criticism of the effects of the drought, hot winds, locusts, and short crops in Kansas caused by the burning of the prairie grasses as stated in Kansas Farmer, 23 September, 1874; benefits resulting from prairie fires by the destruction of hibernating Blissus leucopterus, Nysius destructor [= angustatus], etc.; relations of prairie fires to the origin and maintenance of prairies.
- 1454. RILEY, C. V. [Address on entomology.] <Trans. Ill. State Hortic. Soc. for 1874, 1875, n. s., v. 8, pp. 103-111. Partial reprint: <Trans. Kans. State Hortic. Soc. for 1874, 1875, v. 4, pp. 103-104.
 - Egeria rubi [= Bembecia marginata] injurious to blackberry and raspberry; the cause and cure of scab in apples; ravages, food-plants, seasons, habits, and means against Chrysobothris femorata; evidence for the identity of the leaf-and root-forms of Phylloxera vastatrix; means against the same.
- 1455. RILEY, C. V. Discussion of the honey-bee. <Trans. Ill. State Hortic. Soc. for 1874, 1875, n. s., v. 8, pp. 131-132.

 Apis mellifica as an enemy to horticulture; importance of insects in the fer-
 - Apis mellifica as an enemy to horticulture; importance of insects in the fertilization of flowers.
- 1456. RILEY, C. V. Notes on locusts. <Trans. Ill. State Hortic. Soc. for 1874, 1875, n. s., v. 8, pp. 136–137.

 Native habitat of Caloptenus spretus.
- 1457. RILEY, C. V. Nonsense about the Phylloxera. <Colman's Rural World, 12 January, 1876. S.-b. No. 10, p. 3. See: <N. Y. Tribune, 1876. S.-b. No. 10, p. 27.
 - There is no such species as the American corn-grape, which is reported capable of resisting the attacks of *Phylloxera vastatrix*.

- 1458. RILEY, C. V. Small borer in apple-twig. <Colman's Rural World, 26 January, 1876. S.-b. No. 10, p. 1.
 - Answer to letter of F. Holsinger; description and habits of Psenocerus supernotatus.
- 1459. RILEY, C. V. Worms on cottonwood. <Colman's Rural World, 26 January, 1876. S.-b. No. 10, p. 1.
 - Answer to letter of J. H. Davidson; occurrence of larvæ of Drasteria erechthea on Populus monilifera and on Trifolium; description of the image.
- 1460. RILEY, C. V. Ailanthus silk-worm in Missouri. <Colman's Rural World, 27 January, 1876. S.-b. No. 10, p. 2.
 - Answer to letter of "Subscriber;" extent and unprofitableness of the culture of Samia [= Atlacus] cynthia; naturalization of the worm in the United States.
- 1461. R[ILEY], C. V. Cause of smut in wheat. <Colman's Rural World, 26 January, 1876. S.-b. No. 10, p. 2.
 - Critical review of a report by Pulaski Grange, Tenn., on the cause of smut in wheat; absurdity of the report; occurrence of Brachytarsus variegatus in the smut, and of larvæ of Cecidomyia, Meromyza, and Chlorops in the lower joints of wheat; smut caused by Ustilago segetum.
- 1462. RILEY, C. V. Colorado potato-beetle's native home. <N. Y. Tribune, 9 February, 1876. S.-b. No. 10, pp. 12-13.

 Geographical distribution of Doryphora decemlineata prior to 1859.
- 1463. RILEY, C. V. An entomological question. < Prairie Farmer, 26 February, 1876, v. 47, p. 68. S.-b. No. 10, pp. 4, 5; No. 42, pp. 76, 77. See: < Prairie Farmer, 4 March, 1876, v. 47, p. 76. S.-b. No. 10, p. 4.
 - Reply to Proximo's "An entomological question;" commendation of legislative efforts to effect the appointment of a national entomological commission; text of the two bills introduced into Congress; criticism of the same.
- 1464. RILEY, C. V. Insect ravages. An interesting letter from Prof. C. V. Riley. How to protect our agricultural interests; legislation, wise and otherwise; the duty of Congress. <St. Louis Daily Globe-Democrat, 4 March, 1876, v. 1, p. 3. S.-b. No. 10, pp. 5-8.
 - Importance and extent of injuries inflicted by insects in North America; notice of existing legislation upon means against injurious insects; statement of legislation needed for protection against locusts; criticism of bills introduced into Congress for the appointment of a national entomological commission.
- 1465. RILEY, C. V. Notes on the Yucca borer, *Megathymus yucca*, Walk. <Trans. Acad. Sci. St. Louis, 10 January-23 March, 1876, v. 3, pp. 323-344, figs. 25-31. Separate: <St. Louis, Mo., January, 1876, pp. 23, figs. 25-31. Reprint: <8th Ann. Rept. State Ent. Mo., May, 1876, pp. 169-182, figs. 40-55.

See No. 1482 for synopsis of contents; see No. 1602,

- 1466. RILEY, C. V. Entomology. An interesting lecture on the insect world. The subject considered both practically and scientifically. <St. Louis Daily Globe-Democrat, 25 March, 1876, v. 1, p. 3. S.-b. No. 10, pp. 161-164. Reprint, with omissions: <Ware's Valley Monthly, August, 1876, v. —, pp. 281-289. S.-b. No. 14, pp. 163-167.
 - Definition of entomology; claims of the science as a liberalizing study; its economic importance; metamorphoses, abundance, and almost omnipresence of insects.
- 1467. [RILEY, C. V.] The insect world. Lecture by Professor Riley at Washington University. A practical subject for fruit-growers. <St. Louis Republican, 26 March, 1876. S.-b. No. 10, pp. 178, 179.
 - Advantages and interest of the study of insects as compared with that of other animals; economic importance of the study; ravages of Glossina morsitans.
- 1468. RILEY, C. V. Legislation in regard to insects injurious to agriculture. <Nation, 30 March, 1876, v. 22, p. 208.

 Amount of insect injuries in the United States; demand for an independent

commission; duties and limits of such a commission.

- RILEY, C. V. Entomology. Another lecture by Professor Riley. How to counteract the ravages of insects; direct remedies; practical hints to farmers, etc. Some of the duties of the State entomologist. <St. Louis Daily Globe-Democrat, 1 April, 1876, v. 1, p. 3. S. b. No. 10, pp. 135–138; 179–184. Reprint, with omissions: <Ware's Valley Monthly, September, 1876, v. —, pp. 369–380. S.-b. No. 14, pp. 168–173.
 - Classification of means against insects; cause of the destructive occurrence of insects; importation of foreign pests; history of the introduction and spread of *Pieris rapæ*, *Phylloxera vastatrix*, and *Doryphora decemlineata*; natural history of these insects; spread of insects by small degrees; advantageous means against certain insects; encouragement of enemies of and parasites on noxious insects; need of distinguishing friends from foes; need of co-operation and of legislation for the destruction of insects; duties of a State entomologist.
- 1470. [RILEY, C. V.] Scarlet mite. < Colman's Rural World, 12 April, 1876, fig. S.-b. No. 10, p. 160.
 - Answer to letter of G. W. Barnes; occurrence, habits, and figure of *Trombidium sericeum*; habits of *T. holosericeum*; use of *T. tinctorium* as a dye.
- 1471. RILEY, C. V. Hibernation of Amphipyra [=Pyrophila] pyramidoides. <Psyche, March [13 April], 1876, v. 1, p. 152.

Extract from 3d Ann. Rept. State Ent. Mo., pp. 72-73, with additional note; this species sometimes hibernates as a pupa, and doubtless frequently as a moth.

1472. RILEY, C. V. Bag-worms and borers. How to protect our shade-trees and insure their growth. How to render shade-trees healthy. Letter from the State entomologist.

St. Louis Re-

- 1472. RILEY, C. V.—Continued.
 - publican, 14 April, 1876, No. 16843, p. 3, 3 figs. S.-b. No. 10, pp. 173–175; 175–178.
 - Causes of the death of shade-trees in the city of St. Louis; description, figures, and natural history of and means against *Thyridopteryx ephemera-formis* and *Chrysobothris femorata*.
- 1473. RILEY, C. V. The locust plague; how to avert it. < Proc. Amer. Assoc. Adv. Sci. for 1875, 1876, v. 24, pp. 215-222. Separate: < Salem, April, 1876, pp. 215-222.
 - Extent of the ravages of Caloptenus spretus in 1873, 1874, and 1875; classification of and special remarks on the several means to be employed against the same; need of more extensive investigation of the insect in its native haunts and breeding places.
- 1474. R[ILEY], C. V. Honey locust weevil. <Colman's Rural World, 26 April, 1876.
 - Answer to inquiry of E. H. B.; larva of Spermophagus robinia has legs and spins a cocoon.
- 1475. RILEY, C. V. Apple and peach borers. < Colman's Rural World, 9 May, 1876. S.-b. No. 16, p. 11.
 - Answer to inquiry of F. H.; habits, natural history, and means against Chrysobothris femorata and Ægeria [= Sannina] exitiosa.
- 1476. RILEY, C. V. Notes on the codling-moth. <Colman's Rural World, 17 May, 1876. S.-b. No. 13, p. 153.

 Natural history of Carpocapsa pomonella.
- 1477. RILEY, C. V. Plums and cotton. <N. Y. Weekly Tribune, 17
 May, 1876. S.-b. No. 10, p. 167.
 - Answer to letter of J. C.; a belt of cotton-batting around the trunk of the tree is no protection against the injury to plums by Conotrachelus nenuphar.
- 1478. RILEY, C. V. Rose-bug remedy. <N. Y. Weekly Tribune, 17 May, 1876, fig. S.-b. No. 10, p. 167.
 - Answer to letter of Mrs. S. P. Smith; description, figure, habits, and preferred food-plants of and means against *Macrodactylus subspinosus*.
- 1479. RILEY, C. V. Smut in wheat. <N. Y. Weekly Tribune, 17 May, 1876. S.-b. No. 10, pp. 167–168. Reprint: <Colman's Rural World, 14 June, 1876. S.-b. No. 10, p. 169.
 - Critical review of two quoted communications by A. S. and by L. Heskett, on the cause of smut in wheat; smut caused by the growth of *Ustilago segetum*; Brachytarsus variegatus breeds in various smuts; description of the beetle.
- 1480. [ŘILEY, C. V.] Potato-beetle; progress. <N. Y. Weekly Tribune, 17 May, 1876. S.-b. No. 10, p. 168.
 - Arrival of Doryphora decemlineata at the Atlantic coast; its ravages there and means against them.
- 1481. RILEY, C. V. Locusts as food for man. <Proc. Amer. Assoc. Adv. Sci. for 1875, 1876, v. 24, pp. 208–214. Separate: <Salem, May, 1876, pp. 208–214.
 - Reference to previous writings on the use of locusts as food for man; historical evidence of the extensive use of locusts as food; methods of preparing locusts for food; species used hitherto; use of Caloptenus spretus; peculiarities of individual taste or national custom.

134	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	
	RILEY, C. V. Eighth annual report on the noxious, benefit and other insects of the State of Missouri, made to the State of Agriculture, pursuant to an appropriation for purpose from the legislature of the State. <11th Ann. If State Board of Agric. for 1875, May, 1876, pp. 185+4, 55 Separate: <jefferson 10-lineata<="" 185+4,="" 1876,="" 55="" city,="" colorado="" contents="" doryphora="" insects.="" may,="" mo.,="" noxious="" of="" potato-beetle,="" pp.="" preface.="" table="" td="" the=""><td>State this Rept. figs.</td></jefferson>	State this Rept. figs.
	Damage during the year, 1—Abundaut in Atlantic States, 1—Swarming on Coney Island, 2—Injuring egg-plant, 2—Its scientific name, 2—Additional enemies, 3—Eaten by the crow, 3—Remedies, 3—Cost of applying Paris green, 3—Preparing the poison, 3—Use of straw as a protection, 4—Machine for sprinkling, 4—Machine for brashing off the insects, 4—Experience with Paris green, 5—Experiments of Profs. R. C. Kedzie and Wm. McMurtrie show that it may be used with safety, 6—Trial of other remedies, 6—The insect's native home, 8—The theory that it came from the Rocky Mountain region essentially correct, 10—Poisonous qualities of the insect discussed, 10.	
	CANKER-WORMS, Palcacrita vernata; Anisopteryx pometaria. Two species long confounded, 12—They differ generically; new	12
	genus (Paleacrita) proposed for one, 13—The two compared in all stages, 13, 17—Characters of the genus Paleacrita, 17—Distinguished as an ingred fall confer worms 17. Practical consideration	
٠	guished as spring and fall canker-worms, 17—Practical considerations from their differences of habit, 18—Stunting the larvæ does not produce male moths, 19—Traps recommended, 20, 21.	1
	Its generic name, 22—The term "army-worm" applied to various insects, 23—Past history of the army-worm, 24—Known since 1854 in Missouri, 27—It followed the 1871 conflagration around Peshtigo, Wis., 28—Its history in 1875; very general all over the country, 28, 29—Its history in Missouri in 1875, 30—Sexual differences, 30—Sexual organs illustrated, 30, 32—Natural history of the species, 32—Illustrated in all states, 32, 33—It occurs in Europe, Asia, New Zealand, and Australia, 34—Description of the egg, 34—Where the eggs are laid, 34—Conclusions drawn from structure, 36, 37—When the eggs are laid, 40—In what state does the insect hibernate?, 43—Habits of the worm, 45—Why it escapes detection when young, 45—Why it travels in armies, 46—Time of its appearance, 46—Are there one or two broods?, 47—The fall army-worm, 48—How distinguished from the real army-worm, 48—Plants preferred by the army-worm, 49—Its sudden appearance and disappressions.	22
	den appearance and disappearance, 50—It swarms during wet pre- ceded by very dry seasons, 51—Its natural enemies, illustrated, 52—Remedies, 54—Philosophy of winter burning, 54, 55—Preven- tion, 55—Summary of the leading facts concerning it, 56.	
	THE ROCKY MOUNTAIN LOCUST, Caloptenus spretus. Previous experience in spring of 1867, 57—Predictions verified, 58— General outlook in spring of 1875, 60—Extent of country ravaged, 60—The outlook in Missouri, 61—Country ravaged often as bare	5 7

as in midwinter, 61-Account by counties, 62-Atchison County,

THE ROCKY MOUNTAIN LOCUST—Continued.

62-Andrew County, 62-Benton County, 63-Barton County, 63-Bates County, 63-Buchanan County, 64-Caldwell County, 64—Cass County, 64—Clay County, 67—Clinton County, 68—Dade County, 68-De Kalb County, 69-Gentry County, 69-Hickory County, 69-Holt County, 69-Henry County, 69-Jackson County, 69-Johnson County, 72-Lafayette County, 73-Nodaway County, 73-Newton County, 73-Pettis County, 73-Platte County, 73-Ray County, 74-St. Clair County, 75-Vernon County, 76—Condition of things in other States, 76—Kansas, 76— Nebraska, 79-Iowa, 81-Minnesota, 81-Colorado, 84-Dakota, 85-Montana, 87-Wyoming, 88-Texas, 88-Indian Territory, 88—Manitoba, 89—Damage done in Missouri, 89—Destitution in Missouri, 91-Address of relief committee from Saint Louis Merchants' Exchange, 93—Cases of starvation, 94—The Governor's proclamation, 95—The locusts not a divine visitation, 97—Natnral history; mode of molting illustrated, 98-Habits of the nnfledged young, 100-Directions in which the young travel, 101-Rate at which they travel, 102-They reached but a few miles east of where they hatched, 102--Not led by "kings" or "queens," 103—The species taken for such, illustrated, 103, 104—The exodus in 1875, 104—Time of leaving of the winged insects, 104—Direction taken by the winged insects, 105-Destination of the departing swarms, 106—Native home of the species, 109—Views previously expressed confirmed, 110—Conditions of migration, 112-Conditions which prevent the permanent settlement of the species in Missonri, 113-Modification of the species by climatic conditions, 114, 155-Definition of the species, 114-How distingnished in all stages from species most nearly allied, 117-Experience in spring of 1875, 118—Contrast in summer and fall, 119— No evil without some compensating good, 120-Injury to fruit and fruit trees, 121-Food-plants, 121-Only one kind of plant not touched under all circumstances, 121—Changes that followed the locusts, 121—The widespread appearance of a new grass, ordinarily unnoticed, 122—Appearance of large worms, 123—The locusts did not return in the fall, 124-Natural enemies, 124-Remedies against the unfledged insects, 125-Artificial means of destroying the eggs, 125-Various means of destroying the unfledged young, 126—They are within man's control, 126—The proper ditch to make, 128-Machines used in Colorado, 129-Best means of protecting fruit trees, 130—How to avert locust injuries, 131—Prevention, 131—Legislation, both national and local, 132—Bills before the Forty-fourth Congress, 133—Need of a national entomological commission, 133-The bounties offered in Minnesota, 138—The requisites of a good bounty law, 138, 139—How a bounty law would work, 140-Suggestions, 140-Lessons of year, 142-Locusts as food for man, 143—They have been used from time immemorial, and are used extensively at the present day, 145-The Rocky Mountain species quite palatable, 146-Mode of preparation, 147—False opinions and predictions, 148—Unnecessary alarm caused by other species, 148-Injuries of native species in 1875, 150—Locust flights in Illinois in 1875, 151—They were composed of local species, 152, 153-Explanation of these flights, 154-Locust prospects in 1876, 155-No danger from them in Missouri, 156,

THE GRAPE PHYLLOXERA, Phylloxera vastatrix..... The injuries not great in Missouri in 1875, 157—Completion of its natural history, 157—Where the winged female lays her eggs, 157, 161—The sexed individuals illustrated, 158—Description of the true female, 159—Description of the impregnated egg, 159, 162—Practical considerations growing out of these latest discoveries, 163—Decortication of the bark to destroy the impregnated egg, 163—The insect may be imported from one country to another on cuttings as well as rooted plants, 163—Best time to attack the root-lice, 163—Phylloxera ravages in California, 163—Great destruction around Sonoma, 164-Need, of action by the State authorities, 164-Occurrence of Phylloxera in the Southern States, 164—Report of committee appointed by the American Pomological Society, 165—Its occurrence in Georgia, 166, 167—American grape-vines in Europe, 167—Large demand for our vines, 167— The American vines flourishing in Southern France where the European varieties perish, 167—The orders for some varieties

exceeded the supply, 168—Probable future demand, 168. INNOXIOUS INSECTS.

The Yucca borer, Megathymus yucca.

The only North American butterfly whose larva has the boring habit, 169—The arbitrary nature of classificatory divisions, 170—Butterflies and moths not easily separated, 170—Biological history of the species, 171—Illustrations of all states, 171, 172—Habits of the larva, 171, 172, 181—Mode of pupation, 172, 180—Flight of the imago, 173, 181—Position of wings when the imago rests or walks, 173—Bibliographical notes, 173—Detailed descriptions of the different states, 174, 175, 181—Structural characters illustrated, 175—Affinities of the species, 176—It is a true butterfly, belonging to the hesperians, 178—Characters of the castnians contrasted with those of hesperians, 176, 177, 178—In classification it is better to widen than restrict in the higher groups, 179—Enemies of the Yucca borer, 179—Concluding remarks, 179—Unsafe to describe species from mere drawings, 179.

NOXIOUS INSECTS—Continued.

Supplementary notes on the army-worm, Leucania unipuncta....
Completion of its natural history, 182—Oviposition of the moth described, 183—Eggs described, 183—Conclusions previously arrived at verified, 184—Description of the different larval stages, 184.

1483. RILEY, C. V. Is the Colorado potato-beetle poisonous? < Colman's Rural World, 7 June, 1876. S.-b. No. 10, pp. 164–165. Extract from the 8th Ann. Rept. State Ent. Mo., pp. 10–12; abstract and critical review of Grote and Kayser's "Are potato-bugs poisonous?"; considers the experiments inconclusive.

1484. [RILEY, C. V.] The eggs of the army-worm. <Colman's Rural World, 7 June, 1876. S.-b. No. 10, p. 170. Reprint: <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 211 Proc. <Amer. Nat., August, 1876, v. 10, pp. 508-509.

Reasons why eggs of *Leucania unipuncta* have not been observed heretofore; verification of author's conclusions as to the probable method of oviposition of this species; description of eggs and young larvæ.

169

157

- 1485. R[ILEY], C. V. Persian insect powder. <N. Y. Tribune, 7 June, 1876. S.-b. No. 10, p. 168.
 - Answer to letter of G. W. Holmes; impossibility of prescribing means against unknown iusects; Paris green unsuitable for protection of bearing fruit trees; pyrethrum recommended.
- 1486. R[ILEY], C. V. Cocoons of silkworms. <N. Y. Tribune, 7 June, 1876. S.-b. No. 10, p. 168.
 - Answer to letter of A. R. Sprout; description of larva and cocoou of *Callosamia* [= Attacus] promethea; colors of imagos; food-plants of larvæ; deposition of eggs; silk of little value.
- 1487. [RILEY, C. V.] Is Paris green absorbed? <N. Y. Tribune, 7 June, 1876. S.-b. No. 10, pp. 168–169.
 - Answer to letter of M. F.; plants colored by certain tinctures placed at their roots; Paris green not absorbed into the tissues of plants in perceptible quantities, but neutralized in the soil.
- 1488. RILEY, C. V. Ditching for young locusts. < Colman's Rural World, 14 June, 1876. S.-b. No. 10, pp. 171-172; 203-204; 204-205.
 - Critical review of J. Stayman's article on same subject; proper width and depth of ditches to check the march of unfledged Caloptenus spretus and other locusts.
- 1489. RILEY, C. V. Periodical Cicada, "17-year locust." <N. Y. Semi-weekly Tribune, 23 June, 1876, 3 figs. S.-b. No. 10, pp. 166–167.
 - Occurrence of Cicada [= Tibicen] septendecim at Lexington, Va., in 1876; list of localities at which these insects will appear this year; chronological history of a brood; figures of larva, pupe, and imago.
- 1490. RILEY, C. V. Specific for Colorado potato-beetle. < Colman's Rural World, 28 June, 1876. S.-b. No. 10, p. 160. Reprint, with omissions: < N. Y. Tribune. S.-b. No. 14, p. 7.
 - Answer to letter of R. Barbour; directions for the use of Paris green as a means against Doryphora decembineata.
- 1491. RILEY, C. V. [Inquiries concerning the Colorado potato-beetle, and Mayor Brown's answer thereto.] <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 170-172 Proc.
 - Verbal communication; critical review of answer of Mayor Brown to inquiries of A. C. Hardy de Boislieu as to the means of preventing the introduction of Doryphora decemtineata into Belgium; quotes the 6th Ann. Rept. State Ent. Mo., p. 16, showing that the fear of such introduction is not unfounded; inaccuracy of Mayor Brown's opinions; manuer in which Doryphora decemlineata will probably reach Europe if at all.
- 1492. RILEY, C. V. [On the ravages of young locusts in western Missouri.] <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 179–180 Proc.
 - Verbal communication; alarming nature of the ravages of unfledged Caloptenus spretus in western counties of Missouri at the present time; probable future departure of the winged locusts: means to be adopted against the locusts now.

- 1493. RILEY, C. V. [Lecture on the Rocky Mountain locust.] < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 180 Proc.
 - Brief abstract of lecture; prediction of the speedily approaching end of the ravages of *Caloptenus spretus* in Missouri for 1875; plentiful crops to be expected subsequently.
- 1494. RILEY, C. V. [Predictions verified.] < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 185 Proc.
 - Verbal communication; accuracy of author's conclusions as to the probable doings of Caloptenus spretus in Missouri later in the season.
- - Verbal communication; remarks on the extraordinary development of Vilfa vaginæflora for a season after the ravages of Caloptenus spretus have occurred an illustration of "the struggle for existence;" the interesting character of such instances of abnormal multiplication of a species; criticism and defense of the expressions "struggle for existence" and "natural selection."
- 1496. RILEY, C. V. Jumping seeds and galls. <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 190–192 Proc. Reprint: <Gardener's Mo. and Hortic., July, 1878, v. 20, pp. 213–214. See: <Amer. Nat., February, 1876, v. 10, p. 125. Pacific Rural Press, 6 April, 1878.
 - Description of the seeds (of an unknown plant) which are inhabited by the larvæ of Carpocapsa saltitans; manner in which these larvæ cause the seeds to roll and jump; habits of larva within the seed; description of the plant bearing the seeds; the seed of Tamariscus moved by the larva of Nanodes tamarisci, which feeds within it; description of the gall of Cynips [= Neuroterus] quercus-saltatorius and of the jumping of these galls.
- 1497. RILEY, C. V. [On the use of Paris green as an insecticide.]

 <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 193. See:

 <Amer. Nat., February, 1876, v. 10, p. 126.
 - Verbal communication; confirmation, by experiments of R. C. Kedzie, of author's conclusions in regard to the safety of Paris green as an insecticide; this substance metamorphosed into a less soluble form in the ground and held in the ground if not used to excess; when applied in small but sufficient quantities not injurious to plants; objections to the use of the "Potato-pest poison" made at the Lodi [N. J.] Chemical Works.
- 1498. RILEY, C. V. New use for the American Agave. <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 195–196 Proc. See: <Amer. Nat., February, 1876, v. 10, p. 126.
 - Verbal communication; use of the dried flower-stalk of Agave americanus for lining insect-boxes.
- 1499. RILEY, C. V. [Food of insectivorous plants.] < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 201–202 Proc.
 - Drosera, Dionæa and other plants digest, absorb, and appropriate nitrogenous matters; glands for the appropriation of animal food not yet found in the Sarracenias.
- 1500. RILEY, C. V. Parasites on bees. <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 212 Proc.
 - Bees in California infested with triungulins of a *Meloë* sp. which sometimes kill the bees; usual habits of such triungulins.

- 1501. RILEY, C. V. The locust pest. <Sci. Amer., 1 July, 1876 [v. 49]. n. s., v. 35, p. 9.
 - Effects of water and cold on the eggs of Caloptenus spretus.
- 1502. RILEY, C. V. Swallows; bed bugs. < Colman's Rural World, 5 July, 1876. S.-b. No. 10, p. 166.
 - Answer to letter of R. R. Pierce; nests of Hirundo fulva generally infested by Acanthia lectularia; those of Hirundo americana rarely so; while this habit might cause annoyance about a dwelling, it is of little consequence when the bird nests under the eaves of a church.
- 1503. [RILEY, C. V.] Berry and cherry twigs. < N. Y. Tribune, 5 July, 1876. S. b. No. 10, p. 169.
 - Answer to letter of W. Keyser; eggs of Ecanthus niveus laid in twigs of Rubus and Prunus; description, habits of, and means against Oberea tripunctata.
- 1504. R[ILEY], C. V. Three worms and their work. <N. Y. Weekly Tribune, 12 July, 1876, 2 figs. S.-b. No. 14, p. 132.
 - Answer to letter of "Subscriber;" means against larve of Agrotidide, of Elateridæ, and earth worms, Lumbricus sp.; eggs of Agrotis ypsilon laid in the spring; breeding-habits and economic importance of Lumbricus; figures of an clatrid larva and imago.
- 1505. [RILEY, C. V.] Mite parasites of the Colorado potato-beetle. <Mirror and Farmer, 15 July, 1876, v. 28, No. 29, p. 2. S.-b.</p> No. 14, p. 221. Reprint: < Gardener's Mo. and Hortic., September, 1876, v. 18, p. 279. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 219 Proc. Note: < Amer. Nat., October, 1876, v. 10, p. 636.
 - Ectoparasitism of a gamasid mite [Uropoda americana] on Doryphora decemlineata; list of vertibrate enemies of D. decemlineata.
 - The reprints note the occurrence of D. decemlineata in New Hampshire and
- its ravages along the Atlantic coast. See No. 1610.

 1506. [RILEY, C. V.] [Wheat insects.] < Blair [Nebr.] Times, 20 July, 1876. S.-b. No. 10, pp. 200-201.
 - Quotes from 1st Ann. Rept. State Ent. Mo., pp. 159-161; wheat injured in Nebraska in 1876 by Cecidomyia destructor and Meromyza americana; ravages of larva, description of imago and means against the latter; occurrence of Disonycha flaviventris in Nebraska.
- 1507. RILEY, C. V. A new enemy of wheat. <N. Y. Tribune, 21 July, 1876. S.·b. No. 14, pp. 129–130; 130–131.
 - Answer to letter of W. Robson; occurrence, habits, and ravages of Leucania albilinca in Maryland, Pennsylvania, and Kansas; history and description of larva, pupa, and imago of the insect; evolution of new habits and forms among insects.
- 1508. [RILEY, C. V.] Chinch-bug; bee-moth. < Colman's Rural World, 26 July, 1876. S.-b. No. 10, pp. 201-205.
 - Answer to letter of G. R. Christian; means against Blissus leucopterus and Galleria cercana; food, seasons, habits, and description of the latter.
- The grape-root borer, Egeria polistiformis. 1509. RILEY, C. V. < Colman's Rural World, 26 July, 1876, 2 figs. S.-b. No. 10, pp. 205-206.
 - Answer to letter of F. J. Kron; experiments on means against Ægeria [= Sciapteron] polistiformis; habits of larva; figures of imago; geographical distribution.

- 1510. RILEY, C. V. Sweet-potato beetles; "beautiful bugs." <N. Y. Weekly Tribune, 26 July, 1876, 3 figs. S.-b. No. 10, p. 213.
 - Answer to letter of W. Snowden; characters and habits of Cassididæ; description and figures of larvæ and imagos of Coptocycla [=Cassida] bivittata and C. [= C.] nigripes; descriptions of C. aurichalcea and C. guttata; these species feed on leaves of Ipomæa batatas and C. aurichalcea, also on leaves of Convolvulus and of Solanum dulcamara; means against them.
- 1511. [RILEY, C. V.] Those centennial insects. <N. Y. Weekly Tribune, 26 July, 1876. S.-b. No. 10, p. 219.
 - Criticism of communication of L. A. M.; list of principal insects preying upon stored corn.
- 1512. R[ILEY], C. V. Wheat midge; "rue-worms." < N. Y. Tribune, 2
 August, 1876. S.-b. No. 10, p. 202.
 - Larvæ of Papilio asterias feeding on Ruta graveolens; description, habits, seasons, ravages of, and means against Cecidomyia [= Diplosis] tritici.
- 1513. RILEY, C. V. Apple and peach borers. <Colman's Rural World, 9 August, 1876. S.-b. No. 14, pp. 3-4; 55-56. Extract: <Cultivator and Country Gentleman, 7 September, 1876, v. 41, p. 566. Answer to inquiries of F. H.; Chrysobothris femorata distinguished from Sa
 - perda bivittata [= candida] and Ægeria [= Sannina] exitiosa; habits and description of the first; habits of the last and means against both.
- 1514. RILEY, C. V. Large saw-fly. <Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 4.
 - Answer to inquiry of J. B. J.; description of larva and image of Cimber laportei [= americana]; its food-plants and transformations.
- 1515. RILEY, C. V. Cottony scale-insect on maples. < Colman's Rural World, 9 August, 1876. S.-b. No. 14, pp. 4-5.
 - Answer to inquiry of C. F. Mills; Lecanium acericola [= Pulvinaria innumerabilis] injurious to Acer dasycarpum at Springfield, Ill.; habits, description of the scale, and manner of oviposition; no great injury ever done by insects of this genus.
- 1516. RILEY, C.V. Hickory vs. locust borer. < Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 5.
 - Answer to inquiry of F. M. D.; description of larva and image and natural history of Arhopalus [= Cyllene] pictus; characters distinguishing this from A. [= C.] robiniæ; food-plants and seasons of the latter.
- 1517. RILEY, C. V. Stag-beetle. < Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 5.
 - Answer to inquiry of S. Lee; occurrence and characters of Lucanus elaphus; food of larva.
- 1518. RILEY, C. V. Eggs of the angular-winged katydid. ' < Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 5.
 - Answer to inquiry of V. Kriegel; descriptions of eggs, imagos, and song of Microcentrum retinerve; habits and ravages and means against the same.
- 1519. RILEY, C. V. Experience with the Colorado potato-beetle. < Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 6.
 - Letter from N. Coleman; larvæ of Doryphora decemlineata in confinement pupated on the surface of the ground; images cat the tubers underground.

- 1520. [RILEY, C. V.] An elm enemy. <N. Y. Semi-weekly Tribune, 11

 August, 1876. S.-b. No. 14, pp. 54-55.
 - Habits and ravages of and means against Galeruca calmariensis [=xanthome-lena].
- 1521. RILEY, C. V. The locust in 1876. <N. Y. Weekly Tribune, 16 August, 1876. S.-b. No. 14, p. 73.
 - The occurrence and ravages of Caloptenus spretus in Minnesota and Colorado in 1876; their absence from other portions of the country occasionally inhabited by them; their destruction by Astoma gryllaria [= Trombidium locustarum] and inability to maintain themselves in parts of the region they infest.
- 1522. [RILEY, C. V.] The war on "corn-worms." <N. Y. Semi-weekly Tribune, 18 August, 1876. S.-b. No. 14, p. 73.
 - Review of article by A. Brewster; ravages of and means against larvæ of Elateridæ, Agrotididæ and Lachuosterna quereina [=fusca] and the "grubworm," injurious to growing maize.
- 1523. RILEY, C. V. Spined soldier-bug. <Ohio Farmer, 19 August, 1876, v. 50, p. 118.
- 1524. RILEY, C. V. "Potato-pest poison." Sci. Amer., 19 August, 1876 [v. 49], n. s., v. 35, p. 116.
 Condemning some of the patent poisons for Doryphora decemlineata.
- 1525. RILEY, C. F. Cottonwood borers. <N. Y. Weekly Tribune, 23

 August, 1876. S.-b. No. 14, p. 55.

 Answer to inquiry of J. R.; food-plants and ravages of and means against

Plectrodera scalator.

- 1526. RILEY, C. V. Harmless insects. <N. Y. Weekly Tribune, 23 August, 1876. S.-b. No. 14, p. 55.
 - Answer to letter of a "Subscriber;" occurrence of *Psocus venosus* on appletrces; habits and molting.
- 1527. RILEY, C. V. Locust prospects. <N. Y. Tribune, 6 September, 1876. S.-b. No. 10, pp. 213-215; No. 14, pp. 74-75. Reprint: <Prairie Farmer, 16 September, 1876, v. 47, p. 298. S.-b. No. 10, pp. 207-209.
 - Answer to letter of G. H. H.; movements and ravages of Caloptenus spretus in Aug., 1876, in Dakota, Minnesota, Iowa, Nebraska, and Colorado.
- 1528. RILEY, C. V. Some notes on potato-beetles. <Sci. Amer., 9
 September, 1876 [v. 49], n. s., v. 35, p. 164. Reprint: <Ohio
 Farmer, 16 September, 1876, v. 50, p. 179. S.-b. No. 14, pp.
 6, 7.
 - Critical review of S. R. M.'s "Facts about potato-beetles;" Doryphora decemlineata flies in the day-time, but not at night; feeds upon several species of plants; hibernates as an imago, which should be destroyed in early spring.
- 1529. RILEY, C. V. Entomological works wanted. < Colman's Rural World, 20 September, 1876. S.-b. No. 14, p. 2.
 - Answer to letter of J. W. Newman; notice of some works on entomology for the use of students.

- 1530. [RILEY, C. V.] Grape-leaf Philloxera enemy. < Colman's Rural World, 20 September, 1876. S.-b. No. 14, p. 3.
 - Answer to letter of A. Engleman; Leucopis sp.? parasitic in galls of Phylloxera vastatrix and P. rileyi.
- 1531. [RILEY, C. V.] Grape-leaf gall. < Colman's Rural World, 20 September, 1876. S.-b. No. 14, p. 3.
 - Answer to letter of A. A. Briggs; *Phylloxera vastatrix* does comparatively little damage on the leaves of grape-vines; Clinton and Taylor vines peculiarly subject to the attacks of the leaf-inhabiting form; Concord vines seldom attacked by it and not seriously injured by the root-inhabiting form; means against the leaf-inhabiting form.
- 1532. RILEY, C. V. New locust theory wanted. <Colman's Rural World, 27 September, 1876. S.-b. No. 14, p. 62.
 - Answer to letter of W. T. D.; facts not theories wanted in regard to the flights of Caloptenus spretus.
- 1533. RILEY, C. V. Cecropia worm on elder. <Colman's Rural World, 27 September, 1876, 1 fig. S.-b. No. 14, p. 63.
 - Answer to letter of A. S. Van Winkle; description of larva, cocoon, and imago of Samia [= Attacus] cecropia; figure of larva; habits and seasons; availability of the cocoon for silk.
- 1534. RILEY, C. V. The harlequin cabbage-bug. <Colman's Rural World, 4 October, 1876. S. b. No. 14, p. 56.
 - Answer to letter of R. J. Waters; habits, ravages, distribution, and means against Strachia [= Murgantia] histrionica; colors of early stages and of images.
- 1535. [RILEY, C. V.] Butterfly chrysalis. <N. Y. Tribune, 13 October, 1876. S.-b. No. 19, p. 224. Reprint: <N. Y. Tribune, 21 October, 1876. S.-b. No. 10, p. 223.
 - Answer to inquiry of E. B. S.; occurrence of *Danais archippus* in swarms in central United States in autumn of 1876; colors of larva and pupa; larva feeds on *Asclepias*.
- 1536. [RILEY, C. V.] Domesticated katydid. <N. Y. Tribune, 18 October, 1876. S.-b. No. 19, p. 214. Reprint: <N. Y. Tribune, 21 October, 1876. S.-b. No. 10, p. 219.
 - Answer to letter of C. A. P.; longevity and food of Microcentrum retinerve.
- 1537. [RILEY, C. V.] Unjust accusation? <N. Y. Tribune, 21 October, 1876. S.-b. No. 10, p. 209; No. 19, p. 242.
 - Answer to inquiry of a "Correspondent;" Harpalus erraticus accused of destroying shrubbery; predaceous habits of Carabida.
- 1538. RILEY, C. V. The Rocky Mountain locust. < Colman's Rural World, 1876, v. —, 30 October, 6 November, 13 November. S.-b. No. 10, pp. 185–188. See: < Kansas Farmer, November, 1876. S.-b. No. 10, pp. 225–235. < N. Y. Tribune, October, 1876. S.-b. No. 10, p. 215.
 - Verification of predictions in regard to the limitation of the ravages of Caloptonus spretus in western Missouri in 1876; gathering eggs, ditching, rolling, and burning recommended as means against them; recommends the employment of soldiers, of hogs, and of poultry, late planting, and the destruction of the locusts in their native country; Blissus leucopterus more

- 1538. RILEY, C. V.—Continued.
 - injurious in Western Missouri in 1876 than Caloptenus spretus; prospect of freedom from injury by the former for the next two years; Vilfa vaginæflora introduced into the eastern prairies by the locusts; limit to the eastern range of the locusts; they are not led by kings or queens; occurrence of larvæ of Deilephila lineata after the disappearance of the locusts; locust flights in Illinois.
- 1539. [RILEY, C. V.] Canker-worms at the West. <N. Y. Tribune, 31 October, 1876. S.-b. No. 17, p. 59.
 - Occurrence of *Paleacrita vernata* in Michigan in 1872 and in Ohio in 1874, 1875, and 1876; brief account of *P. vernata* and *Anisopteryx pometaria*; their habits, ravages, and means against them.
- 1540. RILEY, C. V. [The venation of Anisopteryx and variation in imagos.] <Ca. Ent., September [October], 1876, v. 8, pp. 178–179.
 - Variation in the venation of the wings of Anisopteryx pometaria and in the imagos of so-called species.
- 1541. RILEY, C. V. A new enemy of the grasshopper. <Lawrence [Kans.] Journal. Reprint: <Industrialist [Manhattan, Kans.], 2 November, 1876, v. 2, No. 30, p. 2. S.-b. No. 10, pp. 209-210.
 - Letter of F. H. Snow with comments; eggs of Caloptenus spretus destroyed by larvæ of Anthomyia calopteni [=-augustifrons]; characters of the larvæ; eggs destroyed by larvæ of some Ichneumou? [=Systæchus oreas]; need of destroying eggs by artificial means.
- 1542. RILEY, C. V. Silk culture in Kansas. Nationalist, 10 November, 1876. S.-b. No. 14, p. 1.
 - Importance of silk culture; practicability of the culture and prospects of its increase in the United States; measures for its promotion in Kansas; successful raising of silkworms on osage orange [Maclura aurautiaca].
- 1543. RILEY, C. V. Bee killers: Asilus flies. <N. Y. Weekly Sun, 15 November, 1876. S.-b. No. 10, p. 201.
 - Habits of Asilidae, especially of Trupanea [= Promachus] apirora and Asilus missouriensis [= Proctacanthus milberti]; description of T. [= P.] apivora; larva of Asilus sericeus feeds on roots of Rheum rhapouticum.
- 1544. RILEY, C. V. How to use Paris green for the cotton-worm. <Colman's Rural World, 15 November, 1876. S.-b. No. 10, pp. 210–211. See: <N. Y. Tribune, 15 December, 1876. S.-b. No. 14, p. 7.
 - Answer to letter of C. W. Niver; directions for the use of Paris green by the dry and wet method; description of sprinkling machines.
- 1545. [RILEY, C. V.] Notodonta concinna. <Colman's Rural World, 15 November, 1876. S.-b. No. 10, p. 219.
 - Answer to letter of J. Barritt; description of larva of Notodonta [= Œdema-sia] eoneinna; habits, food-plants, and means against the same.
- 1546. [RILEY, C. V.] The dog-day harvest fly. <Colman's Rural World, 15 November, 1876. S.-b. No. 10, p. 220.
 - Answer to inquiry of C. A. U.; description of Cicada pruinosa [= tibicen]; method of sonifaction; habits; notice of some manuals of entomology.

- 1547. [RILEY, C. V.] Snake-worms. < Colman's Rural World, 15 November, 1876. S.-b. No. p. 10, p. 220.
 - Answer to letter of J. Armstrong; description of larvæ of Sciara sp.; their habit of moving in congregations; their abode and enemies.
- 1548. RILEY, C. V. Locust eggs. < Colman's Rural World, 15 November, 1876. S.-b. No. 10, p. 223.
 - Answer to letter of C. T.; female Acridida lay more than one litter of eggs; the occurrence of egg-like parasites in males has led to the supposition that the males bore eggs.
- 1549. RILEY, C. V. Locust flights east of the Mississippi. <Colman's Rural World, 22 November, 1876. S.-b. No. 10, pp. 211–212; 221–222; 223–225. Reprint: <Sci. Amer., 16 December, 1876 [v. 49], n. s., v. 35, p. 392, 2 figs. S.-b. No. 10, pp. 216, 217. <Trans. Kans. Acad. Sci., 1877, v. 5, pp. 62–64.
 - Limitation of plants and animals to certain geographical regions; regions in which alone Caloptenus spretus survives; species confounded with C. spretus; occurrence of swarms of Acridium americanum in Ohio; description, geographical distribution, and ravages of the same; swarms of Caloptenus differentialis, C. atlanis, and C. femur-rubrum in Illinois; ravages of the same. The reprint in the Sci. Amer. contains figures of Caloptenus spretus and Acridium americanum.
- 1550. RILEY, C. V. Locusts again. <N. Y. Tribune, 22 November, 1876. S.-b. No. 10, p. 222.

 Effects of winter on the vitality of the cggs of Acridida.
- 1551. RILEY, C. V. The army-worm; its natural history complete. <Sci. Amer., 9 December, 1876 [v. 49], n. s., v. 35, p. 372, 4 figs. S.-b. No. 10, pp. 217-219. Reprint, with slight changes: <Proc. Amer. Assoc. Adv. Sci. for 1876, 1877, v. 25, pp. 279-
 - Geographical distribution of *Leucania unipuncta*; figures of all stages and of ovipositor of female; place and manner of oviposition; description of eggs and young larvæ; number of annual broods; summary of the natural history of this species.
- 1552. RILEY, C. V. The apple-bark louse. <Ca. Farmer, 15 December, 1876. S.-b. No. 14, p. 50.
 - Seasons, habits, and ravages of Mytilaspis pomicorticis [=pomorum]; description of young larvæ; of males and females and formation of scales.
- 1553. RILEY, C. V. The apple maggot; a formidable enemy. <N. Y. Semi-weekly Tribune, 15 December, 1876. S.-b. No. 14, pp. 7–8.
 - Answer to letter of P. M. Augur; description of larva and image of *Trypeta pomonella*; ravages, food-plants, habits, and means against the same; literature of the subject.
- 1554. RILEY, C. V. Entomological notes; confounding friend with foe. <Colman's Rural World, 20 December, 1876. S.-b. No. 14, p. 4.
 - Description of scales and eggs of Diaspis harrisii [= Chionaspis furfurus] and of colors of larva, pupa, and imago of Chilocorus bivulnerus; ravages of the Diaspis; habits and usefulness of the Chilocorus.

- 1555. RILEY, C. V. Locust injury next spring. The territory in Missouri that will probably suffer therefrom. <Colman's Rural World, 20 December, 1876. S.-b. No. 14, pp. 5-6. Reprint: <Industrialist [Manhattan, Kans.], 17 February, 1877, v. 2, No. 44, p. 4. S.-b. No. 14, p. 49.
 - Prediction of ravages of Caloptenus spretus to occur in spring of 1877 in Missouri.
- 1556. [RILEY, C. V.] Amputating insects. <N. Y. Semi-weekly Tribune, 29 December, 1876, 2 figs. S.-b. No. 14, pp. 8-9.
 - Description of the work of Elaphidion putator [= villosum] and Oncideres cingulata; habits of both; figures of larva, pupa, and image of the latter; similarity in habits of Oncideres amputator and E. putator to those of O. cingulata.
- 1557. [RILEY, C. V., et al.] The Rocky Mountain locust or grasshopper, being the report of proceedings of a conference of the Governors of several Western States and Territories, together with several other gentlemen, held at Omaha, Nebr., on the 25th and 26th days of October, 1876, to consider the locust problem; also a summary of the best means now known for counteracting the evil. <St. Louis, 1876, 8°, pp. 3+58, 8 figs. See: <N. Y. Tribune, 1876. S.-b. No. 10, p. 223.
 - Preface, proceedings, pp. 1-36. Practical considerations and suggestions for the suppression of Caloptenus spretus; description, oviposition, transformations, and habits of the same; its enemies and parasites; means against it; description of Anthomyia calopteni n. sp. [= angustifrons]. Figures. Prepared by J. S. Pillsbury, P. Pusey, and C. V. Riley.
- 1558. RILEY, C. V. Potato pests. Being an illustrated account of the Colorado potato-beetle and the other insect foes of the potato in North America, with suggestions for their repression and methods for their destruction. <New York: Orange Judd Company [1876], pp. 108, 49 figs., map. Review: <Cultivator and Country Gentl., 11 January, 1877, v. 42, p. 25. Reply to review: <1bid., 1 February, 1877, v. 42, p. 78. S.-b. No. 14, pp. 56-57.

TABLE OF CONTENTS.

Preface	7
Introduction	9
THE COLORADO POTATO-BEETLE, Doryphora 10-lineata	11
Its past history	11
Prediction that it would reach the Atlantic, 12—Its march across	
the country, 13—It reaches the Atlantic, 14—Its swarming in	
large cities, 16—Its occurrence out at sea, 17.	
The insect's native home	17
When it first attacked the potato, 18.	
Rate at which it traveled	21
How it traveled.	21
Mostly in the beetle state, 21—Assisted by man, 21—Tendency to	
migrate in swarms, 23.	
It spreads, but does not travel in the sense of leaving one district for	
another	23

1558. RILEY,	C.	V.—	Coi	atin	ued.
--------------	----	-----	-----	------	------

THE COLORADO POTATO-BEETLE—Continued.	
Area invaded by it	24
Causes which limit its spread	25
Intense heat in the South, 25—Excessive dryness in the mountains, 26.	
How it affected the price of potatoes	26
The modification it has undergone	27
Its natural history	27
First made known in 1863, 28—The female capable of laying 1,000 eggs, 29—Three broods a year, 28.	,
Its poisonous qualities	29
Exhalations from the crushed bodies injurious, 29.	
Its food-plants	30
The number increases with each year, 32—Varieties of potatoes preferred, 33.	
The beetle eats as well as the larva	33
Its natural enemies	34
Birds which feed upon it, 35–36—Domestic fowls, 36—Reptiles, 36—Spiders and ites, 36–38—True insects, 39—Rust-red social wasp, 40—Lady-birds, 40–43—Ground-beetles, 44–45—Rove-beetles, 46—Blister-beetles, 46—Soldier-bugs, 47–51—Tachina-fly, 52—Asilus-flies, 53.	01
Remedies	54
Encouragement of natural enemies, 54—Preventive measures, 54—	
Mechanical means of destruction, 55—Pincers for, 56—Sun-scalding, 56—Horse-machine, 57—Machines for collecting, 58, 59—Poisonous applications to the plant, 60—Paris green, 61—Different modes of using Paris green, 62–65—Other poisonous applications tested, 66—Patent poisons, 68.	
The use of Paris green	69
Its influence on the plant, 70—Its influence on the soil, 71—Its influence on man indirectly through the soil or through the plant, 74.	
Bogus experiments.	75
Alarm about the insect abroad	76
Unnecessary prohibition of traffic in American potatoes, 77—How the insect will most likely get to Europe, 78—The chances of its getting there, 79–82—Could it become acclimated there?, 82.	
Nomenclature	83
The bogus Colorado potato-beetle, Doryphora juncta	85
OTHER INSECT FOES OF THE POTATO.	
THE STALK-BORER, Gortyna nitela	90
THE POTATO STALK-WEEVIL, Trichobaris trinotata Habits, 92—Remedy, 93.	92
THE POTATO OR TOMATO WORM, Protoparce celeus———————————————————————————————————	93
BLISTER-BEETLES	96
The striped blister-beetle, Epicanta vittata	97
The ash-gray blister-beetle, Macrobasis unicolor	98
The black-rat blister-beetle, Macrobasis unicolor	99

1558.	RILEY, C. V.—Continued.	
	BLISTER-BEETLES—Continued.	
	The black blister-beetle, Epicauta pennsylvanica	99
	The margined blister-beetle, Epicauta cinerea	99
	Remedies, 100.	
	THE THREE-LINED POTATO-BEETLE, Lema trilineata	100
	Habits, 101, 102—Remedies, 102.	
•	THE CUCUMBER FLEA-BEETLE, Crepidodera cucumeris	102
	Habits, 103—Remedies, 103.	
	THE CLUBBED TORTOISE-BEETLE, Coptocycla elavata	103
	Habits, 103—Remedies, 104.	

- pædia, 1876, v. 2, pp. 412-416, 16 figs. S.-b. No. 10, pp. 76-80. Definition of the term "gall-insects"; classification and habits of the same; mention by name and figures of typical species and galls made by them; dimorphism and metagenesis of Cynips.
- 1560. RILEY, C. V. Gall-nuts. < Johnson's New Universal Cyclopædia, 1876, v. 2, p. 417, 1 fig. S.-b. No. 10, p. 81.

 Definition of "gall-nuts" formed by Cynips gallæ-tinctoriæ on twigs of Quer-
- cus infectoria; local origin, ehemical composition, use in the arts.

 1561. RILEY, C. V. Galls. < Johnson's New Universal Cyclopædia,
 - 1876, v. 2, pp. 418-419, 2 figs. S.-b. No. 10, pp. 82-83. Definition of "galls;" their variety of form, texture, and location; their nature and source.
- January, 1877. S.-b. No. 14, pp. 57-58; 58-59.

 Explains means of prognosticating the advent of Calontenus spretus in the
 - Explains means of prognosticating the advent of Caloptenus spretus in the spring of 1877; recommends measures of precaution and the procurement of information concerning means against these insects, and gives assurance that the invasion of the locusts will be but temporary.
- 1563. RILEY, C. V. Bots. <Sci. Amer., 6 January, 1877 [v. 50], n. s., v. 36, pp. 9-10. Reprint: —S.-b. No. 14, pp. 235-236. <Lancaster Farmer, 15 September, 1877, v. 9, p. 142. S.-b. No. 14, p. 129. <Colman's Rural World. S.-b. No. 14, p. 100.
 - Habits, ravages, and means against Gastrophilus equi, Cephalomyia [= Estrus] ovis, and Hypoderma bovis.
- 1564. [RILEY, C. V.] Academy of Science. Brilliant and profound address of Prof. C. V. Riley. The splendid record of the Academy for the past year. Reports of officers and committees and election of managers for the ensuing year. <St. Louis Times, 16 January, 1877, v. —. S.-b. 14, pp. 91-96. Reprint: <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 238-254 Proc. Separate: <St. Louis, 1877, 16 pp. Translation: <Anzeiger des Westerns, 16 January, 1877, v. 43, p. 3. S.-b. No. 14, pp. 50-54.
 - Review of progress made in science in North America, and especially by the Academy of Science of St. Louis in 1876; remarks on entomological observations and publications, and especially on *Doryphora decemlineata*, *Phylloxera vastatrix*, and *Caloptenus spretus*.

- 1565. RILEY, C. V. Is this a grasshopper year? Prof. Riley's opinion concerning the prospect for bugs. It all depends on the kind of weather we have during February. <St. Louis Daily Globe Democrat, 7 February, 1877, v. 2, No. 263, p. 3. S.-b. No. 14, pp. 69-70. Reprint: <Industrialist [Manhattan, Kans.] 17 February, 1877, v. 2, pp. 1, 4. S.-b. No. 14, p. 49. See: <Colman's Rural World, 1877. S.-b. No. 1, pp. 59-60.
 - Replies to questions as to the likelihood of the hatching of eggs of Caloptenus spretus in the spring of 1877; the degree of development attained before winter; possibility of the resumption of development after it has once been arrested.
- 1566. [RILEY, C. V.] Are the locusts hatching? Mistaken identity. <Colman's Rural World, 14 February, 1877, 2 figs. S.-b. No. 14, p. 62.
 - Modified extract from 8th Ann. Rept. State Ent. Mo., May, 1876, pp. 149-150, Tragocephala [=Chortophaga] viridifasciata and Tettix granulatus mistaken for Caloptenus spretus; geographical distribution of the first; hibernation and colors of the two former; figures of both.
- 1567. RILEY, C. V. Condition of locust eggs: Inquiries answered. <Colman's Rural World, 21 February, 1877. S.-b. No. 14, pp. 67-68; 68.
 - Replies to inquiries as to the degrees of development attained by eggs of Caloptenus spretus submitted for examination.
- 1568. RILEY, C. V. Tarred paper for fruit trees. <Colman's Rural World, 7 March, 1877. S.-b. No. 14, p. 61.
 - Critical review of article by E. Gaylord; the inclosure of trunks of fruit trees in tarred paper serviceable as a protection from the sun, rabbits, mice, and borers.
- 1569. RILEY, C. V. Insect on the grape. <Gardener's Mo. and Hortic., March, 1877, v. 19, p. 90.
 - Varieties of grape attacked by Desmia maculalis; means against the same.
- 1570. RILEY, C. V. Ninth annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <12th Ann. Rept. State Board of Agric. for 1876, March, 1877, pp. 7+129+3, 33 figs. Separate: <Jefferson City, Mo., March, 1877, pp. 7+129+3, 33 figs.

There are several species, having different habits, 1—Three which may be destroyed by similar methods, 1—Botanical details as to the current and gooseberry, 2.

3

The Gooseberry Span-worm, Eufitchia ribearia

Its natural history, 3—Most destructive gooseberry insect in Missouri, 3—Generic nomenclature, 3—Characters of the moth, 4—
Description of the egg, 4—Where the eggs are laid, 4—The in-

7

23

27

29

32

34

1570. RILEY, C. V.—Continued.

THE GOOSEBERRY SPAN-WORM—Continued.

sect single-brooded, 5—How it spreads, 5—A native species, 5—Its past history, 5—It prefers the gooseberry to the currant, 6—The moth is closely imitated by one which greatly differs structurally, 6—Parasites, 6—Remedies, 7—Other currant spanworms, 7.

Belongs to the "false-caterpillars," 7—The different specific names it has received, 7—An imported species, 8—Its introduction and spread, 8—Independently imported at several eastern points, 9—Its natural history, 9—How the eggs are laid, 10—Nature and habits of the worm, 11—Characters of the parent flies, 12—Preventive measures, 13—Remedies, 13—White hellebore the best, 13—How best used, 14-15—The worm is not poisonous, 16—Natural enemies, 17—It furnishes a forcible example of arrenotoky, or the power of producing male offspring without impregnation, 18—Results of parthenogenesis in different insects, 18—It also furnishes an interesting instance of defunctionation of special parts, 19—The saw of the female imperfect compared with others, 20—Evolutionary bearings of this fact, 21—Descriptive, 21—Variation in the antennæ and wing veins, 22.

THE NATIVE CURRANT-WORM, Pristiphora grossulariæ

Wherein it differs from the imported species, 23—Its habits, 24—
Where the eggs are laid, 25—How the winter is passed, 25—Its occurrence in Missouri, 26—Remedies, 26—Descriptive, 26.

THE STRAWBERRY-WORM, Harpiphorus maculatus.

Also a false-caterpillar, 27—It has a wide range, 27—How the eggs are deposited, 28—Character and habits of the worm, 28—Remedies, 28—Descriptive, 28.

Abbot's white pine-worm, Lophyrus abbotii.

Destructive power of the insects of its genus in Germany, 29—
Evergreens which it prefers, 30—Habits and characters of the worm, 30—Variation in the antenne, 30—Characters of the perfect flies, 30–31—How the eggs are laid, 31—Natural enemies, 31—Remedies, 32—Descriptive, 32.

LE CONTE'S PINE-WORM, Lophyrus lecontei.

A more general feeder than Abbot's species, 32—The close resemblance of the worms, 33—How they differ, 33—Descriptive, 33—Other species of the genus, 34.

Injury in the West in 1876, 34—Spread of the insect during the year, 34–35—Its great abundance on the Atlantic coast, 35—Rate at which it traveled since 1859, 37—An average of 88 miles a year, 37—How it traveled, 37; principally in the beetle state, and greatly assisted by man, 37—Its migrating habit, 38—Area invaded by it nearly 1,500,000 square miles, 38—Causes which limit its spread, 38—Will it reach the Pacific slope?, 39—How it affected the price of potatoes, 39—The modification it has undergone, 40—A mite parasite added to its natural enemies, 41—Its introduction to Europe, 42—A living specimen found last summer in the Bremen dock-yards, 42—Could it live and multiply in Europe?, 43—Action taken by European governments to prevent its introduction, 44—Consideration of the Kearney "potato-pest poison," 45.

1570. RILEY, C.	V.—Continued.
-----------------	---------------

THE ARMY-WORM, Leucania unipuncta

Further notes and experiments thereon, 47—Two generations produced annually at St. Louis, and a probable third generation, exceptionally, 48—Summary of its natural history, 49.

47

50

57

THE WHEAT-HEAD ARMY-WORM, Leucania albilinea....

A new enemy to wheat, 50—First complaint of it in the East, 51—First appearance in Kansas, 51—Habits and natural history, 52—The egg differs from that of the army-worm, 53—Wherein the worm is distinguished from its destructive congener, 54—Two broods each year, 54—Natural enemies, 54—Remedies, 55—Descriptive, 55.

THE ROCKY MOUNTAIN LOCUST, Caloptenus spretus.

It continues to interest the people of the West, 57-Previous opinions justified, 57—The invasion of 1876, 59—Few in British America, 59-Condition of things in Montana, 59-In Wyoming, 59in Dakota, 59-In Minnesota, 60; locusts and alkali soil, 61; good done by Governor Pillsbury, of Minnesota, 61—In Colorado, 62—In Iowa, 63—In Nebraska, 64—In Kansas, 65—In Missouri, 66—Flights in opposite directions at the same time, 66—Counties in Missouri that were overrun, 67-Red-legged locust troublesome in east Missouri, 68—Detailed reports from counties in Missouri, 68—Andrew County, 68—Atchison County, 68—Barry County, 68-Barton County, 69-Bates County, 69-Benton County, 69—Buchanan County, 69—Cass County, 69—Cedar County, 70—Caldwell County, 70—Clay County, 70—Dade County, 70-De Kalb County, 70-Gentry County, 70-Greene County, 71—Harrison County, 71—Henry County, 71—Hickory County, 71-Holt County, 71-Jasper County, 72-Jackson County, 73-Johnson County, 73-Lafayette County, 73-Lawrence County, 73—McDonald County, 74—Newton County, 74— Nodaway County, 74-Pettis County, 74-Platte County, 74-Polk County, 75—Ray County, 75—Saint Clair County, 75—Vernon County, 75-In Indian Territory, 76-In Texas, 76-In Arkansas, 76—Destination of the departing swarms of 1875, 77—They reached into British America, 78—Source of the swarms of 1876, 79—Eastern line reached, 80—Rate at which the insects spread, 80—Direction of flight, 81--Influence of wind in determining the course of locust swarms, 81-Locust flights east of the Mississippi, 81—Geographical range of species, 82—Causes which limit the spread of the Rocky Mountain locust, 83-Flights of Acridium americanum, 84-Does the female of the Rocky Mountain locust lay more than one egg-mass?, 85-How the eggs are laid, 86—Philosophy of the egg-mass, 87—How the young locust escapes from the egg, 88-How it escapes from the ground, 90-Additional natural enemies, 91—Animals which destroy the eggs, 91—The Anthomyia egg-parasite, 92—The common flesh-fly, 95— Other undetermined enemies of the eggs, 96—Insects which destroy the active locusts, 98-Experiments with the eggs and conclusions therefrom, 99—Experiments to test the effects of alternately freezing and thawing, 99—Experiments to test the influence of moisture upon the eggs, 104-Experiments to test the effects of burying at different depths and of pressing the soil, 104—Experiments to test the effects of exposure to the free air, 104- The Omaha conference, 106-Remedies and suggestions, 108—Des ruction of the young or unfledged locusts, 108—Protec-

1570. RILEY, C. R.—Continued.

THE ROCKY MOUNTAIN LOCUST—Continued.

tion of fruit-trees, 110—Legislation, 111—Act passed by the Missouri legislature, 111—Acts passed by the Kansas legislature, 112—Act providing for the destruction of locusts in Minnesota, 114—Area in which eggs were laid, 116—Condition of eggs, 117—Temperature of the winter of 1876—77, 120—Prospects for 1877, 121.

INNOXIOUS INSECTS.

- The Hellgrammite, Corydalus cornutus.

 Its curious egg-mass described, 126—Resembling bird-dung at a distance, 126—Where laid, 127—The egg-burster, 127—Characters and habits of the newly-hatched larva, 127—Difficulty of rearing it in still water, 128—The eggs that have been hitherto mistaken for those of Corydalus, 128—They are probably those of Belostoma grandis, 128.
- 1571. RILEY, C. V. Important observations on the Rocky Mountain locust or grasshopper pest of the West. <Sci. Amer., 28 April, 1877 [v. 50], n. s., v. 36, pp. 260-261, 5 figs.
 - Observations on the egg-laying habits of Caloptenus spretus; figures egg, egg-masses, method of oviposition, and female anal characters.
- 1572. RILEY, C. V. Experiments with locust eggs and conclusions therefrom. <Sci. Amer., 5 May, 1877 [v. 50], n. s., v. 36, pp. 276-277. S.-b. No. 14, p. 101.
 - Experiments upon the vitality of eggs of Caloptenus spretus; effects of alternately freezing and thawing, of exposure to great moisture or to the free air, and of burying them at different depths.
- 1573. RILEY, C. V. Prof. Riley's report to the Governor of Kansas:
 The grasshopper question: Interesting information. <Commonwealth [Topeka, Kans.], 12 May, 1877, No. 2500, p. 2. S.-b.
 No. 14, pp. 63–67. Reprint: <St. Louis Daily Globe-Democrat,
 14 May, 1877, v. 2, No. 359, p. 3. S.-b. No. 14, pp. 70–72..
 Notice: <Ibid., p. 4. S.-b. No. 14, p. 70. See: <Kansas
 Farmer, 16 May, 1877. S.-b. No. 14, p. 59.
 - Reply to letter of Governor G. T. Anthony; area within Kansas in which eggs of Caloptenus spretus were laid in 1876; the degree to which the young from these eggs had perished or were likely to commit ravages in the summer of 1877; causes of the destruction of a great proportion of the young locusts; means of completing this destruction.
- 1574. RILEY, C. V. The strawberry leaf-roller, Anchylopera fragariæ. <Gardener's Mo. and Hortic., May, 1877, v. 19, pp. 143-144, fig. S.-b. No. 14, p. 45.
 - Reprint, with additional introduction and note, from 1st Ann. Rept. State Ent. Mo., March, 1869, pp. 142-143, fig. 80. See No. 1059 for synopsis of contents.

- 1575. [RILEY, C. V., et al.] U.S. Entomological Commission. Circular No. 1. [Riley, Packard, Thomas.] < Washington: 1877, 8°, 4 pp.
 - Queries regarding the migrations, appearances, habits, and ravages of Caloptenus spretus; directions for making replies.
- 1576. RILEY, C. V. U. S. Entomological Commission. Circular No. 2. <Washington: 1877, 8°, 4 pp.
 - Plans of work; request for information regarding the natural history, insect enemies, and parasites of *Caloptenus spretus* and other locusts; means against the same.
- 1577. [RILEY, C. V., et al.] Bulletin of the United States Entomological Commission. Destruction of the young or unfledged locusts. No. 1. [Riley, Packard, Thomas.] < Washington: 1877, 8°, 12 pp. See: <Sci. Amer., 2 June, 1877 [v. 50], n. s., v. 26, p. 344.
 - Enumeration of means of destroying young or unfledged locusts; quotations of laws passed by States of Missouri, Kansas, and Minnesota to provide for the destruction of locusts and their eggs.
 - Supplement to Bulletin No. 1, 2 pp. Summary of means for the destruction of young locasts.
- 1578. [RILEY, C. V., et al.] Bulletin of the United States Entomological Commission. On the natural history of the Rocky Mountain locust, and on the habits of the young or unfledged insects as they occur in the more fertile country in which they will hatch the present year. No. 2. [Riley, Packard, Thomas.] < Washington: May, 1877, 8°, 15 pp., figs. 1–11, map.
 - Description, oviposition, hatching, transformations, and habits of *Caloptenus* spretus; map of the country that will suffer most severely, showing the eastern limit of injury the present year.
- 1579. RILEY, C. V. The grape leaf-folder. <Journ. and Farmer, 14
 June, 1877. S.-b. No. 14, p. 133.
 - Description of larva, imago, habits, ravages, distribution, seasons, and foodplants of Desmia maculalis.
- 1580. RILEY, C. V. The rascal leaf-crumpler. <Journ. and Farmer, 14 June, 1877. S.-b. No. 14, pp. 133-134.
 - Description of larva, larva-case, imago, habits, ravages of, distribution, seasons, food-plants, and means against *Phycita nebulo* [= Acrobasis indiginella].
- 1581. RILEY, C. V. The Hessian-fly. <Journ. and Farmer, 14 June, 1877. S.-b. No. 14, p. 135.
 - Description of eggs, larva, puparium, and imago of Cecidomyia destructor; civil history, seasons, habits, ravages of and means against it; habits of Semiotellus [= Merisus] destructor.
- 1582. RILEY, C. V. Locust prospects. <Sci. Amer., 16 June, 1877, [v. 50], n. s., v. 36, p. 369.
 - Causes of the destruction of a great proportion of the young of *Caloptenus* spretus; a general and disastrous invasion improbable for some years.
- 1583. [RILEY, C. V.] Insect enemies. <N. Y. Tribune, 16 June, 1877. S.-b. No. 14, pp. 234–235.
 - 1. Apple-borers; answer to letter of V. B. P.; young apple-trees killed by Xyleborus obesus.

- 1583. RILEY, C. V.—Continued.
 - 2. Rose-chafer; answer to letter of I. M. H.; means against *Macrodactylus subspinosus*.
 - 3. Bark-louse; answer to letter of J. L. K.; seasons and ravages of and means against Mytilaspis pomicorticis [= pomorum].
 - 4. Flea-beetle; answer to letter of J. E. R.; habits, ravages, and means against Graptodera [= Haltica] chalybea.
- 1584. RILEY, C. V. The Hellgrammite. <Sci. Amer., 23 June, 1873 [v. 50], n. s., v. 36, pp. 392–393, 3 figs.
 - Structure and habits of *Corydalus cornutus*, description and figures of eggmasses, eggs, larvæ, pupa, and imagos; adaptations of sexual structure to conditions; figure and description of eggs of *Belostomå grande* [= americanum].
- 1585. RILEY, C. V. The 'hopper in Iowa. Report by Professor Riley, Chief of the U. S. Entomological Commission. <Chicago Daily Tribune, 4 July, 1877, v. 32, p. 3. S. b. No. 14, pp. 112–114. Extract: <N. Y. Tribune, 4 July, 1877. S. b. No. 14, p. 128.
 - Report to the governor of Iowa as to the observed and anticipated ravages of Caloptenus spretus in southwestern Iowa in 1877; the young locusts mostly destroyed by the weather; their natural enemies and the efforts of man; localities infested; recommends diversification of crops; summary of means against the young locusts; claims for the first recommendation of proper ditching; description of apparatus for catching and killing locusts.
- 1586. RILEY, C. V. Strawberry worm and remedy. <N. Y. Tribune, 18 July, 1877. S.-b. No. 14, p. 231.
 - Description of egg, larva, and imago, habits of and means against Emphytus [= Harpiphorus] maculatus.
- 1587. RILEY, C. V. Fighting the Hessian-fly. <N. Y. Tribune, 18
 July, 1877. S.-b. No. 14, pp. 232–233. Reprint: <Colman's
 Rural World, 5 December, 1877. S.-b. No. 14, pp. 268–269.
 - Habits, seasons, parasites, introduction, spread, and ravages of and means against *Cecidomyia destructor*; description of eggs and larva.
- 1588. RILEY, C. V. The grasshopper. Considered practically and scientifically with a retrospective and prospective glance at his history. <Daily Rocky Mountain News [Denver, Colo.], 1 August, 1877, v. 18, p. 4. S.-b. No. 14, pp. 81–84; 110–112. Reprint: <Colorado Farmer, 2 August, 1877, v. 9, No. 31, p. 4. <Chicago [Ill.] Daily Tribune, 7 August, 1877, v. 32, p. 7. S.-b. No. 14, pp. 84–86.
 - Observations on the past and present conditions of Caloptenus spretus in 1877 in the regions invaded by it; work of its enemies and parasites; complication of the locust problem in Colorado owing to the climate of the State; prospects of future injury; means against the locusts in Colorado; work of the U. S. Entomological Commission.
- 1589. RILEY, C. V. In reference to wheat-worms. <Prairie Farmer, 11 August, 1877. S.-b. No. 14, p. 127; No. 20, p. 86.
 - Critical review of C. Thomas' "A wheat insect" (Prairie Farmer, 21 July, 1877), with correction of some obscurities in regard to Meromyza americana and Gortyna nitela.

- 1590. RILEY, C. V. Locust flights. An appeal from Prof. C. V. Riley. <Manitoba Daily Free Press, 8 September, 1877. S.-b. No. 14, p. 91.
 - Reprint of questions of Circular No. 1 of U.S. Entomological Commission, with request for answers and other information from observers in Manitoba.
- 1591. RILEY, C. V. The locusts in Kansas. <Sci. Amer., 8 September, 1877 [v. 51], n. s., v. 37, p. 164.
 - No danger of locust invasion in Kansas and adjacent States in the fall of 1877; diseased conditions of late swarms; seasons of drought and locust swarms generally followed by rainy seasons and a scarcity of locusts.
- 1592. RILEY, C. V. A satisfactory grasshopper-machine. <Sci. Amer., 8 September, 1877 [v. 51], n. s., v. 37, p. 169.

 Description and figure of a machine for the killing of locusts.
- 1593. [RILEY, C. V.] Mistaken identity. <N. Y. Tribune, 12 September, 1877. S.-b. No. 14, p. 233.
 - Answer to letter of Subscriber; Lema trilineata mistaken for Doryphora 10-lineata; first occurrence of the latter in Vermont in 1876.
- 1594. RILEY, C. V. Injured orchard. <N. Y. Tribune, 12 September, 1877. S.-b. No. 14, p. 233.
 - Answer to letter of G. W. T.; occurrence of numerous nests of Formicidæ about the roots of orchard trees; probability that the ants are harmless; means against ants.
- 1595. [RILEY, C. V.] The stalk-borer. <N. Y. Tribune, 12 September, 1877. S.-b. No. 14, p. 233.
 - Answer to letter of Wisc.; food-plants of Gortyna nițela; description of larva.
- 1596. RILEY, C. V. Locust prospects in southwest Missouri this fall. <Journ. and Farmer, 27 September, 1877. S.-b. No. 14, pp. 236-237.
 - Ravages of Caloptenus spretus in Missouri in 1876 and prospects of same in 1877; history of locust flights in 1877.
- 1597. RILEY, C. V. The Colorado potato-beetle in Europe. German thoroughness. <Sci. Amer., 29 September, 1877 [v. 51], n. s., v. 37, p. 198. S.-b. No. 14, pp. 123-124.
 - Occurrence of *Doryphora* 10-lineata at Bremen and at Mülheim, in Germany; thorough measures taken to eradicate the pests; economic value of the measures.
- 1598. [RILEY, C. V.] The cussed red-leg. <Chicago [Ill.] Times, 29
 September, 1877. S.-b. No. 14, pp. 119–123. Reprint: <Mo.
 Rept. Kansas State Board of Agric., 1877, pp. 32–41. S.-b. No.
 14, pp. 146–161. Abstract: <Amer. Nat., November, 1877, v.
 11, pp. 663–673. <Ca. Nat. and Quart. Jour. Sci., December, 1877, v. 8, pp. 363–374.
 - Distinction between the terms locust and grasshopper; nature of varieties and species; distribution of Caloptenus spretus; laws governing its migrations and distribution; exemplification of these laws by records of flights during 1877; means against locusts; distinctions between Caloptenus spretus and C. femur-rubrum; habits and transformations of Astoma gryllaria (= the young of Trombidium sericeum [= locustarum]); habits of the latter.

- 1599. RILEY, C. V. [White-grub-fungus.] <N. Y. Weekly Tribune, 4 October, 1877. Notice: <Amer. Ent., June, 1880, [v. 3], n. s., v. 1, p. 140.

Views of earlier authors as to the habits of meloid larvæ; nature of the newly hatched larva (triungulin); natural history of Meloe and Sitaris, with reference to the literature, descriptions of the triungulins, especially of M. barbarus?; figures of the several stages; nature of hypermetamorphosis; food-habits and breeding habits of Macrobasis and Epicauta, especially of E. vittata; summary of what is known of the larval habits of other genera of Meloidæ; descriptions and figures of all the earlier stages of E. vittata; descriptions of eggs and triungulins of E. cinerea, E. pennsylvanica, and Henous confertus.

1601. RILEY, C. V. On a remarkable new genus in *Meloidæ* infesting mason-bee cells in the United States. <Trans. Acad. Sci. St. Louis, 16 November, 1877, v. 3, pp. 563–565, fig. 40; fig. 13 of pl. 5, with 1 p. expl. of pl. Separate: <[St. Louis, Mo., 1877], pp. 20–22, fig. 40; fig. 13 of pl. 5, with 1 p. expl. of pl. Abstract: <Amer. Nat., April [19 March], 1878, v. 12, pp. 218–219.

Description of imagos, ultimate stage of second larva, and coarctate larva of Hornia minutipennis, n. g. et n. sp., parasitic in cells of Anthophora sponsa [= abrupta]; figures & imago and coarctate larva of the same; illustrates the stages of degradation in tarsal claws of several genera of Meloidæ; synoptic table of the North American genera of Meloini.

- 1602. RILEY, C. V. Additional notes on Megathymus yucca. <Trans. Acad. Sci. St. Louis, 16 November, 1877, v. 3, pp. 566-568. Separate: <[St. Louis, Mo., 1877], pp. 23-25.
 - Supplementary to No. 1465; habits, seasons, and number of molts of the larva; description of larva in second and third stage and of variations in the imago; description of coloradensis n. var.; Egiale cofaqui a variety of Megathymus yuccw; Erynnis alcew bores in stems of Malva sylvestris in Europe; Elaphidion tectum and Scyphophorus yuccw in stems of Yucca.
- 1603. RILEY, C. V. Further remarks on *Pronuba yuccasella* and on the pollination of *Yucca*. <Trans. Acad. Sci. St. Louis, 16 No-

- 1603. RILEY, C. V.—Continued.
 - vember, 1877, v. 3, pp. 568-573. Separate: <[St. Louis, Mo., 1877], pp. 25-30. Translation: <Stett. Ent. Zeit., 1878, Jahrg. 39, pp. 377-382.
 - Critical review of remarks on *Pronuba* by Chambers, Zeller, and Boll; *Hyponomenta quinquepunctella* mistaken for *Pronuba yuccasella*; variations in the former; uniformity in the latter; behavior of *Pronuba* in flowers of *Yucca*; question of the method of fertilization of *Yucca*.
- 1604. RILEY, C. V. On the differences between Anisopteryx pometaria, Harr. and Anisopteryx ascularia W.-V., with remarks on the genus Paleacrita. <Trans. Acad. Sci. St. Louis, 16 November-20 December, 1877, v. 3, pp. 573-577. Separate: <[St. Louis, Mo., 1877], pp. 30-34.
 - Characters of Anisopteryx ascularia show that it is congeneric with A. pometaria; criticism of Packard's Monograph of Geometrida; structural characters separating Anisopteryx and Paleacrita.
- 1605. RILEY, C. V. Wheat rust and Hessian fly. <N. Y. Tribune, 19 December, 1877. S.-b. No. 14, p. 235.
 - Means against Puccinia graminis; P. graminis and Æcidium berberidis are alternate generations of one species.
- 1606. RILEY, C. V. A new oak-gall on acorn cups. <Trans. Acad. Sci. St. Louis, 20 December, 1877, v. 3, pp. 577-578.
 - Description of gall of Cynips quercus-glandulus n. sp. [p. 578] on cupules of all species of prinus group of Quercus; remarks on the allied structure of the galls of C. fecundatrix in Europe, and of C. quercus-frondosa; occurrence of galls of C. [= Amphibolips] quercus-prunus on cupules of acorns.
- 1607. [RILEY, C. V.] [Maggots in sauce.] <N. Y. Tribune, 26 December, 1877. S.-b. No. 14, p. 234.
 - Larvæ of *Drosophila* sp. in canned sauce; *D. nigricornis* common in similar substances.
- 1608. [RILEY, C. V.] [Round-headed apple-tree borer.] <N. Y. Tribune, 26 December, 1877. S.-b. No. 14, p. 234.

 Ravages of and means against Saperda bivittata [= candida].
- 1609. RILEY, C. V. Entomological notes. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 217-218 Proc. See: <Amer. Nat., October, 1876, v. 10, p. 635.
 - Generic resemblance of Carpocapsa saltitans to C. pomonella; correction of vernacular name of Cicada [= Tibicen] septendecim; occurrence of the same in Virginia in 1876; yearly development of C. [= T.] tredecim; Sericaria mori reared for five years on Maclura aurantiaca with increased vigor and healthfulness; Salix nigra stripped by larvæ of Vanessa antiopa; habits of the same.
- 1610. RILEY, C. V. Entomological notes. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 219 Proc. See: <Colman's Rural World, 1876. S.-b. No. 10, p. 165. <N. Y. Tribune, 1876. S.-b. No. 9, p. 215. <Amer. Nat., October, 1876, v. 10, p. 635.
 - Occurrence and ravages of Leucania albilinea in Kansas; ectoparasitism of Uropoda americana on Doryphora decemlineata; list of vertebrate enemies of

- 1610. PILEY, C. V.—Continued.
 - D. decemlineata; occurrence of D. decemlineata in New Hampshire; its ravages along the Atlantic coast; see No. 1505.
- 1611. RILEY, C. V. Centennial insects. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 220-221 Proc.
 - List and characterization of insects observed injuring exhibits in the Centennial Exposition.
- 1612. RILEY, C. V. Parasites on eggs of Caloptenus spretus. < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 226 Proc.
 - A large proportion of locust eggs throughout the country are being destroyed by parasites, predaceous insects, and other causes; precautions against threatened injury; discovery of four new enemies of locust eggs.
- 1613. RILEY, C. V. Locust flights east of the Mississippi. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 226-228 Proc. Locust swarms in Ohio, Illinois, Georgia, and South Carolinia not composed of Caloptenus spretus; causes which limit the eastward flight of C. spretus.
- 1614. RILEY, C. V. Geographical range of species. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 230-233 Proc.
 - Range of species in our time affected only by the interposition of human influence; the limits of this range definitely established by classified knowledge, and impassable, except by man's assistance; Caloptenus spretus can not change its habits; the difference in the susceptibility of different species to change inexplicable.
- 1615. RILEY, C. V. [Anticipated locust injury next summer.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 236 Proc.
 - Abundance and advanced state of development of eggs of Caloptenus spretus in the States invaded in 1876; probable abundance and ravages in 1877.
- 1616. RILEY, C. V. [Japanese mode of packing silk-worm eggs.]

 Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 236
 Proc.
 - Mode in which the eggs of Sericaria mori are packed in Japan for transportation.
- 1617. RILEY, C. V. Locust experience. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 267 Proc. See: <Colman's Rural World, 1877. S.-b. No. 14, p. 265.
 - Northern limit of the permanent breeding-grounds of Caloptenus spretus; confirmation of views regarding laws governing locust invasions; discovery of a new law.
- 1618. RILEY, C. V. Mite transformations. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 267–268 Proc. See: <Colman's Rural World, 1877. S.-b. No. 14, p. 265.
 - Habits and transformations of *Trombidium sericeum* [= T. locustarum]; Astoma gryllaria an immature form of the same.
- 1619. RILEY, C. V. [Mygale hentzii and Pepsis formosa.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 269 Proc.
 - The bite of Mygale hentzii not so deadly as is generally supposed; habits of Pepsis formosa in preying on the spider.

- 1620. RILEY, C. V. [Ravages of Termes flavines.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 269 Proc.
 - Note to communication of R. D. Grant on some ravages of Termes flavipes.
- 1621. RILEY, C. V. On the oviposition of Saperda bivittata Say. < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 269-270 Proc. See: < Colman's Rural World, 28 November, 1877. S.-b. No. 20,
 - Description of egg and manner of oviposition of Saperda bivittata [=candida].
- 1622. RILEY, C. V. On migratory butterflies. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 273-274 Proc. See: < Valley Naturalist, January, 1878, v. 1, p. 2.
 - Phenomena and causes of the migrations of butterflies, especially of Danais archippus.
- 1623. RILEY, C. V. [Phylloxera and grape-rot.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 275 Proc. See: < Colman's Rural World, 28 November, 1877. S.-b. No. 14, p. 268.
 - Disagreement with views of A. J. Cook as to there being any connection between the work of Phylloxera vastatrix and the ordinary grape-rot.
- 1624. RILEY, C. V. The periodical Cicada. < Western Farmer's Almanac for 1878, 1877, p. 48. S.-b. No. 14, p. 138. Reprint: < Colman's Rural World, 28 November, 1877. S.-b. No. 14, pp. 265-266.
 - Regular periodicity of the appearance of Cicada [= Tibicen] septendecim and C. [=T.] tredecim; popular description and natural history of the same; chronology of twenty-two different broods, geographically classified.
- 1625. RILEY, C. V. The locust plague in the United States: being more particularly a treatise on the Rocky Mountain locust, or so-called grasshopper, as it occurs east of the Rocky Mountains, with practical recommendations for its destruction. < Chicago: Rand, McNally & Co., 1877, 236 pp., 3 pl., 42 figs. tract: < Manitoba Standard, 1877. S.-b. No. 14, pp. 97-98. Notice: < Cultivator and Country Gentl., 30 August, 1877, v. 42, p. 557. < Gardener's Mo. and Hortic., October, 1877, v. 19, p. 317. <Ent. Mo. Mag., October, 1877, v. 14, p. 118.

TABLE OF CONTENTS.

Preface	7
Introduction	
CHAPTER I:	
Characters of the species	13

Characters of the species Classificatory position of the Rocky Mountain locust, 13-How it. differs from other species, 14-Easily confounded with the common red-legged locust, 14-Detailed descriptions of both, 15-Its still closer resemblance to the Atlantic migratory locust, 22— Characters of this last, 22—Species vs. variety and race, 23—Comparisons of these three closely-allied species in their early stages, 26-A green variety of the Rocky Mountain locust not infrequent, 27-Purely an American insect, 28.

	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	219
1625.	RILEY, C. V.—Continued. CHAPTER II:	
	Chronological history The locust plague in the "Old World," 29—Extent of its injuries, 30—Migratory species in Europe, Asia, and Africa, 30—The ravages of the locust in America, 31—Its earliest visitations, 31—Injuries on the Pacific coast, 32—Injuries east of the Rocky Mountains, 33—Invasions of 1818–'19, 33—Of 1845–'49, 34—Of 1855, 34—Of 1856, 35—Of 1857–'67, 35, 36—Of 1866, 36—Damage the following year, 37—The invasion of 1873, 38—That of 1874, 39—Why so disastrous, 41—General outlook in the spring of 1875, 42—Severity of the injuries from the young insects that year, 43—Destination of that prevailed, 44—Amount of loss sustained, 45—Destination of departing swarms of 1875, 47—The invasion of 1876, 49—Eastern limit reached, 53—Omaha conference, 53.	2
	CHAPTER III: Native home and geographical range of the species east of the	
-	mountains Source of the devastating swarms that reach into the Mississippi Valley, 55—Their origin in the extreme Northwest country lying east of the mountains, 56—Cause of their emigration, 57—Difference between summer and fall swarms, 58—The species not at home in the Mississippi Valley, 62—Not permanent or able to perpetuate itself there, 62—Conditions which prevent such permanence, 63—Not likely to do serious harm east of the ninety-fourth meridian, 65—Reasons why, 65.	58
•	CHAPTER IV:	
	Natural history and transformations How the eggs are laid, 69—Philosophy of the egg-mass, 71—The female capable of laying more than one egg-mass, 72—The escape of the young locust from the egg, 73—Its emergence from the ground, 78—Growth and transformations, 78—Acquisition of wings, 80—Number of molts, 82—Flight at night, 83.	69
	CHAPTER V:	
	Habits and power for injury Flight and ravages, 85—Migratory instinct and great destructive power confined to a single specie's west of the Mississippi, 88— Food-plants, 89—Injury to fruit trees, 93—Time of appearance of invading swarms, 94—Rate at which they spread, 95—Direction of their flight, 96—Where the eggs are preferably laid, 96—Time of hatching, 97—Habits of the young or unfledged locusts, 98—Directions in which the young travel, 100—Rate at which they travel, 100—Limit of their east ward spread, 100—Not led by kings or queens, 101—Direction taken by the departing swarms, 103—Their destination, 104.	85
	CHAPTER VI:	
	Effects of the young insects in the country where they hatch, but where they are not indigenous Experience with the young locusts in spring, 107—Contrast in summer and fall, 108—No evil without some compensating good,	107

108-Changes that follow the locusts, 109-The prevalence of large green worms, 110-The sudden appearance of a peculiar

grass, 110.

1625. RILEY, C. V.—Continue	1625. Ri	LEY.	U.	V.—	Coi	ntin	ued
-----------------------------	----------	------	----	-----	-----	------	-----

CH	APTER	≀ VII •

Natural enemies of the Rocky Mountain locust

Birds and other vertebrate animals, 113—The good offices of birds.

113—Invertebrate animals, 114—Animals that attack the eggs,
115—The silky mite, 115—The Anthonyia egg-parasite, 118—The
common flesh-fly, 123—Undetermined larvæ, 123—Ichneumonflies, ground-beetles, click-beetles, and myriapods, 126—Insects
that destroy the active locust, 126—The locust-mite, 128—The
anonymous Tachina-fly, 131—Yellow-tailed Tachina-fly, 134—
Flesh-fly, 135.

113

139

187

207

CHAPTER VIII:

Practical considerations How best to prevent locust injuries, 139—Encouragement of natural enemics, 139—Destruction of the eggs, 139—Experiments with the eggs and conclusions drawn therefrom, 140-Effects of alternately freezing and thawing, 141—Influence of moisture, 143—Exposure to the free air, 148—Burying at different depths, and pressure of the soil, 149—Table of temperatures, 152—Harrowing in the fall, 153—Collecting the eggs, 153—Plowing, how most effectual, 153-Irrigation, tramping, 154-Destruction of the young or unfledged locusts, 155-Burning, 155-Crushing, 157—Trapping, 157—Ditching and trenching, 161—Catching, 161-Different contrivances for this purpose, 162-Use of destructive agents, 162-Coal-oil pans, 163-The use of coal-tar, 164—The protection of plants by special applications, 165—The best means of protecting fruit and shade trees, 166-Sulphur fumes and smudges, 167—Destruction of the winged locusts, 167-Preventive measures, 169-Suggestions that may prove of service, 170-Use of hogs and poultry, 171-The Signal Service, 172—Military aid, 173—Diversified agriculture, 174—Organized effort, 176-State legislation, 176-Missouri locust law, 177-Kansas locust laws, 178, 179-Minnesota locust law, 180-Nebraska locust law, 184-How to avert locust invasions, 186.

CHAPTER IX

CHAPTER X:

General considerations

Nomenclature, 207—Locust vs. grasshopper, 207—Prairic fires vs. locust ravages, 209—Fasting and prayer, 213—Not a divine visitation, 216—Influence of the wind in determining the course of locust flights, 216—Locusts as food for man, 217.—Unnecessary alarm caused by comparatively harmless species, 227.

1626. RILEY, C. V. On an extensile penetrating organ in a gamasid mite. <Proc. Amer. Assoc. Adv. Sci. for 1876, 1877, v. 25, pp. 273–275, 1 fig. See: <Ca. Ent., September [October], 1876, v. 8, p. 180.

- 1626. RILEY, C. V.—Continued.
 - Habits and description of *Uropoda vegetans* and *U. americana* n. sp.; nature of the adhesive filament of these mites; structure of the supposed maxillary penetrating organ in *Uropoda*.
- 1627. RILEY, C. V. On the curious egg-mass of Corydalus cornutus, Linn., and on the eggs that have hitherto been referred to that species. <Proc. Amer. Assoc. Adv. Sci. for 1876, 1877, v. 25, pp. 275–279, 1 fig. Reprint: <9th Ann. Rept. State Ent. Mo., March, 1877, pp. 125–129, figs. 30–33. See: <Ca. Ent., October [November], 1876, v. 8, pp. 181–182. See No. 1570 for synopsis of contents.
- 1628. RILEY, C. V. Phylloxera and grape-rot. <N. Y. Tribune, January, 1878. S.-b. No. 19, pp. 193; 224. Reprint: <Colman's Rural World, 1878. S.-b. No. 14, pp. 266-267. Extract: <Gardener's Mo. and Hortic., July, 1879, v. 21, pp. 213-214. S.-b. No. 23, pp. 118; 143-145. <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 237-238. See: <Gardener's Chronicle, 9 August, 1879. S.-b. No. 26, p. 15. See No. 1721 for synopsis of contents.
- 1629. RILEY, C. V. Address by Prof. C. V. Riley, retiring, before the St. Louis Academy of Science. <St. Louis Republican, 8 January, 1878. -S.-b. No. 14, p. 269. Reprint: <Trans. Acad. Sci. St. Louis, 1880, v. 4, pp. 1–6 Proc.
 - Review of progress made in science in 1876; success of the investigation into the Rocky Mountain locust scourge.
- 1630. RILEY, C. V. New facts about the round-head apple-tree borer. <N. Y. Weekly Tribune, 20 February, 1878. S.-b. No. 19, pp. 193–194. Reprint: <Colman's Rural World, 20 March, 1878. S.-b. No. 14, pp. 267–268. Reprint, with slight omission: <Ibid., 12 March, 1879. S.-b. No. 23, pp. 101–102. See: <Mirror and Farmer, 6 April, 1878. S.-b. No. 20, p. 3.
 - Description of egg and manner of oviposition of Saperda bivittata [= can-dida]; seasons of oviposition; means against the same.
- 1631. [RILEY, C. V.] Buggy beans. <N. Y. Tribune, 20 February, 1878. S.-b. No. 14, p. 234.

 Habits and distribution of and means against Bruchus faha: edibility of the
 - Habits and distribution of and means against Bruchus faba; edibility of the same.
- 1632. RILEY, C. V. On the transformations of the red mites. < Amer. Nat., March [23 February], 1878, v. 12, pp. 139-146, figs. 1-6. Habits, description, and figures of Trombidium locustarum [p. 142], T. giganteum [p. 143], T. muscarum [p. 144], and Hydrachna belostoma [p. 146], n. sp.; figures and description of most of the stages of the same; definition of
- 1633. RILEY, C. V.] Clothes moths. <Sci. Amer., 23 March, 1878 [v. 52], n. s., v. 38, p. 177. S.-b. No. 19, p. 238. Reprint: <Kansas Farmer, 17 April, 1878. S.-b. No. 19, p. 225. <Colman's Rural World, 3 April, 1878. S.-b. No. 19, pp. 198–199.

Trombidium. [Advance extract from No. 1643.]

- 1633. RILEY, C. V.—Continued.
 - Indiscriminate use of names and habits of several species of *Tinea* injurious to woolen goods, fur, hair, and similar substances; means against the moths.
- 1634. RILEY, C. V. Pieris vernalis and P. protodice. <Ca. Ent., February [March], 1878, v. 10, p. 39.
 - Conformation of T. E. Bean's conclusion that P. vernalis is the spring form of P. protodice.
- 1635. RILEY, C. V. Migratory butterflies. <Sci. Amer., 6 April, 1878 [v. 52], n. s., v. 38, p. 215, fig. S.-b. No. 19, p. 250. Extract: <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 102, fig. 34.
 - Occurrence of migratory swarms of Danais archippus in central southern United States in the autumn of 1877, and of return migrations in spring; explanation of these migrations; mentions other migratory Rhopalocera; distribution of Danais archippus and Cynthia [= Pyrameis] cardui; figure of Danais archippus.
- 1636. RILEY, C. V. The horn-bug. <Sci. Amer., 20 April, 1878 [v. 52], n. s., v. 38, p. 249, 1 fig. S.-b. No. 19, p. 241.

 Natural history, description of egg, larva and pupa, and figures of larva, pupa, and imago of Passalus cornutus.
- 1637. RILEY, C. V. Egg-feeding mites. <Ca. Ent., March [April], 1878, v. 10, pp. 58-59.

 Citation of records of Acari feeding on eggs.
- 1638. [RILEY, C. V.] Bad work of the grain Aphis <N. Y. Tribune, 26 June, 1878. S.-b. No. 19, p. 224.

 Habits and ravages of and means against Aphis avenæ [= Nectarophora granaria].
- 1639. [RILEY, C. V.] Of Doryphora. <N. Y. Tribune, 26 June, 1878. S.-b. No. 19, p. 243.
 - Natural history, enemies, and spread of *Doryphora decemlineata*; description of larva and imago; methods of using Paris green.
- 1640. RILEY, C. V. The apple-tree borer. <N. Y. Tribune, 24 July, 1878. S.-b. No. 19, p. 195.

 Means against Chrysobothris femorata.
- 1641. [RILEY, C. V.] The raspberry saw-fly. <N. Y. Tribune, 24 July, 1878. S.-b. No. 19, pp. 197; 226.

 Habits of and means against Selandria [= Monophadnus] rubi.
- 1642. [RILEY, C. V.] Inquiring friends. <N. Y. Tribune, 24 July, 1878.
 S.-b. No. 19, p. 198.

 Means against Ægeria tipuliformis.
- 1643. [RILEY, C. V., et al.] First annual report of the United States Entomological Commission for the year 1877, relating to the Rocky Mountain locust and the best methods of preventing its injuries and of guarding against its invasions, in pursuance of an appropriation made by Congress for this purpose. With map and illustrations. < Washington, 29 July, 1878, pp. 16+

1643. RILEY, C. V.—Continued.

477+ 295, 111 figs., 5 pls., 2 maps. See: <Sci. Amer., 5 July, 1879 [v. 55], n. s., v. 41, p. 8. S. b. No. 23, p. 121.

TABLE OF CONTENTS.

DETTER OF TRANSMITTAL	ΔI
LETTER OF SUBMITTAL	XIII
Preface	xv
Introduction: By C. V. Riley.	
Introductory remarks	1
Creation and organization of the Commission, 1-Division of labor,	
1—Circulars sent out, 2-6—Area over which eggs were laid in	
1876, 6—Outlook in spring in more southern States, 7—Letter to	
Governor Anthony, of Kansas, on the condition of things in May,	
8-10—State of things in Minnesota in spring, 11—Ontlook in Ne-	
braska in June, 13—Second meeting of the Commission, 14—	
Outlook in Iowa in June, 15—Visit to Colorado, Utah, and Mon-	
tana in June, 17—Visit to Colorado in July, 17–20—Third meet-	
ing of the Commission, 21—Trip to the Pacific coast, 21—Trip to	
British America, 22—List of chapters, 23—Character and synop-	
sis of chapters, 24-29—Prospects for 1877, 29.	
CHAPTER I: By C. Thomas.	
Classification and nomenclature; characters of the species	13
Families of the Orthoptera, 32—Locust vs. grasshopper, 33—Ge-	
neric nomenclature, 37—Generic diagnosis, 40—Species of the	
genus Caloptenus, 42-Full definition of Caloptenus spretus and	
of its nearest congeners, 43-52.	
CHAPTER II: By C. Thomas.	
Chronological history of locust injuries	53
Compared with locust ravages in the Eastern Hemisphere, 53-In-	
jury in the Northwest early in the present century, 54-Brief	
reviews from 1820 to 1877, 54-56—Locust history in Texas, 57-62;	
in Indian Territory, 63; in Arkansas, 64; in Missouri, 64-68; in	
Kansas, 68-74; in Nebraska, 74-77; in Iowa, 77-80; in Minne-	
sota, 80-87; in Dakota, 88-92; in Montana, 92-96; in Idaho, 96;	
in Wyoming, 97; in Colorado, 99-102; in Utah, 102-104; in New	
Mexico and Arizona, 105; in Nevada, 105; in Oregon and Wash-	
ington Territory, 106; in British North America, 108-112-Tabu-	
lar view of locust years, 113.	
CHAPTER III: By C. Thomas.	
Statistics of losses	114
Difficulty of obtaining reliable data, 114—Estimates in Kansas,	114
115; in Minnesota, 116; in Missouri, 117—Loss in Kansas, Ne-	
braska, Iowa, and Missouri in 1874, 118—Loss to different crops,	
120—Loss to Missouri in 1875, 121—Total loss during the years	
1874-77, \$200,000,000, 122.	
CHAPTER IV: By C. Thomas.	100
Agricultural bearing of the locust problem	123
Drawback to the settling of the West, 124—What is likely to be	
the effect in the future, 125—Modification and settlement of the	
Western plains, 129—Crops which suffer most, and those which	
suffer least, 130—Small grains not affected by invading swarms,	
128—Need of judgment in planting, 129.	

1643. RILEY, (J.	$\nabla \cdot \cdot$	Co	ntin	ued.
----------------	----	----------------------	----	------	------

RILEY, C. V.—Continued.	
CHAPTER V: By A. S. Pachard, jr.	
Permanent breeding-grounds of the Rocky Mountain locust	131
Permanent breeding-grounds, 131—Definition of the permanent re-	
gion, 133—Its character and extent, 134—The Rocky Mountain	
locust a sub-borcal insect, 135—The sub-permanent region, 136—	
The temporary region, 136.	
CHAPTER VI: By C. Thomas and A. S. Pachard, jr.	
Geographical distribution	136
Eastern limits of spread, 137—Northern limits, 139—Western lim-	
its, 140—Southern limits, 141—Sub-permanent region, 142.	
CHAPTER VII: By A. S. Pachard, jr., and C. Thomas.	
Migrations	143
Classification of flights, 143—Invading swarms, 143—Returning	
swarms, 143—Local flights, 143—Height at which swarms move,	
144—Effect of change of wind and weather on flights, 145—	
Flight at night, 147—Migrations previous to 1877 east of the	
Rocky Mountain Plateau, 148-Direction of invading swarms	
prior to 1877, 149—Direction of flight in 1876, 151—Migrations	
within the permanent region, 153-158; in Montana, 153; in	
Wyoming, 156; in Colorado, 157; in Eastern Idaho and Utah,	

158; in the lower Snake Valley, 158—Return migrations from the temporary region previous to 1877, 159-162—Return migrations in 1877, 162-165—Local flights in 1877, 165—Record of flights for July 3 and July 20, 1877, 165-169—Summary of flights by States, 170-174—Southward flights in 1877, 175—Movements in different directions at one time, 176—Destination of return swarms, 177.

212

CHAPTER VIII: By C. V. Riley.

Habits and natural history..... Destructive powers of locusts, 212-215-Stoppage of railroad trains, 215—Rate at which locust swarms move, 215—Velocity of flight, 216—Direction of invading swarms, 217—Time of appearance of invading swarms, 217-Flight at night, 218-Height of flight, 219—General habits at night, 219—Where the eggs are laid, 222— Manner in which the eggs are laid, 223—Philosophy of the eggmass, 225—The female lays more than one egg-mass, 226—Interval between different egg-layings, 227—Number of eggs laid, 228-The hatching process, 228-231-Where and under what conditions of soil the young hatch most freely, 231-Time of hatching, 231-Habits of the young or unfledged locusts in the temporary region, 232—Directions in which the young travel, 234— Rate at which the young travel, 235—They reach but a few miles east of where they hatch, 235—Not led by kings and queens, 236— Time of year when wings are acquired, 237—Direction taken by swarms departing from the temporary region, 238—Destination of departing swarms, 238—Do the return swarms breed? 239—Do return swarms from the temporary region retrace their course? 240—The species essentially single-brooded, 240-243—Reasons why it cannot produce two generations annually, 243—The species can not permanently dwell in the temporary region, 244-The insects which hatch there do not remain, 246—Extensive and thick egg-laying seldom occurs twice consecutively in the same locality, 247—Reasons why it does not, 248—Causes of migration, 249 - Food-plants, 251-Crops and plants most liked and those feast liked, 252-254-Unnecessary alarm caused by comparatively harmless species, 255.

1643.	RILEY, C. V.—Continued.	
	CHAPTER IX: By A. S. Pachard, jr.	
	Anatomy and embryology	257
	External anatomy, 257—Divisions of the body, 258—Sexual differ-	
	ences, 259—Internal anatomy, 261—The digestive system, 262-	
	264. —The nervous system, 264—The heart, 266—The tracheæ and	
	dilated air-sacs, 267-Mode of breathing, 269-The locust an	
	aëronaut, 270—The male reproductive system, 270—The female	
	reproductive system, 271—Organs of special sense, 272—The fine	
	anatomy of the locust, 273-Embryology of the Rocky Mountain	
	locust, 277—Different egg-layers, 278—How the embryo lies	
	within the egg, 278—How it bursts the egg, 279.	•
	CHAPTER X: By C. V. Riley.	
	Metamorphoses	279
	The six stages of growth, 279-Variability in the depth of coloring,	
	281—The process of molting, 281—Time required for it, 283—Dif-	
	ferences in the immature stages between the Rocky Mountain,	
	the lesser, and the red-legged locusts, 283.	
	CHAPTER XI: By C. V. Riley.	
	Invertebrate cnemies	284
	Value of the locust's minute cnemies, 284—Animals that destroy	
	the eggs, 285—The Anthomyia egg-parasite, 285—The common	
	flesh-fly, 289—Ground-beetles and their larvæ, 289—Harpalus	
	larvæ, 289—The egg-feeding Amara, 291—Blister-beetle larvæ,	
	292—Their character and locust egg-feeding habits, 293—History	
	of the oil-beetle, 294—History of Sitaris, 295—History of Hornia,	
	296-History of Epicauta, 297-Macrobasis and Henous, 301-	
	Other meloid genera, 302—Soldier-beetle larvæ, 302—Asilid	
	larvæ, 303—Click-beetle larvæ, 304—Miscellaneous species, 305—	
	Chalcid-fly, 306—Animals that prey on the locust after it is born,	
	306—The locust-mite, 306—The efficacy of its work, 308—Its	
	transformations, 309—Other mites, 312, 313—Ground-beetles,	
	313—Tiger-beetles, 314—Asilus-flies, 317—Digger-wasps, 317—	
	Tachina-flies, 319—Their efficacy in destroying locusts, 321—	
	Flesh-flies, 323—Ichneumou-flies, 324—Hair-worms, 326—Their	
	curious life-history, 327-332-Insects attacked by hair-worms,	
	327—How hair-worms get into locusts, 332—Miscellaneous locust	
	enemies, 334.	
	CHAPTER XII: By C. Thomas.	
	Vertebrate enemics	334
	Good offices of birds probably underrated, 334—Experience of cor-	
	respondents, 336—Some of the most useful birds, 338—Paper by	
	Professor Aughey on the beneficial work of birds, 338—Enormous	
	number of birds destroyed for market, 346—Damage done to in-	
	sectivorous birds by birds of prey, 348—The English sparrow,	
	349—What public sentiment necds, 349.	
	CHAPTER XIII: By C. V. Riley.	
	Remedies and devices for destruction	350
	Encouragement of natural agencies, 351—Destruction of the eggs,	
	351—Harrowing in the autumn, 353—Experiments to test the	
	effect of alternately freezing and thawing, 353—Experiments to	
	test the effects of exposure to air, 354—Experience with harrow-	
	ing, 355—Plowing, 356—Experiments to test the effects of bury-	
	ing at different depths, 356—Experience in plowing, 358—Irriga-	
	tion, 359-Experiments to test the effects of moisture on the	
	13 ENT——15	

1643. RILEY, C. V.—Continued.

CHAPTER XIII-Continued.

eggs, 359—Tramping, 361—Collecting, 361—Destruction of the young or unfledged locusts, 362—Burning, 363—The burning of prairies, 363—The Hetzel burning machine, 363—The Horner burning contrivance, 363-Hand burners, 364-The Atwood machine, 364—Use of wire and kerosene, 364—Crushing, 364—The Drum locust-crusher, 365—The Simpson locust-crusher, 366— The Hoos locust-crusher, 367—The Hansberry locust-crusher, 368—The Kenworthy locust machine, 370—J. C. Melcher's machine, 371—The Peteler machine, 371—The King suction-machine, 374—The Flory locust-machine, 376—Trapping locusts, 377—Nets and seines, 377—Ditching and trenching, 378—Protection by barriers, 381—Coal-oil, 381—Coal-oil pans, 383—The Canfield pan, 384—The Adams pan, 385—The Anderson coal-oil contrivance, 386-Mr. Long's contrivance, 386-Mr. Watrous's contrivance, 387-Mr. Swearingen's contrivance, 387-Use of coaltar, 387—The Robbins pan, 388, 390—Use of coal-tar previous to 1877, 388-Other machines for the use of coal-tar, 391-Catching and bagging, 391-Principles and facts to be borne in mind in catching locusts, 391—The Riley machine, 392—Mr. Thompson's net, 392—The Elliot catcher, 394—The Wilson-Rhode catcher, 395—Contrivance for catching the pupe, 394—The Godard catcher, 395—The Benson catcher, 396—The Hutchins catcher, 396-The Sylvester catcher, 397-The Horo hoppor-catcher, 398-The Belt device, 398—Hand-nets, 399—Use of destructive agents, 399—Buhach or Persian insect-powder, 400—Veith's insect-fluid, 400—Sulpho-carbonate of potassium, 401—Naphthaline, 401— Paris green, 401—The protection of fruit trees, 403—Destruction of the winged insects, 404—Fumigation, 405—The effect of concussion, 405—Diversified agriculture, 406—Legislation, 407— Bounty laws, 409—Missouri locust act, 409—Kansas locust acts, 410-Minnesota locust acts, 412-Nebraska locust act, 413-Suggestions that may be of service, 414-More attention needed to the growth of root-crops, 415-The benefits of irrigation, 415-Hogs and poultry, 415-Non-planting, 415-Use of soldiers, 416-Preventive measures against the winged insects, 417—Further investigation needed, 417-Means that have been suggested against the winged insects, 418-Systematic burning of young, 418-Co-operation with Dominion Government, 419-Protection by smoke, 419—Farmers should receive the locust probabilities,

CHAPTER XIV: By C. V. Riley.

Influence of prairie fires on locust increase.

Reasons given why prairie fires do not influence locust increase,
421—Possible indirect connection in the past, 422.

Influence of weather on the species.....

421

423

CHAPTER XV: By C. V. Riley.

Effects of weather on the young locusts, 424—Effects of weather on the eggs, 424—Possibility of predicting consequences upon meteorological grounds, 424—The temperature of the soil, 425—Thermal constant for the hatching of locust eggs, 426—Number of hours which eggs must be exposed to hatch at a temperature of 60° F., 431—Number of hours required in 1875.

164	3. RILEY, C. V.—Continued.	
	CHAPTER XVI: By C. V. Riley.	
	Effects that generally follow severe locust injury	432
	Contrast between summer and autumn, 432—No evil without some	
	compensating good, 433—Changes that follow the locusts, 433—	
	Sudden appearance of plants and insects not ordinarily noticed,	
	434—The white-lined morning Sphinx, 435—Tame grasses per-	
	manently injured, wild grasses not, 435—Permanent effects on	
	different plants, 436—Injury to fruit trees, 436.	
	CHAPTER XVII: By A. S. Pachard, jr.	400
	Uses to which locusts may be put.	437
	Locusts as food for man, 438-441—As fish-bait, 441—Analysis of locust-juices, 442—They furnish a large quantity of formic acid	
	and a new oil, 442-443—Locusts as manure and as poultry food,	
	443.	
	CHAPTER, XVIII: By C. V. Riley.	
	Ravages of other locusts in the United States	443
	Ravages of locusts that are occasionally migratory in the Atlantic	
	States, 443—Great destruction in past years in New England,	
	444—The most common species concerned in this work, 446—	
	Locust flights in Illinois, 446—The species composing them, 447—	
	The phenomenon exceptional, 447—Locusts ordinarily non-migra-	
	tory, occasionally become so under favoring conditions, 448-	
	Locust flights in Ohio, 449—Importance of discriminating be-	
	tween species, 450—Geographical limits of species, 450—The mi-	
	gratory locusts of the Pacific, 451—Chronological account of	
	their ravages, 451—Species concerned, 452—Late injuries in	
	southern California, 454—Tabular view of locust years, 456—Injury from other non-migratory locusts, 456—The species con-	
	cerned, 459.	
	CHAPTER XIX: By A. S. Pachard, jr.	
	Ravages of locusts in other countries	460
	Locust injuries in Central America, 460—Great destruction in	100
	Honduras and Guatemala, 462—The locusts in South America,	1
	465—The locusts in the Old World, 467—Injuries in Germany,	
	468—Habits of the European species, 469—Injuries in Russia,	
	470—Different species affecting the Old World, 471—Geographical	
	distribution of the European migratory locust, 472-476—Notes	
	on Algerian locusts, 476—Injury in China, 477—In southern Aus-	
	tralia, 477.	
	APPENDICES.	
	APPENDIX I:	
	Mr. Allen Whitman's report from Minnesota	[3]
	State of things in spring, [4]—Exertions of farmers, [4]—Different	
	rieans employed to destroy the eggs, [4]—Effects of temperature	
	and rain-fall on the eggs, [5]—Dates of hatching, [6]—Late	
	hatching, [7]—Progress during spring, [8]—First winged, [10]—	
	Statistics of damage, [11].	
	APPENDIX II:	F107
	Aughey on locust-feeding birds.	[13]
	Letter of transmittal, [13]—Examinations of the contents of the	
	stomachs of birds of Nebraska, giving the number of locusts, number of other insects, and number of seeds of the different	
	birds considered in their proper classificatory position, [14]-[62].	
	Land of the Land o	

1643.	RILEY, C. V.—Continued.	
	APPENDIX III:	F.007
	Report of Jacob Boll, special assistant, [62]—Reports from correspondents and other miscellaneous reports, chronologically arranged, [64]-[82].	[63]
	APPENDIX IV:	
	Missouri data for 1877	[83]
	APPENDIX V:	
	Kansas data for 1877	[85]
	APPENDIX VI:	
	Iowa data for 1877	[104]
٠	APPENDIX VII:	
	Colorado data for 1877 Report from William Holly, special assistant, [111]—Data from correspondents and from other sources, [113]-[116].	[111]
	APPENDIX VIII:	
	Nebraska data for 1877	[116]
	APPENDIX IX:	
	Narrative of the first journey made in the summer of 1877, by Mr. Packard Diary notes through Kansas, Colorado, and Utah, [135]—Observa-	[134]
	tions in Montana, [137]—Results of the journey, [138]. APPENDIX X:	
	Narrative of a second journey in the summer of 1877, by Mr. Packard.	[1397
	From Chicago to Utah, [139]—Observations in Oregon and Washington Territory, [140]—From Vancouver Island to California, [141]—Variations in Caloptenus spretus, atlanis, and femur-rubrum, [143].	
	APPENDIX XI:	
	British-American data	[145]
	Characteristics and prevailing winds of Manitoba, [145]—Data from the Cypress Hills region, [146]—The locust breeds permanently in the third prairie steppe, [146]—No damage ever done	
	in the Peace River country, [146]—Ravages in the Northwest in the early part of the century, [147]—Records by the Hon. Don-	
	ald Gunn, [148].	
	APPENDIX XII:	F4#0#
	Autumn flights in 1877 Southward movements in Minnesota and Iowa, [150]—In Nebraska and Missouri, [151]—In Arkansas and Kansas, [152]—In Colorado and Texas, [155]	[150]

643. RILEY, C. V.—Continued.	
APPENDIX XIII:	
Flights and migrations in 1877	[156]
region, [156]-[164]—Flights in the more northern parts of the locust country, [164]-[197].	
APPENDIX XIV:	
Answers to the question: Did any locusts remain in the temporary region after the departure of the return swarms? Answers from Arkansas, Colorado, and Iowa, [198]—Answers from Kansas, [199]—Answers from Missouri, [200]—Answers from Minnesota, Nebraska, and Texas, [201].	[198]
APPENDIX XV:	
Prevailing direction in which the young insects travel	[202]
APPENDIX XVI:	
Time of year when the bulk of the insects become winged	[205]
APPENDIX XVII:	#10.04##
Nocturnal habits of locusts in Iowa, Dakota, and Nebraska, [207]— In Minnesota and elsewhere, [208]–[210].	[207]
APPENDIX XVIII:	
Facts relating to the eggs Data as to time of hatching of eggs, proportion destroyed, causes of destruction, etc., in Colorade and Dakota, [211]—In Iowa and Kansas, [212]—In Minnesota, [213]—In Missouri and Nebraska, [216]—In Texas and Utah, [217].	[211]
APPENDIX XIX:	
Means of destruction	[218]
Means employed in Dakota and Nebraska, [218]—In Iowa, [219]—In Minnesota, [220]—Elsewhere, [221].	
APPENDIX XX:	
Injury: crops affected	[222]
APPENDIX XXI: Previous visitations—Aid of animals	[226]
Data from Nebraska and Iowa, [226]—From Minnesota, [227]—From Dakota, [229].	
APPENDIX XXII:	
Minnesota data	[230]
APPENDIX XXIII:	
Are the eggs ever laid thickly for two consecutive years in the same ground?	FORET
Answers to the above question from Arkansas, Colorado, and Iowa, [236]—From Kansas, [237]—From Missouri, [238]—From Minne-	[200]
sota, Nebraska, and Texas, [239].	
APPENDIX XXIV: Miscellaneous data	[240]
From Minnesota and Nebraska, [240]—From Iowa, [241]—From Dakota, [242].	[240]

1643.	RILEY, C. V.—Continued.
	APPENDIX XXV:
	Data from Dakota, Montana, Utah, and New Mexico [243]
	From Dakota, [243]-[247]-From Montana, 248-From Utah,
	[253]—From New Mexico, [259].
	APPENDIX XXVI:
	List of correspondents
	In Arkansas, British America, and California, [261]—In Colorado
	and Dakota, [262]—In Idaho and Iowa, [263]—In Kansas, [264]—
	In Minnesota, [265]—In Missouri, [267]—In Montana, [269]—
	In Nebraska, Nevada, and New Mexico, [270]—In Texas, Utah,
	and Washington Territory, [271]—In Wyoming, [272].
	APPENDIX XXVII:
	Bibliography on the locusts of America. By B. P. Mann [273]
	Index
	Errata[295]
1644.	RILEY, C. V. Attractive but untrue. <n. 31="" july,<="" th="" tribune,="" y.=""></n.>
đ(h	1878. Sb. No. 19, p. 198.
() ()	

Absurdity of statement that "a parasite of the strawberry plant has been found to wage a war of extermination against the Phylloxera."

1645. RILEY, C. V. Locusts eat the castor bean. <N. Y. Tribune, 14 August, 1878. S.-b. No. 19, pp. 198, 227-228.

Acrididæ relish Fagopyrum and Linum but eat Ricinus communis with reluctance, though with impunity.

1646. [RILEY, C. V.] The stalk-borer. <N. Y. Weekly Tribune, 21 August, 1878, v. 37. S.-b. No. 19, pp. 227; 242-243; 244.

Description, food-plants, and habits of larva and description of image of Gortyna nitcla; G. nebris a variety of G. nitela.

1647. RILEY, C. V. That hundred and fifty million dollars. Sci. Amer., 24 August, 1878 [v. 53], n. s., v. 39, p. 117. S.-b. No. 19, p. 245.

Critical review of sensational reports in current agricultural papers; an appropriation of \$5,000 made to the U.S. Department of Agriculture for cotton insect investigation.

1648. RILEY, C. V. Silk-worm breeding. <Sci. Amer., 24 August, 1878 [v. 53], n. s., v. 39, p. 119. S. b. No. 19, p. 244.

Number of annual generations of Sericaria mori; inferiority of races breeding more than once annually.

1649. RILEY, C. V. Cotton-worm. < Daily Constitution [Atlanta, Ga.], 8 September, 1878, v. 11, No. 73, p. 1.

Aletia argillacea [=xylina] feeds from the extra-floral nectar-glands of the cotton-plant.

1650. RILEY, C. V. That "fatherless and motherless race." The basket-worm, alias drop-worm, alias bag-worm, Thyridopteryx ephemeræformis. <Sci. Amer. Suppl., 28 September, 1878, fig. S.-b. No. 19, pp. 245-246.

Critical review of W. H. Gibson's article extracts from author's "The bagworm," with additions; degradation and breeding habits of female Psychida; method of imagination, fecundation, and oviposition of Thyridop-

- 1650. RILEY, C. V.—Continued.
 - teryx ephemeræformis; description of eggs and figures of all other stages; down in which the eggs are enveloped is extruded from the abdomen with the eggs; food-plants, parasites, and means against the species; its follicles available for silk production; this species proterogynous but most insects proterandrous.
- on the structure and development of *Hornia*. < Kansas City Review of Science and Industry, September, 1878, vol. 2, No. 6, p. 353. S.-b. No. 19, p. 209. Reprint: < Proc. Amer. Assoc. Adv. Sci. for 1878 [14 July], 1879, v. 27, pp. 284–285. See: < Ca. Ent. September [October], 1878, v. 10, pp. 177–178.
 - Occurrence in *Epicauta* and *Macrobasis* of the hypermetamorphoses characteristic of *Meloida*; specification of the stages of the same; summary of the life-history of *Hornia minutipennis* parasitic on *Anthophora abrupta*.
- 1652. RILEY, C. V. On the larval characteristics of Corydalus and Chauliodes, and on the development of Corydalus cornutus.

 Kansas City Review of Science and Industry, September, 1878, v. 2, No. 6, p. 354. S.-b. No. 19, pp. 210-211. Reprint:
 Ca. Ent., May, 1879, v. 11, pp. 96-98.
 Proc. Amer. Assoc. Adv. Sci. for 1878, [14 July], 1879, v. 27, pp. 285-287. See:
 Ca. Ent., September [October], 1878, v. 10, p. 178.
 - Characters and economic value of larva of Corydalus cornutus; characters in brief of the imago; characters of larva of Chauliodes; description of eggs of both genera; situation of egg-masses of Corydalus and physical character of their covering; description of respiratory apparatus and method of respiration of larvæ of Corydalus.

Method of hibernation of gall-making Pemphigina on Ulmus.

- NOTE.—Nos. 1651–1653 and 1668—1669 were reprinted from Proc. Amer. Assoc. Adv. Sci., v. 27, under a separate cover. Salem, February, 1879.
- 1654. [RILEY, C. V.] The cottonwood killer. <N. Y. Weekly Tribune, 9 October, 1878. S.-b. No. 19, p. 242. Extract: <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 159-160.
 - Habits, ravages, and food-plants of *Lina scripta*; description of and means against the same; *Trypeta pomonella* an example of the acquisition of new habits in insects.
- 1655. [RILEY, C. V.] A new insect foe to green corn. <N. Y. Tribune,
 9 October, 1878. S.-b. No. 19, pp. 234; 244.
 Ravages of Cetonia [= Euphoria] inda.
- 1656. [RILEY, C. V.] Inquiring friends. <N. Y. Tribune, 9 October, 1878. S.-b. No. 19, p. 245.

Ravages and food-plants of Hylesinus opaculus; impracticability of means against bark-borers in large trees.

- 1657. RILEY, C. V. Notes from the South. Facts about the cotton-worm. <Sci. Amer., 16 November, 1878 [v. 53], n. s., v. 39, pp. 312–313. S.-b. No. 19, pp. 213–214; 239–241. See: <Amer. Rural Home, 27 September, 1878. S.-b. No. 23, p. 128.
 - Note of a trip in the southern United States in 1878; commission to investigate insects injurious to the cotton-plant; food-plants and habits of the larva and image of Aletia argillacea [=xylina]; use of poisonous baits and of Paris green; Anomis exacta injurious in certain regions.
- 1658. RILEY, C. V. Some further facts regarding that "fatherless race." <Sci. Amer. Suppl., 30 November, 1878. S.-b. No. 19, pp. 238-239.
 - Description and figures of male copulatory organs of Thyridopteryx ephemeræformis; description of method of copulation; figures larva, pupa, imagos,
 and follicles of the same; "parthenogenesis, though not improbable, seldom takes place in this species;" nature of the down intermingled with
 the eggs; previous error concerning this down.
- 1659. [RILEY, C. V.] "The carpet bug." <N. Y. Tribune, 1 December, 1878. S.-b. No. 19, p. 236.
 - Ravages of and means against Anthrenus scrophulariæ; figures larva, pupa and imago.
- 1660. [RILEY, C. V.] A bug that eats bees. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237. Reprint: <Prairie Farmer, 4 January, 1879, v. 50, p. 3. S.-b. No. 23, p. 110.
 - Description and habits of Phymata erosa.
- 1661. [RILEY, C. V.] Tomato worm. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237.
 - Excessive abundance of larva of Sphinx quinquemaculata [= Protoparce celeus] near Port Hope, Ontario, in 1878.
- 1662. [RILEY, C. V.] Worm snake. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237.
 - Habits and abode of larvæ of Sciara sp. at Orange, Conn.
- 1663. [RILEY, C. V.] Carpet pests. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237.
 - Food habits, spread of, and means against Anthrenus scrophulariæ.
- 1664. RILEY, C. V. Corn worm. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, pp. 235; 237.
 - Food-plants and means against Heliothis armigera.
- 1665. [RILEY, C. V.] Hessian-fly. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, pp. 235; 237.
 - Indications of the ravages of and food-plants of Cecidomyia destructor.
- 1666. [RILEY, C. V.] Apple-worm. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, pp. 235; 237.
 - Baldwin apples more exempt than other varieties from the attacks of the second broad of Carpocapsa pomonella.
- 1667. RILEY, C. V. Anent the English sparrow. <Evening Star [Washington, D. C.], 28 December, 1878, v. 52, No. 8030, p. 1. S.-b. No. 19, pp. 214–215; No. 23, p. 151.
 - Value of Passer domesticus as a destroyer of insects and in other respects; objections to the special protection of the same.

- 1668. RILEY, C. V. Silk-culture; a new source of wealth to the United States. <Kansas City Review of Science and Industry, 1878, v. 2, pp. 419–423. S.-b. No. 19, pp. 216–218. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1878, [14 July], 1879, v. 27, pp. 277–283. Extract: <Kansas Farmer, 1879: 26 February, 12 November. S.-b. No. 23, pp. 105–106; No. 39, pp. 91–92. See: <Ca. Ent., September [October], 1878, v. 10, p. 178. <St. Louis Globe-Democrat, 28 August, 1878. S.-b. No. 19, p. 203.
 - Practicability and desirability of the extensive establishment of silk-culture in the United States; sketch of efforts made towards such establishment; requisites to its success.
- Mountain locust. <Kansas City Review of Science and Industry, 1878, v. 2, pp. 424–427. S.-b. No. 19, pp. 218–220. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1878, 14 July, 1879, v. 27, pp. 271–277. See: <St. Louis Daily Globe-Democrat, 24 August, 1878, v. 4, No. 95.
 - Limits of breeding-grounds of and region invaded by Caloptenus spretus; causes, periods, and directions of flights of the same; causes of the limitation of the flights; ravages; generalizations regarding locust invasions; work of the U. S. Entomological Commission.
- 1670. RILEY, C. V. A complete life-history of the army-worm, *Leucania unipuncta*, and its parasites. <25th Ann. Rept. Secr. Mass. Board Agric. for 1877, 1878, pp. 243–253, figs.
 - Early history, synonymy, distribution, seasons, enemies, and parasites of Leucania unipuncta; habits of larvæ and of female imagos; descriptions of all stages; figures of all stages except the egg; figures of two unnamed parasitic Ichneumons. Parasitic habits of Exorista [= Nemoræa] leucaniæ, E. flavicauda, Microgaster [= Apanteles] militaris, Pezomachus minimus, and Ophion purgatus; descriptions of Microgaster [= A.] militaris, Mesochorus vitreus, and Ophion purgatum. Ichneumon leucaniæ also a parasite.
- 1671. RILEY, C. V. Phylloxera. < Johnson's New Universal Cyclopædia, 1878, v. 3, pp. 1241-1243, 8 figs. S.-b. No. 10, pp. 192-194.

 Definition of the genus *Phylloxera*; list of the sixteen (16) described United States species; descriptions of their galls; distribution, history, description, and figures of various stages and ravages of and means against *Phylloxera vastatrix*.
- 1672. RILEY, C. V. Potato-bug. <Johnson's New Universal Cyclopædia, 1878, v. 3, pp. 1361-1364, 9 figs. S.-b. No. 10, pp. 195-198. List of principal enemies of potato-plant; spread, natural history, food-plants, enemies of, and means against Doryphora decemlineata; figures of the same in its different stages, and of Lydella [= Exorista] doryphora, Calosoma calidum, Hippodamia convergens, Mysia [= Anatis] 15-punctata, Arma [= Podisus] spinosus, Harpactor [= Milyas] cinctus, Perillus circumcinctus, and Doryphora juncta.
- 1673. RILEY, C. V. Weevil. < Johnson's New Universal Cyclopædia, 1878, v. 4, p. 1338, fig. S.-b. No. 14, p. 46.
 - Definition of "weevil;" mention of numerous species, with statement of their food-plants and manner of obtaining food; habits and means against Sitophilus = Calandragranaria and S. = C.granaria.

- 1674. RILEY, C. V. The locust swarms that devastate the trans-Mississippi country; their source, movements, and eastern limit. <Western Farmer's Almanac for 1879, 1878, pp. 48-50. S.-b. No. 19, pp. 247-248.
 - Limits of the breeding-grounds of and of regions invaded by *Caloptenus spretus*; causes, periods, and directions of their flights; causes of the limitations of the same physical and principally atmospheric; amount of ravages.
- 1675. [RILEY, C. V.] Prof. Riley in favor of the birds. <Evening Star [Washington, D. C.], 9 January, 1879, v. 53, No. 8039, p. 3. S. b. No. 19, pp. 223–224; No. 23, p. 94.
 - From St. Louis [Mo.] Globe-Democrat. List of some birds which are beneficial to horticulture and some which are noxious.
- 1676. RILEY, C. V. Letter from Prof. C. V. Riley. <Colman's Rural World, 15 January, 1879. S.-b. No. 23, pp. 107–108.
 - Letter to N. J. Colman transmitting report as former treasurer of the Missouri State Horticultural Society; transmission of documents for distribution; proposal for a revised and condensed edition of author's reports as State entomologist of Missouri.
- 1677. [RILEY, C. V.] Michigan apples and codling-moth. <N. Y. Tribune, 15 January, 1879. S. b. No. 23, p. 109.
 - Freedom of apples from *Carpocapsa pomonella* in 1878 in Michigan due to the smallness of the apple crop in 1877 and to the efforts made by the orchardists to exterminate the insects.
- 1678. RILEY, C. V. Notes on the Aphididæ of the United States, with descriptions of species occurring west of the Mississippi. <Bull. U. S. Geol. and Geog. Surv. Terr., 28 February, 1879, v. 5, pp. 1–32, pl. 1–2, with 2 pp. expl. of pl. Separate: <Washington, 22 January, 1879, 32 pp., 2 pl., with 2 pp. expl. of pl. Abstract: <Sci. News, 15 April, 1879, v. 1, pp. 184–186.
 - Part 1, pp. 1-17, by C. V. Riley, entitled "Biological notes on the *Pemphiginæ*, with descriptions of new species;" history and description of *Schizoneura americana* n. sp. [p. 4], Colopha ulmicola, Pemphigus populi-monilis n. sp. [p. 13], P. populi-transversus n. sp. [p. 15], P. p.-ramulorum n. sp. [p. 16], P. acerifolii n. sp. [p. 16], P. fraxinifolii n. sp. [p. 17], and Hormaphis spinosus.
 - Part 2, pp. 18-32, by J. Monell, entitled "Notes on Aphidina, with descriptions of new species." Describes nine (9) new species of Siphonophora, five (5) of Aphis, two (2) of Rhopalosiphum, nine (9) of Callipterus, three (3) of Chaitophorus; description of other species of these genera and of Drepanosiphum and the genera above named; synoptic tables of some of the species.
- 1679. [RILEY, C. V.] . [Plant-lice on potatoes.] <N. Y. Tribune, 12 February, 1879. S.-b. No. 23, p. 109.
 - Aphididæ reputed injurious to potato plants, probably Aphis solani.
- 1680. RILEY, C. V. Missouri entomological reports. < Colman's Rural World, 19 February, 1879. S.-b. No. 23, p. 102.
 - Proposal for republication of the reports of the State entomologist of Missouri; reasons for the non-publication of a tenth report; sketch of what that report would have contained.

- 1681. RILEY, C. V. Entomological notes. The chinch-bug. <Farmer's Review. February, 1879, 2 figs. S.-b. No. 23, pp. 111-112.
 - Figures of all stages, ravages, natural history, and means against *Blissus* leucopterus; its method of hibernation, and the effect of weather upon it.
- 1682. RILEY, C. V. Notes on the apple-worm. <Colman's Rural World, 5 March, 1879. S.-b. No. 23, p. 102. Reprint with slight changes: <Amer. Nat., August [July], 1879, v. 13, pp. 523–524. <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 238–239.

See No. 1721 for synopsis of contents.

- 1683. RILEY, C. V. The rice-weevil. Serious injury to stored and to cribbed corn. <Farmer's Review, March, 1879, fig. S.-b. No. 23, p. 99; 112.
 - Figures larva, pupa, and imago of Calandra oryzæ and imago of C. granaria; habits, ravages, and means against the same; description of egg and method of oviposition of C. oryzæ; use of carbon bisnlphide against these insects; poisonous effects of eating the comminuted beetles.
- 1684. RILEY, C. V. [Letter to W. G. Le Duc.] < Boston Daily Herald, 4 April, 1879. < Psyche Advertiser, 11 April, 1879, v. 2, p. 9. < Ent. Nach., 1 July, 1879, jahrg. 5, p. 177.
 - The reason for which the author resigned as entomologist of the U.S. Department of Agriculture was not ill-health.
- 1685. RILEY, C. V. Preventing rot in plums. <N. Y. Tribune, 9 April, 1879. S.-b. No. 23, p. 110; 129.
 - Trapping and jarring recommended as a means against Conotrachelus nenuphar.
- 1686. [RILEY, C. V.] A new insect pest. < Colorado Farmer, 10 April, 1879, v. 12, No. 15, p. 6. S.-b. No. 23, p. 108.
 - Injnry to orchards and ornamental trees in California by *Dorthesia* [characias?], introduced from Australia; ravages of *D. characias* in southern Africa; prospective ravages in California; means against it.
- 1687. RILEY, C. V. The nervous system and salivary glands of Phylloxera. <Psyche, 11 April, 1879, v. 2, pp. 225–226.
 - Correctness of E. L. Mark's conclusion that the author had mistaken nerve cords for tracheæ; dissent from M. Cornu's view that the root swellings cansed by *Phylloxera* are due to mechanical action of the puncture and the subsequent absorption of liquids; swelling considered to be caused by the introduction of a salivary secretion.
- 1688. RILEY, C. V. Mr. Henderson's experiments. < Gardener's Mo. and Hortic., April, 1879, v. 21, pp. 120–121. S.-b. No. 23, p. 107.
 - Review of P. Henderson's "Carnivorous plants;" believes that Dionwa and Drosera are nourished by the insects digested by their leaves.
- 1689. RILEY, C. V. The migrations and hibernation of *Aletia argillacea*. <Washington [D. C.] World, 10 May, 1879. <Science News, 1 June, 1879, v. 1, pp. 230–232. S.-b. No. 23, pp. 119–120. <Sci. Amer., 14 June, 1879. [v. 54], n. s., v. 40, p. 375. <Galveston [Tex.] Daily News, 24 October, 1879, v. 38, p. 2.

- 1689. RILEY, C. V.—Continued.
 - S.-b. No. 23, pp. 166–167, No. 42, pp. 38–39. Abstract: <Amer. Nat., November [25 October], 1879, v. 13, p. 726. <Farmer's Review, September, 1879. S.-b. No. 23, p. 108.
 - Ravages of insects, particularly of Aletia argillacea [=xylina] in the United States; hibernation theories; author's belief in the probable hibernation of the moth in the southern parts of the cotton belt; species mistaken for Aletia; description and food-plants of Aspila virescens.
- 1690. RILEY, C. V. Insects affecting clover. <N. Y. Tribune, 14 May, 1879. S.-b. No. 23, p. 90; 97; 129.
 - Habits, ravages of and means against Hylesinus trifolii and Cecidomyia leguminicola; prior observations upon the same; characters distinguishing C. leguminicola from C. [= Diplosis] tritici and C. destructor.
- 1691. RILEY, C. V. [*Œcanthus niveus.*] <N. Y. Tribune, 14 May, 1879. S.-b. No. 23, p. 95; 98.
 - Occurrence of eggs of *Ecanthus niveus* in twigs of apple, raspberry, grape, and other plants; slight injuries to be expected from the same; means against them.
- 1692. [RILEY, C. V.] Insect powders and their use. <N. Y. Tribune, 14 May, 1879. S.-b. No. 23, p. 95; 96; 130–131.
 - Abstract of W. Saunders' "Insect powder," with additions; differences between Persian and Dalmatian insect powder; methods of using them against insects.
- 1693. [RILEY, C. V.] The grasshopper prospect. <N. Y. Tribune, 14 May, 1879. S.-b. No. 23, p. 98.
 - Prospect of injuries by Caloptenus spretus in 1879; a new species injurious in the Sierra Valley of California.
- 1694. RILEY, C. V. The westward progress of the imported cabbageworm. <Amer. Nat., June [20 May], 1879, v. 13, p. 393. See: <Colorado Farmer, 10 April, 1879. S.-b. No. 23, p. 108. <Farmer's Review, April, 1879. S.-b. No. 22, p. 64. <Colman's Rural World, 1879. S.-b. No. 23, p. 108. <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, p. 239. Spread of *Pieris rapæ* into Illinois and Missouri; means against the same.
- 1695. [RILEY, C. V.] Lures for moths. <N. Y. Tribune, 28 May, 1879. S.-b. No. 23, p. 95.
 - Trapping Carpocapsa pomonella in shallow dishes of sweetened liquid more harmful than beneficial, as many useful insects are destroyed; the method serviceable against Heliothis armigera.
- 1696. [RILEY, C. V.] The currant-worm. <N. Y. Tribune, 11 June, 1879. S. b. No. 23, p. 106.
 - Description, habits, distribution, and spread of and means against *Nematus* ventricosus [=ribesii]; methods of using hellebore.
- 1697. [RILEY, C. V.] Sweet-potato beetle. <N. Y. Tribune, 11 June, 1879. S.-b. No. 23, pp. 106—107; 131.
 - Description of all stages, habits, food plants, and means against *Coptocycla* [= Cassida] nigripes.

- 1698. RILEY, C. V. [The seventeen-year Cicada.] <N. Y. Tribune, 1879. Reprint: <Colman's Rural World, 25 June, 1879. S.-b. No. 23, p. 152.
 - Boundaries of the areas in which Cicada [= Tibicen] septendecim is expected to occur in 1879; request for information of its appearance.
- 1699. RILEY, C. V. The imported carpet-beetle, Anthrenus scrophularia, L. <Farmer's Review, 1879, fig. S.-b. No. 23, pp. 96–98. Extract: <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 54.

Letter of J. H. Parsons, with answer; description of larva and image of *Anthrenus scrophularia*; its history, habits, ravages, and means against them; figures larva, pupa, and image.

- 1700. [RILEY, C. V.] The cheese-skipper. <N. Y. Tribune, 2 July, 1879. S.-b. No. 23, p. 93. Notice: <Farmer's Review, 7 September, 1879. S.-b. No. 23, p. 103.
 - Description of larva, pupa, and imago, habits and means against *Piophila* casei.
- 1701. RILEY, C. V. The thick-thighed walking-stick. <Sci. Amer., 5
 July, 1879 [v. 55], n. s., v. 41, pp. 7-8, fig. S.-b. No. 23, p. 121.
 Reprint, with slight changes: <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 241-245, pl. 3.
 Treats of Diapheromera femorata. See No. 1721 for synopsis of contents.
- 1702. RILEY, C. V. The cotton-worm. < Mobile Register, 9 July, 1879. S.-b. No. 68, p. 204. See: < Colorado Citizen, 17 July, 1879. S.-b. No. 23, p. 91. < Galveston News, 19 July, 1879. S.-b. No. 23, p. 93.
 - Remarks at the Cotton Exchange about prospective injuries from Aletia argillacea [=xylina]; sudden appearance and means against the same; more injurious in wet than dry weather.
- 1703. RILEY, C. V. Other insects affecting cheese. <N. Y. Tribune, 9 July, 1879. S.-b. No. 23, p. 92. Reprint: <Amer. Dairyman. <Western Rural, 9 August, 1879, v. 17, No. 32, p. 250. S.-b. No. 23, p. 92.
 - Habits and transformations of Tyroglyphus siro, Corynetes [=Necrobia] rufipes, and Dermestes lardarius; tenacity of life of the Tyroglyphus and assumption of Hypopus form by the same; anecdote of Latreille; injury done in cheese by Musca corvina.
- 1704. RILEY, C. V. Pupation of the Nymphalidæ. Abstract: <Psyche, 11 July, 1879, v. 2, pp. 249-251.
 - Notice of prevalent explanations of the manner in which suspended pupa of Lepidoptera support themselves while shedding the larval skin; explanation of the process in *Vanessa antiopa* and *Paphia glycerium*.
- 1705. RILEY, C. V. Fire-flies. <Sci. Amer., 26 July, 1879 [v. 55], n. s., v. 41, p. 49. S.-b. No. 23, p. 104.
 - Definition of "Fire-flies;" description of larva, pupa, and image of *Photinus* pyralis; Photuris pennsylvanicus, the commonest species in the more northern states; absence or imperfect development of wings in certain female Lampyridae.

- 1706. RILEY, C. V. Grape-scale insect, new species. < Pacific Rural Press, 16 August, 1879. S.-b. No. 32, p. 66.

 Answer to inquiry of editor of Press; a new coccid, allied to *Pulvinaria vitis*.
- 1707. RILEY, C. V. Dragon-flies. <Sci. Amer., 23 August, 1879 [v. 55], n. s., v. 41, p. 113. S.-b. No. 23, p. 100.

 Description and natural history of Libellulidæ.
- 1708. RILEY, C. V. Failure of tea roses. Habits of Fuller's rose beetle, Aramigus fulleri Horn. <Sci. Amer., 30 August, 1879 [v. 55], n. s., v. 41, p. 129, fig. S.-b. No. 23, pp. 114–116. Reprint: <Gardener's Mo. and Hortic., October, 1879, v. 21, pp. 310–311, fig. S.-b. No. 23, pp. 126–127. Reprint, with additions: <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 255–257, pl. 17, f. 2.

 See No. 1721 for synopsis of contents.
- 1709. RILEY, C. V. The "Devil's darning needle." <Sci. Amer., 6 September, 1879 [v. 55], n. s., v. 41, p. 148. S.-b. No. 23, p. 104. Review: <Sci. Amer., 27 September, 1879 [v. 55], n. s., v. 41, p. 194. S.-b. No. 23, p. 103½.
 - Diapheromera femorata winters in the egg-state; W. J. McGee confounds the above with a species of Ranatra; the term "Devil's darning needle" mostly associated in the popular mind with the dragon-flies (Libellulida).
- 1710. [RILEY, C. V.] The red-spider on roses. <Sci. Amer., 13 September, 1879 [v. 55], n. s., v. 41, p. 161. S.-b. No. 23, pp. 116–117.
 - Nature, ravages, and means against *Tetranychus telarius*; habits and transformations of *Acarina*.
- 1711. RILEY, C. V. Philosophy of the pupation of butterflies. <Sci. Amer. Suppl. No. 193, 13 September, 1879, p. 3069, 3 figs. S.-b. No. 23, p. 125. Reprint: <Science News, 15 September, 1879, v. 1, pp. 346-350. Reprint with changes: <Nature, 16 October, 1879, v. 20, pp. 594-595. <Kosmos, January, 1880, bd. 6, pp. 313-318. <Proc. Amer. Assoc. Adv. Sci. for 1879, 1880, v. 28, pp. 455-463, figs. <Separate: Salem, Mass., July, 1880, 9 pp., 6 figs. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 162-167, figs. 65-70. Abstract: <The Field Nat. and Sci. Student, January, 1883, pp. 179-180. S.-b. No. 42, p. 1. <Bull. Philos. Soc. Wash., 1879, v. 3, pp. 41-43.
 - Illustrates the prevalent explanation of the manner in which suspended chrysalids become attached to their supports; describes and illustrates the more correct process; structure of the terminal joint of the chrysalis especially in Vanessa antiopa, Terias sp., Danais archippus, and Paphia glycerium.
- 1712. RILEY, C. V. Parasites of the cotton-worm. <Ca. Ent., September, 1879, v. 11, pp. 161--162.
 - List of parasites of Aletia argillacea [=xylina]; descriptions of Trichogramma pretiosa n. sp. [p. 161], Cirrospilus [= Tetrastichus] esurus n. sp. [p. 162], and Tachina aletia n. sp. [p. 162].

- 1713. RILEY, C. V. [Insect injurious to junipers.] <Ca. Ent., September, 1879, v. 11, p. 177.

 Junipers on Long Island injured by Dapsilia rutilana.
- 1714. RILEY, C. V. The imported cabbage-worm in the South. <Farmer's Review, September, 1879. S.-b. No. 23, p. 106.

 Spread of *Pieris rapæ* westward and southward; its present distribution; southern limit of distribution of *Doryphora* 10-lineata.
- 1715. RILEY, C. V. The shedding of the tracheæ and double cocoons.

 <Amer. Nat., October, 1879, v. 13, p. 652.

 Review of two notes by E. Potts.
- 1716. RILEY, C. V. Leaf-galls on the grape-vine. <N. Y. Tribune, 1
 October, 1879. S.-b. No. 23, pp. 117; 173. See: <Kansas
 Farmer, 26 November, 1879. S.-b. No. 26, p. 9. <West. Rural,
 10 December, 1879.
 - . Habits, ravages, reproduction, and distribution of Phylloxeva vastatrix.
- 1717. RILEY, C. V. The Croton bug as a library pest. <Library Journal, September-October, 1879, v. 4, p. 376.

 Letter to Weston Flint; ravages of and means against Blatta [=Ectobia] germanica.
- 1718. RILEY, C. V. The Ailanthus silk-worm, *Attacus* (Samia) cynthia. <Science News, 15 October, 1879, v. 1, pp. 377-383. Extract: <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 56-58, fig. 16. <Farmer's Review, 8 January, 1880, v. 4. S.-b. No. 23, pp. 167-168.
 - Description of eggs, larvæ, and imagos of Samia [=Attacus] cynthia; history, food-plants, acclimation, and artificial rearing of the same, and of Samia [=Attacus] riciui; differences between the eggs, larvæ, and cocoons of the two species; availability of these and other species for silk-culture; superiority of Sevicaria mori for this purpose; occurrence of parthénogenesis and of retardation of development in Bombycidæ.
- on some recent cotton-worm. Letter from Prof. C. V. Riley on some recent cotton-worm articles in the News. <Galveston [Tex.] Daily News, 24 October, 1879, v. 38, No. 185, p. 2. S.-b. No. 23, pp. 164–165. Reply: <Ibid., 31 October, 1879, No. 191, p. 4.
 - *Critical review of W. J. Jones's "The cotton caterpillar," and of Investigator's "The cotton-worm;" denies that Aletia argillacea [=xylina] hibernates in the pupa state; author not interested in entomological discoveries for mercenary ends.
- 1720. RILEY, C. V. [Spread of *Pieris rapw* into Alabama.] <Ca. Ent., October, 1879, v. 11, p. 196.
 - Pieris rapæ now quite common in Alabama; it has reached Selma but not Mobile.
- 1721. RILEY, C. V. Report of the entomologist. <Ann. Rept. [U. S.]
 Commissioner Agric. for 1878, 22 November, 1879, pp. 207-257,
 + 1 p. expl. of pl., pls. 1-7. Separate: <Washington, Octo-

1721. F	RILEY, C. V.—Continued.	
	ber, 1879, pp. 52 + 1 p. expl. of pl., 7 pl. Notice: <garden< th=""><th>er's</th></garden<>	er's
	Mo. and Hortic., November, 1879, v. 21, pp. 349-350. Sb.	No
	23, p. 29.	
	LETTER OF SUBMITTAL	207
	INSECTS AFFECTING THE COTTON PLANT	210
	Report of special investigations carried on in 1878, 210—Organization and personnel of the commission of inquiry, 210—Circular letter of inquiries, 210—Report of A. R. Grote, 213—Seasons, hibernation, and migrations of Aletia argillacea [=xylina], 213—Discovery of parasites, 214—Food-plants of the imago, 214.	
r	THE SILK-WORM: A BRIEF MANUAL OF INSTRUCTIONS FOR THE PRO-	045
	[A reprint of special report No. 11.] Culture of Sericaria mori in the United States, 215—Profits of producing cocoons, 217—Profits of reeling, 217—Profits in raising eggs, 218—Nature of the silkworm, 218—The egg, 219—The larva, 219—The cocoon, 220—The chrysalis, 221—The moth, 221—Enemies and diseases, 221—Muscardine, 221—Pébrine, 222—Other diseases, 223—Varieties or races, 223—Wintering and hatching the eggs, 224—Feeding and rearing the worms, 225—Preparation for spinning, 228—Gathering the cocoons, 229—Choking the chrysalis, 229—Egg-laying; reproduction, 230—Reeling, 232—Food-plants, 235—Glossary of terms used, 236.	215
r	The grape Phylloxera	237
	Importance of American grape-culture, 237—Grafting of vines, 237—Distribution of Phylloxera, 238—Hoplessness of means against, 238.	
1	NOTES ON THE APPLE-WORM	238
	Causes of the scarcity of Carpocapsa pomonella in Michigan, 238— Means against the same, 239.	+
T	THE WESTWARD PROGRESS OF THE IMPORTED CABBAGE-WORM Spread of <i>Pieris rapa</i> , 239—Means against the same, 239.	239
(CHAPIN'S APPLE-LEAF SEWER, Phoxopteris nubcculana Clem Habits and natural history, 239—Its injury to apple-trees, 239— Means against it, 239—Description of larva and chrysalis, 239— Synonymy, 239.	239
ם	THE THICK-THIGHED WALKING-STICK, Diapheromera femorata Say Vernacular names, 241—Nomenclature, 241—Characters, 241—Destructive powers, 241—Habits and natural history, 243—Its appearance every alternate year, 244—Natural enemies, 244—Remedies, 245—Description of egg, 245—The larva, 245.	241
7	THE GREAT ELM-LEAF BEETLE, Monocesta coryli, Say*	245
	Past history, 245—Ravages, 246—Habits, 246—The eggs, 246—The larvæ, 246—Habits of larvæ, 246—The pupa, 246—Remedies, 247.	
	THE JUNIPER WEB-WORM, Dapsilia rutilana, Hübn Importation, 247—Distribution in Europe, 247—Habits, 247—Remedies, 248—Description of larva, 248; of pupa, 248; of imago, 248.	247

^{*} Translation: <Acker- und Gartenbau-Zeitung, 1 June, 1880, v. 11, p. 87, fig. S.-b. No. 23, p. —.

17 21.	. RILEY, C. V.—Continued.	
	THE CLOVER-ROOT BORER, Hylesinus trifolii, Müller *	248
	Occurrence in New York, 248—Habits in Europe, 249—Food-plants,	
	249-Natural history, 249-Remedies, 250-Differential charac-	
	ters of H. opaculus and H. trifolii, 250—Enemies, 250—Descrip-	
	tion of larva, 250; of pupa, 250.	
	THE CLOVER-SEED MIDGE, Cecidomyia leguminicola, Lintner	250
	Ravages in New York, 250—Habits, 251—Remedies, 251—Descrip-	
	tion of imago, 251-Larva, 252-Cocoon, 252-Pupal excuvium,	
	252—Comparative characters of C. tritici, C. destructor, and C.	
	leguminicola, 252.	
• 0	THE APPLE COLEOPHORA, Coleophora malivorella, n. sp. †	253
	History, 253—Habits, 253—Injuries, 253—Larval habits, 254—Ene-	
	mies and remedies, 254—Description of larva and imago, 254.	
	Fuller's Rose-Beetle, Aramigus fulleri Horn	255
	History, 255—Distribution, 255—Ravages, 255—Habits and charac-	
	ters of image 256—Remedies 257—Description of egg larva and	

pupa, 257.

1722. RILEY, C. V. [Parasites bred from the cotton-worm.] <Ca. Ent.,
November, 1879, v. 11, p. 205.

Nine species of parasites have been bred from Aletia argillacea [=xylina].

- 1723. RILEY, C. V. The bee-moth. <N. Y. Tribune, 1879. Reprint: <Farmer's Review, 3 January, 1880. S.-b. No. 23, p. 158. Description of larva and imago, habits, ravages, and means against Galleria cereana.
- 1724. RILEY, C. V. The silk-worm; being a brief manual of instructions for the production of silk. <Special Report No. 11 [U. S.], Dept. Agric., Washington, 1879, pp. 31, 8 figs. Second edition: <Washington, 1882, 37 pp., 8 figs. Third edition: <Washington, 1883, 37 pp., 8 figs. Fourth edition: Fifth edition: Sixth edition: <Bull. No. 9 Div. Ent. U. S. Dept. Agric., 11 July, 1886, 65 pp., 29 figs., 2 pl. Seventh edition: <Ibid., April, 1888.

Treats of Sericaria mori. See No. 1721 for synopsis of contents.

- 1725. RILEY, C. V. London purple as an insecticide. <Farmer's Review, 29 January, 1880, v. 4, No. 5, p. —.
- 1726. [RILEY, C. V.] [The food-habits of thrushes.] < Amer. Ent., January, 1880 [v. 3], n. s., v. 1, pp. 2-3.
 - Notice of S. A. Forbes' "The food-habits of thrushes;" incompleteness of our knowledge of the food-habits of birds and of Carabida; author's opinion favorable to the birds.
- 1727. [RILEY, C.V.] The grape Phylloxera in California. <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 3.
 - Phylloxera vastatrix as destructive to Vitis vinifera in California as in Europe; the winged female supposed not to have appeared yet in California; speculations upon this modification of habit, if true.

^{*} Translation: <Acker- und Gartenbau-Zeitung, 1 August, 1880, v. 11, p. 119, 2 figs. S.-b. No. 23, p. —. Extract: <Prairie Farmer, 22 November, 1879, v. 50, p. 370. S.-b. No. 23, pp. 168-169. <Amer. Rural Farmer, 18 December, 1879. S.-b. No. 45, pp. 23-24.

[†] Translation: < Acker- und Gartenbau-Zeitung, 1 September, 1880, v. 11, p. 135, fig. S.-b. No. 23, p. 182.

- 1728. [RILEY, C. V.] On the hibernation of the cotton-worm, *Aletia* argillacea Hübn. <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, pp. 6-11, figs. 3-5; p. 15.
 - Advance print from Bull. No. 3 U. S. Ent. Commission, pp. 24-31. See No. 1736 for synopsis of contents.
- 1729. [RILEY, C. V.] [Inquilines in galleries of the common white-ant.] < Amer. Ent., January, 1880, [v. 3], n. s., v. 1, p. 15.
 - Trichopscnius depressus and three undescribed Aleocharini found inquilinous in galleries of Termes flavipes in Texas by E. A. Schwarz; the Trichopscnius previously known by a single specimen from Georgia.
- 1730. [RILEY, C. V.] Large white scale on acacias, etc. <Amer. Ent. January, 1880 [v. 3], n. s., v. 1, p. 20.
 - Occurrence and ravages of *Dorthesia characias*? in California and southern Africa; characters of the insect.
- 1731. [RILEY, C. V.] [Ravages of moths in cushions.] < Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 20.
 - Conclusions of discussion by German Society of Railway Companies as to the best method of preventing the ravages of moths in the cushions of railway carriages.
- 1732. [RILEY, C.V.] [Grape Phylloxera in Geelong, Victoria.] < Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 20.

 Occurrence of Phylloxera vastatrix in Geelong, Victoria.
- 1733. RILEY, C. V. Borers in black-ash: Fall web-worm: Apple-tree insects. . <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, pp. 22–23.
 - Black-ash trees in New York killed by the borings of Parandra brunnea; ash trees bored also by Trochilium [= Fatua] denudata; the leaves of the same and other trees destroyed by Hyphantria textor [= cunea]; apples injured by the "gimlet-worm"; habits, ravages of, and means against Bucculatrix pomifoliella; figure of cocoons and imago of this moth.
- 1734. [RILEY, C. V.] Skippers injuring smoked hams. < Amer. Ent., January, 1880, [v. 3], n. s., v. 1, pp. 23-24.
 - Answer to communication from E. A. Smith; hams injured by larvæ of *Pio-phila casei*; means against these and other insects injuring hams.
- 1735. [RILEY, C. V.] *Tipula* eggs in the stomach of the cat-bird. <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 24.

 Answer to communication from S. A. Forbes; description of eggs and method

. of oviposition of Tipula trivittata?

1736. RILEY, C. V. The cotton-worm. Summary of its natural history, with an account of its enemies and the best means of controlling it; being a report of progress of the work of the commission. <Bull. No. 3, U. S. Ent. Commission, 28 January, 1880, 144 pp., 1 pl., 84 figs. Extract: <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 67-68, figs. 19-20. <Farmer's Review, January, 1880, 2 figs. S. b. No. 23, pp. 163-164.

Introductory	1
Text of circular sent out by the commission, 1—History of the investigation, 1—Need thereof, 2—Early work of Townend Glover, Thomas Affleck, and D. L. Phares, 2—Special report ordered by Congress, 3—Character of the Bulletin, 3—Division of the cotton-belt in the southern or hibernating and the northern or non-hibernating portions, 4—Practical aim of the commission,	
4—Persons who have assisted, 4.	
1. Destructiveness of the worm Losses occasioned by it Increase of injury westward, 7—Least injury in northern portion of belt, 7—Tabular statement of loss for past 15 years, 8.	7 7
2. Popular and scientific names for the insect Most desirable popular name, 8—Different technical names by which the insect has been known, 8—Earliest description of it, 8.	8
3. Characters and stages of the insect	9
The egg. The worm always hatched from an egg, 9—Where the egg is laid, 9—Number of eggs to a leaf, 10—All eggs perish during winter, 10—Other eggs mistaken for those of Aletia, 10.	9
The worm or larva. Character of newly-hatched worm, 10—Number of molts, 11— Variation in color, 11—Peculiar habits, 11—Flourishes on the cotton-plant only, 12.	10
The chrysalis	12
The moth or imago Distinguishing characters of the moths, 14—Sexual differences, 14—Different habits at night and at day, 14—Fecundity, 15— It finds its natural food on the cotton-plant, 15—Fond of fruits, 15—Character of its tongue, 15.	13
4. Time elapsing from one generation to another	16
5. Time of year when the first worms appear. Prevailing opinions erroneous, 16—The worms hatch in April, 16—First worms always in small numbers, 17—The third generation usually ealled the first, 17.	16
6. Condition of soil and plant connected with the appearance of the first worms. The worms appear first on low, moist, and rich lands, 18—Few exceptions to the rule, 18—The plant must be well advanced and luxuriant, 18—Some shelter generally near, 19—Reoccurrence year after year in the same spots, 19—Theory explaining first appearance under the circumstances observed, 19.	18
7. WET WEATHER FAVORS THE DEVELOPMENT OF THE WORMS Humid and hot atmosphere most favorable, 20—Heavy storms destroy, 20—Effect of dryness, 21—Indirect influences of wet weather, 21—It prevents the working of the worm's natural enemies, 21.	20

1736. RILEY, C. V.—Continued.	
8. Number of annual generations.	21
Prevailing general belief erroneous, 21—Early generations often overlooked, 22—Also often exterminated by natural enemies,	
22—At least seven annual generations in southern portion of	
belt, and probably more, 22.	
9. MIGRATIONS AND POWER OF FLIGHT OF THE MOTH	21
The moth has great power of wing, 23—It flies many hundreds	
of miles north of eotton belt, 23—Appearance of the worm on eotton first planted great distances from any other cotton thus	
accounted for, 23—Migrating habit developed only in later	
month's when they become numerous, 24.	
10. HIBERNATION	24
Different opinions and beliefs, 24—Difficulty surrounding the question, 24—Erroneous conclusions easily drawn, 25—Fallacy	
of the belief that the ehrysalis winters underground, 25-26—	
Ability of the moth to survive the winter, 26—Other moths	
easily mistaken for it, 27—Theory of annual introduction of	
the species from some exotic country, 28—Arguments for and against the theory, 29, 30—Hibernation of the moth in the	
southern portion of the belt substantiated, 30—But a very	
small proportion of those which abound in the fall live to per-	
petuate their kind the next spring, 31—Localities where the	
moth hibernates, 31.	0.1
11. Natural enemies	31
Quadrupeds, birds, and reptiles that feed upon the worm, 32—	00
Introduction of English sparrow, 33.	
Invertebrates	33
Predaeeous insects	33
ants, 34—How they are drawn to a cotton-field by the aphides,	
34—Beetles which prey upon the worm and others which pre-	
sumably do so, 35—Enemies among the half-winged bugs or	
Heteroptera, 36—Among two-winged flics or Diptera, 36—Among nerve-winged insects or Neuroptera, 37.	
Parasites	38
Previous belief that none preyed on Aletia, 38—Infesting the	
egg, 38—Infesting and issuing from the worm, 39—Flesh flies and Tachina flies, differences between them, 40-42—Issuing	
and Tachina flies, differences between them, 40-42.—Issuing from the ehrysalis, 42—A parasite known but not named in	
1852—Ten distinct parasites infesting Aletia, and two other	
undetermined species, 47.	
12. Remedies: Means of coping with the insect	47
South since the war, 48.	
Prevention	48
Modes of cultivation, 48—Topping, 48—Use of carly varieties,	
48—Isolating fields, 48—Diversified agriculture, 49—Rotation of groups, 49—Protection of natural angular 49—Farly as late	
of erops, 49—Protection of natural enemies, 49—Early vs. late planting, 49—Interspersing corn with cotton, 49—Jute, 49.	
Mechanical means of killing the worms	49
Simple methods of brushing and dragging, 50.	F 0
Machines for brushing off the worms	50

1736.	RILEY.	C. V.	—Contin	med.
		O	COHUIL	iccoca.

12. Remedies: Means of Coping with the Insect—Continued. The Ewing brushing machine, 50—The Helm brushing machine, 51—The Wood-Smith brushing machine, 52—The Iske brushing machine, 53—Use of kerosene and coal oil with shallow pans or cloth frames, 54.	
Poisoning the worms. Impetus given to invention by the use of mineral poisons, 54— Value of concerted action, 54—General principles to be observed in poisoning, 55—Dry vs. wet applications, 55—Advantage in being prepared, 55.	54
Arsenical compounds. Their value, 55—Safe and harmless where carefully used, 55—Cantion necessary, 56—Antidote to arsenic poison, 56—Paris green, 56—When first recommended against the cotton-worm, 56—Its value, advantages, and disadvantages, 56—Modes of using it, 57—Patents issued for Paris green mixtures, 57—They should be disregarded, 58—Arsenic, 58—It causes the squares to fall, 58—Patent granted for its use in 1871, 59—Johnson's dead-shot, 59—Arseniate of soda, 59—Texas cotton-worm destroyer, 60—Lodi pest poison, 60.	55
London purple	60
Pyrethrum powder Its origin and nature, 62—Its remarkable insecticide qualities, 62—Most effectual on the cotton-worm, 63—Perfectly harmless to man, 63—The extract of the plant has the same virtne, 63— Growth of the plant in California, 63—Experiments conducted, 64—The alcoholic extract very potent, 64—Probable cost of using it, whether as powder or extract, 65—Highly encouraging prospects from its use, 65.	62
Keroscne oil. It canses instant death to the worm, but is difficult to apply without injuring the plant, 66—Vaporized by means of steam, 66—Used as a soap, 66.	65
Acts much as kerosene, 66—Made into soap with ashes from hull of seed, 66—Might be profitably used around oil factories, 66.	66
Carbolic acid	67
Sulphnr	67
Extracts or decoctions from various plants	67
Yeast ferment; fungns infection. Proposition by Dr. Hagen, 68—Dr. Bail's experiments, 68—Not in accord with the leading mycologists, 68—Experiments with	68
beer-mash, 69—With beer-mash and yeast ferment, 70. Machines and contrivances for powdering	74

2, Remedies: Means of Coping with the Insect—Continued. and sprinkler, 78—The Davis duster, 78—The Levy duster, 79—The Taylor duster, 80—The Allen duster, 81—The Young duster, 81—The Goodheart duster, 81.	
Machines and contrivances for sprinkling and spraying	81
Machines for applying vapors The Steinmann vaporizer, 97—The Pearl vapor generator, 99.	97
Destruction of the moth	100
Lamps for attracting the moth Usually resorted to when foo late to do practical good, i00— Fires, 100—Trouble of keeping them up throughout the night, 101—Beneficial insects killed by lamps and fires, 101—Ad- vantage of lamps, 101—The Garrett lamp, 101—The Binkley lamp, 102—The McQueen lamp, 103—The Rigel lamp, 103— The Walker lamp, 103—Simple form of lamp, 104—The Pugh lamp, 104—Lamps in motion, 105—Le Blanc's cotton-moth destroyer, 105—The Fordtran cotton-moth destroyer, 105. Poisoned sweets and fluids as means of destroying the moth. Sweets attractive, 106—Ripe fruits and melons often suffer, 106— Baits that allure and kill, 106—Experiments with them, 107— Moths less attracted during summer than in autumn, 107—Lit- tle inducement to use baits in autumn, 108—They are most	100
beneficial in spring, 108—White rags, 108—Theory of Dr. J. L. Lupton, 108.	
Answers to circular from West Feliciana Parish, Louisiana, 109— From Evergreen, Ala., 111—From Henderson, Tex., 113—From Hempstead, Tex., 115—Different views as to hibernation and mode of perpetuation of the insect, 116—Auswers to circular from Livingston, Ala., 117—From Parish of Concordia, La., 119—From Natchitoches, La., 122—From Jackson, Miss., 126— From Larissa, Tex., 127—From Perry County, Ala., 128— From Walterborough, S. C., 128—From Jamestown, Fla., 130— From San Antonio, Tex., 131—From Kirkwood, Miss., 133— From Woodville, Miss., 135.	
	1

1737. [RILEY, C. V.] The 17-year Cicada in Iowa. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, pp. 25-26.

Review of C. E. Bessey on the 17-year Cicada in Iowa; limits of the broods of 1854-1871, 1861-1878, and 1862-1879 of Cicada [= Tibicen] septendecem in Iowa; occurrence of the last brood in Missouri; comparison of the distribution of these broods with the distribution of timber trees.

- 1738. [RILEY, C. V.] Vegetal-feeding ground-beetles. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26. Notes from various sources upon the phytophagous habits of Carabidæ.
- 1739. [RILEY, C. V.] The pear-leaf blister. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26. Notice: <Ibid., March, 1880 [v. 3], n. s., v. 1, p. 74.
 - Review of T. J. Burrill's "The pear-leaf blister"; ravages of *Typhlodromus* pyri on leaves of pear-tree; characters of this mite; noxions habits of Acarina, especially of Phytoptus.
- 1740. [RILEY, C. V.] Fuller's rose-beetle in California. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26.

 Occurrence and ravages of Aramigus fulleri in California.
- 1741. [RILEY, C. V.] Lepidium vs. bed-bugs. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26.

 Notice of S. M.'s "Lepidium, the bed-bug destroyer;" Acanthia lectularia said to be attracted to and killed by Lepidium sp.
- 1742. [RILEY, C. V.] Aniseed and grain weevils. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26.

 Notice of reports that Calandra granaria were attracted from grain to anisced and killed by it.
- 1743. RILEY, C. V. Two valuable insecticides. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, pp. 41-45.

 From Bull. No. 3, U. S. Ent. Commission, pp. 60-65. See No. 1736, for synopsis of contents.
- 1744. [RILEY, C. V.] Use of buckwheat to destroy insects. < Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 48.

 Extract from "The Farmer" with remarks on the planting and plowing in of buckwheat as a means against injurious insects.
- 1745. [RILEY, C. V.] Tipula eggs in stomach of cat-bird. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 50.

 Answer to communication from S. A. Forbes; two kinds of eggs found in stomach of eat-bird; these birds eat gravid females of Tipula.
- 1716. [RILEY, C. V.] Beetles supposed to be feeding on wheat. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 50.

 Answer to letter from P. H. M.; larva of Cryptorhopalum sp. supposed to have injured grains of wheat, was probably feeding on the remains of true grain insects; food-habits of larvæ of Dermestidæ.
- 1747. RILEY, C. V. The apple-twig borer. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, pp. 50-51, figs. 11-12.

 Answer to letter from H. G. Woleott; habits, food-plants, and means against Amphicerus bicaudatus; figure of male and female imago and of bored apple-twigs; habits and food-plants of Sinoxylon basillare and of Bostrichus.
- 1748. [RILEY, C. V.] Stinging caterpillars. Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 51.

 Answer to letters from T. Pollard and G.W. S.; food-plants, general appear-

Answer to letters from T. Pollard and G.W. S.; food-plants, general appearance, vernacular name, urticating properties and description of imago of Lagoa opercularis.

- 1749. RILEY, C. V. A new genus of *Proctrotrupidæ*. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 52, figs. 13–14. Review: <1bid., December, 1880 [v. 3], n. s., v. 1, p. 293.
 - Description and figures of Didictyum [=Hexaplasta] zigzag n. g. et n. sp. reared from chrysalis of Aletia argillacea [=xylina].
- 1750. RILEY, C. V. Agricultural advancement in the United States. <Farmer's Review, 1880, v. 4: 4 March, p. 158; 11 March, p. 174. S.-b. No. 23, pp. 153-158. Notice: <*Ibid.*, 4 March, 1880, p.—. S.-b. No. 23, p. 150. Reprint: <Journ. Amer. Agric. Assoc., April, 1881, v. 1, pp. 47-54. Separate of reprint: <August, 1881, pp. 47-54.
 - Suggestions for the organization of an agricultural association; comparison of the relations of government to the advancement of agriculture in England, Germany, France, and the United States; necessary changes in the United States Department of Agriculture.
- 1751. RILEY, C. V. Parasites of the plum Curculio. <Farmer's Review, 4 March, 1880, v. 4, p. —, 2 figs. S.-b. No. 23, pp. 169–170.
 - Figures of larva, pupa, cocoon, and imagos of Sigalphus curculionis; habits, variations, usefulness, and dissemination of the same; habits, colors, and seasons of Porizon [= Thersilochus] conotracheli.
- 1752. [RILEY, C. V.] Trapping the carpet-beetle. < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 53-55, fig. 15.
 - Figures larva, pupa, and imago of Anthrenus scrophularia; food-plants and habits of imago and ravages of larva; importation, distribution, vernacular names of and means against the same.
- 1753. [RILEY, C. V.] Silk-worm eggs: Silk culture. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 55.
 - Method of obtaining eggs of Sericaria mori and information on silk culture.
- 1754. [RILEY, C. V.] [Hickory Scolytus.] < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 58.

 Occurrence of Scolytus quadrispinosus in Washington Territory.
 - 1755. [RILEY, C. V.] [Tenacity of life.] <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 68.

 Resistance of Cleonus sp. to the influence of various insecticides.
 - 1756. [RILEY, C. V.] [Danais archippus.] < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 73.
 - Occurrence of a colorational variety on the island of Antigua.
 - 1757. [RILEY, C. V.] Reports of the U. S. Entomological Commission. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 73.

 Method of obtaining the first report of the U. S. Entomological Commission.
 - 1758. [RILEY, C. V.] [Habits of the cotton-moth.] < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 74.
 - Request for information on the habits of the imago and the food-plants of the larva of Aletia argillacea [=xylina] during March and April.
 - 1759. [RILEY, C. V.] Typhlodromus pyri. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 74.
 Reference to early account of the pear-leaf blister-mite.

- 1760. [RILEY, C. V.] Food-habits of ground-beetles. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 75.
 - Note on request of S. A. Forbes for Carabidæ found in situations suggesting herbivorous habits.
- 1761. [RILEY, C. V.] Moths and butterflies caught by the tongue. Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 75.
 - Notice of several articles on the capture of Noctuida and Sphingida by the flowers of Physianthus albens; capture of Sphingida by the flowers of Nerium oleander and Enothera grandiflora, and of Syrphus sp. by the flowers of Bidens chrysanthemoides.
- 1762. [RILEY, C. V.] [Queen bees in the mails.] Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 75.
 - Circumstances under which queen bees may be sent by mail; exclusion of pinned insects from the same.
- 1763. [RILEY, C. V.] Common tiger-beetle. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 77–78.
 - Answer to inquiry of J. L. Seney; habitat, distribution, and food-habits of Cicindela repanda.
- 1764. [RILEY, C. V.] Gall on *Pelargonium*. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Answer to letter from Mary Treat; gall at base of *Pelargonium* probably made by a mite; the *Poduræ* found in it were undoubtedly feeding on diseased gall-tissue.
- 1765. [RILEY, C. V.] Catalogues and monographs of insects. < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Answer to letter of P. S. B.; mention of monographic works and catalogues of North American insects.
- 1766. [RILEY, C. V.] Leaf-hoppers injuring wheat fields. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Answer to letter of R. L. B.; habits and ravages of Cicadula exitiosa, Diedrocephala flaviceps, and Jassus sp. in the southern United States, and of Jassus sexnotatus in Europe; means against the same.
- 1767. RILEY, C. V. A new leaf-hopper injurious to small grain. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Description of Diedrocephala flaviceps n. sp. injurious to wheat and oats in Texas in 1876.
- 1768. RILEY, C. V. The bird question dispassionately considered. <Farmer's Review, 1 April, 1880, v. 4, p. 211. S.-b. No. 23, pp. 147–148.
 - Extract from E. Perris's "Birds vs. Insects," with introductory; conclusions as to the value of birds in agriculture as destroyers of noxious insects.
- 1769. RILEY, C. V. The cotton-worm in the United States. <Amer. Ent., April, 1880 [v.3], n. s., v.1, pp. 93-95. Reprint, with slight changes: <Proc. Amer. Assoc. Adv. Sci. for 1879, 23 February, 1881, v. 28, pp. 464-466. Separate of reprint: <Salem, Mass., August, 1880, 3 pp.
 - Date and manner of first appearance of larvæ of Aletia argillacea [=xylina]; number of annual generations and the existence of parasites upon it; his-

- 1769 RILEY, C. V.—Continued.
 - tory and refutation of the annual immigration theory; division of the cotton belt into regions wherein, respectively, A. argillacea [=xylina] is permanent and temporary; food-habits and enemies of the same.
- 1770. [RILEY, C. V.] The migrations of butterflies. < Amer. Ent., April, 1880 [v. 3], n. s., v. 1, pp. 100-102, figs. 34-35.
 - Migratory flights of Lepidoptera supposed to be due to excessive multiplication of individuals and to a true migratory instinct; direct ous, times, and extent of migrations of Danais archippus; explanation of the migratory instinct; distribution of D. archippus and Pyrameis cardui; figures the former and its manner of clustering.
- 1771. [RILEY, C. V.] Gouty gall on blackberry and raspberry canes. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 107.
 - Answer to letters of P. S., T. A. C., and J. W.; seasons, ravages of and means against Agrilus ruficollis.
- 1772. [RILEY, C. V.] Sowing cotton seeds in hot-beds and transplanting as a means of preventing injury from the cotton-worm. <Amer. Ent., April, 1883 [v. 3], n. s., v. 1, p. 107.

 Impracticability of W. J. W.'s suggestion as indicated in title.
- 1773. [RILEY, C. V.] Not the cotton-moth. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 107.
 - Answer to inquiry of D. B. Woodbury; food-plants of Tolype velleda.
- 1774. [RILEY, C. V.] Moths caught in Alabama: Muscle-shaped barklouse on apple-trees South <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, pp. 107-108.
 - Mentions several insects received from J. F. Bailey, of Marion, Ala.; larva of Papilio philenor feeds on Aristolochia; Bombus virginicus Q robs hives of Apis mellifica; Mytilaspis pomicorticis [=pomorum] not before received from so far south; means against the same.
- 1775. [RILEY, C. V.] Chrysalides supposed to be those of Aletia. Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 108.
 - Answer to letter of W. J. Jones; pupe of Agrotis inermis [=saucia] and Prodenia lineatella from cotton field.
- 1776. [RILEY, C. V.] Apple-twig borer. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 108.
 - Answer to inquiry of T. V. M.; occurrence and habits of Amphicerus bicaudatus.
- 1777. [RILEY, C. V.] Clover-weevil. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 108.
 - Answer to inquiry of W. W. F.; several "weevils" infest clover-seeds; Hylesinus trifolii infests the roots and lower part of the stem.
- 1778. [RILEY, C. V.] Effects of cold applied to the chrysalides of butterflies. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 110-111.

 Notice of the experiments of W. H. Edwards; effects of hibernation in the chrysalis and imago states.
- 1779. [RILEY, C. V.] Moth issuing from a larva. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 114.
 - Notice of J. J. Weir on the issuance of *Orgyia* sp. from the larva without passing through the pupal stage.

- 1780. [RILEY, C. V.] The rose-slug. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 115-116, figs. 42-43.
 - Description of eggs, larvæ, and imago, history and means against Selandria [=Monostegia] rosæ; figures, egg, larvæ, imago, with details of structure, and injured leaf.
- 1781. [RILEY, C. V.] Dr. Asa Fitch. <Amer. Ent., May, 1880 [v. 3], n. s , v. 1, pp. 121-123.
 - Biographical sketch of Asa Fitch, with notice of his collection of insects and his writings.
- 1782. [RILEY, C. V.] Flea-beetle on young tobacco plants. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 123.
 - Leaves of Nicotiana tabacum eaten by Epitrix hirtipennis [= Crepidodera parvula] and probably by E. [= C.] cucumeris; leaves of Solanum eaten by the first named and by E. [= C.] brevis; distribution of these and other species of Epitrix [= Crepidodera].
- 1783. [RILEY, C. V.] The proboscis of the common house-fly. Amer.
 Ent., May, 1880 [v. 3], n. s., v. 1, p. 125.
 Review of paper by G. Macloskie.
- 1784. [RILEY, C. V.] Notes on South American Lepidoptera. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 125-126.
 - Report of meeting of the Entomological Society of London; length of proboscis of and presence of scent organ in Sphingidæ; perception of colors in Rhopalocera; secondary sexual character in Callidryas and other genera; according to R. Meldola the proboscis of Macrosila cluentius is 23.5cm (9½ inches) long.
- 1785. [RILEY, C. V.] Floating apiaries. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 126-127.
 - Notice of an unsuccessful attempt to increase the product of an apiary by floating it southward late in the season and back to the north in the spring, on the Mississippi River.
- 1786. [RILEY, C. V.] American *Staphylinida* wanted. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 127.
 - Notice of A. Fauvel's works on *Staphylinida* and his request for additional material.
- 1787. [RILEY, C. V.] Probable parthenogenesis in the Hessian fly. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 127.
 - Review of paper by H. A. Hagen; Cecidomyia destructor less injurious than formerly in the older wheat-growing regions, its area of abundance having moved westward with the westward extension of wheat culture.
- 1788. [RILEY, C. V.] Raspberries destroyed by weevils. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 127.
 - Ravages of Otiorhynchus picipes in England; warning against its introduction into the United States.
- 1788a. [RILEY, C. V.] Cotton culture and the insects affecting the plant at Bahia, Brazil. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 128-129.
 - Letter of R. A. Edes, with remarks on specimens sent.

1789. [RILEY, C. V.] Hemispherical larva at bottom of ant hill. < Amer. Ent., May, 1880 | v. 3], n. s., v. 1, p. 129.

Answer to inquiry of D. S. Sheldon; habitat of larva of Microdon globosus.

1790. [RILEY, Č. V.] Chrysalides dug up in cotton field, mistaken for those of the cotton-worm. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 129.

Answer to letter of R. Worrel.

1791. [RILEY, C. V.] Aleurodes on Oxalis. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 129-130.

Answer to letter of S. A. Conrad; description of all stages of Aleurodes sp. from leaves of Oxalis sp.

1792. [RILEY, C. V.] Larvæ in stomach of black-bass. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 130.

Answer to inquiry of S. A. Forbes; two coleopterous larvæ from the stomach of *Micropterus salmoides*; one a dytiscid, the other perhaps a dascyllid.

1793. [RILEY, C. V.] Insects from stomach of rock-bass. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 130.

Answer to letter of S. A. Forbes; coleopterous larva from the stomach of common sun-fish (*Lepiopomus pallidus*) and rock-bass (*Roccus lineatus*).

1794. [RILEY, C. V.] New enemy to sugar-cane. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 130, fig. 48.

Answer to letter of D. Th.; ravages of Ligyrus rugiceps upon sugar-cane, maize, and grasses; recommends use of lamp and kerosene pan, and figures one form of such apparatus.

1795. [RILEY, C. V.] Parasites of the plum Curculio. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 131-132, figs. 49-50.

Answer to letter of W. S. Barnard; quotes from 3d Ann. Rept. State Ent. Mo., pp. 24-26; habits of Sigalphus curculionis; habits, varieties, and figures of all stages of the same; food-habits of Semasia [= Grapholitha] prunivora on which the Sigalphus is parasitic.

1796. [RILEY, C. V.] Blister-beetles from New Mexico. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 132.

Answer to inquiry of J. M.; Macrobasis albida common in the Southwest and valuable as a vesicant; Diplotaxis not known to be a vesicant.

1797. [RILEY, C. V.] Abnormal cocoon. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 132.

Answer to letter of W. S. Barnard; occurrence of Callosamia [= Attacus] promethea and Orgyia antiqua at Ithaca, N. Y.; description of abnormal cocoon of the latter found on sugar-maple.

- 1798. [RILEY, C. V.] Insects found about orange-trees. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 132.
 - Answer to inquiry of J. S. Barnwell; list of insects from Darien, Ga.; most of them have no relation to orange trees; notes on Aphis sp., Chilocorus bivulnerus, Syrphus sp., Neoclytus erythrocephalus, Drasterius amabilis, Platynus punctiformis, Forficula sp., Gryllotalpa borealis, and Psocus venosus.

- 1799. [RILEY, C. V.] Luperus brunneus, Crotch. <Amer. Ent., May, 1880, [v. 3], n. s., v. 1, p. 132.
 - Luperus noxius Riley MSS., is a pale form of L. brunneus; description of the typical and other forms of the same; distinctive characters of L. brunneus and L. morulus from other North American species and from each other; characters of the genus Calomicrus.
- 1800. [RILEY, C. V.] Professor Riley on army-worms. <N. Y. Weekly Sun, 20 June, 1880. S.-b. No. 26, p. 89.
 - Interview with reporter; summary of present knowledge of Leucania unipuncta.
- 1801. RILEY, C. V. Cotton-caterpillars. <Selma [Ala.] Times, 25 June, 1880.
- 1802. RILEY, C. V. Notes on our commoner insects. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 133, 134, figs. 51-52.
 - Description and figures of larva, pnpa, cocoon, and image of Arctia [= Pyrrharctia] isabella; habits, seasons, food-plants, parasites, vernacular names and variations of the same; figure of Ophion macrurum; descriptions of Ichneumon carulcus, I. signatipes, and Trogus obsidianator; description of larva of Ecpantheria scribonia.
- 1803. RILEY, C. V. The white-grub fungus. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 137-140, figs. 53-55.
 - Description and figures of *Torrubia ravenelii* and of its fructification; forms in which it occurs on larvæ of *Lachnosterna quercina* [= fusca]; list of some articles upon it; its synonymy and distribution; list of related species and their distribution; occurrence of related species on other insects.
- 1804. RILEY, C. V. The true and the bogus Yucca moth, with remarks on the pollination of Yucca. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 141–145.
 - History of observations on *Prodoxus decipiens*; habits, parasite, and foodplants of the same; evolution of *Prodoxus* and *Pronuba*; criticism of errors resulting from mistaking *Prodoxus decipiens* for *Pronuba yuccasella*; failure of *Yucca angustifolia* to become fertilized in the absence of *Pronuba yuccasella*; *Hyponomeuta quinquepunctella* Chambers is the same as *Prodoxus decipiens*.
- 1805. [RILEY, C.V.] Intermittance of phosphorescence in fire-flies. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 146.
 - 'Criticism of a discussion before the London Entomological Society; the winged imagos of Lampyrida can intermit their light; the larva and wingless imagos can suppress their light.
- 1806. [RILEY, C. V.] Grain Aphis vs. rust. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 147.
 - Ravages of Aphis avenæ [Nectarophora granaria] and of rust on wheat and oats in Georgia; the work of aphides promotes the growth of some kinds of rust.
- 1807. [RILEY, C. V.] Mold and Phylloxera. <Amer. Ent.; June, 1880 [v. 3], n. s., v. 1, p. 147.
 - Criticism of statements by Rommier; development of mycelinm on phyllox-erized roots does not result in the destruction of the *Phylloxera*.

- 1808. [RILEY, C.V.] Infecting Phylloxera with fungus disease. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - Abstract of discussion at meeting of Académie des Sciences de Paris on the infection of *Phylloxera vastatrix* by parasitic fungi; particular fungi infect only particular insects; method of experimentation to determine which fungi to use and how to apply them; improbability of success in these experiments.
- 1809. [RILEY, C. V.] Fungus in Cicada. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - Notice of C. H. Peck's description and J. Leidy's note on Massospora cicadina; Cicada [= Tibicen] septendecim and C. [= T.] tredecim subject to the attacks of the same.
- 1810. [RILEY, C. V.] On the nature of the phosphorescence of the glowworm. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - Conclusions drawn by Jousset de Bellesme; probability that the phosphorescent substance is a gascous product; nature of phosphorescence.
- 1811. [RILEY, C. V.] Death of mules caused by insects. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - About 6,000 mules reported killed in the Ouachita Valley, Louisiana, by the attacks of Simulium sp.
- 1812. [RILEY, C. V.] [May beetles swarming in Alabama.] < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - Abundance and ravages of Lachnosterna quercina [=fusca] on oak-trees around Mobile, Ala., in May, 1880.
- 1813. [RILEY, C. V.] Fungus diseases of beneficial insects. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 149.
 - Abstract of communication by C. Brongniart and Max Cornu on an epizootic among Syrphus mellinus caused by a species of Entomophthora; almost all insecticides liable to the objection that they kill useful as well as noxious insects.
- 1814. [Riley, C. V.] Early appearance of cotton-worm. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 149.
 - Extracts from letter of J. M. Bell and from the Goliad [Tex.] Guard and from G. Witting on the date of appearance of Aletia argillacea [=xylina] in Texas.
- 1815. [RILEY, C. V.] The butterfly tongue. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 149.
 - Notice of and extract from paper by E. Burgess; conclusions as to the manner in which butterflies imbibe their liquid food.
- 1816. [RILEY, C. V.] Cottony maple scale. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 149.
 - Notice of J. D. Putnam on Pulvinaria innumerabilis.
- 1817. [RILEY, C. V.] Necrological. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 150.
 - Obituary notices of E. A. H. v. Kiesenwetter, S. C. Snellen van Vollenhoven, and F. L. de Laporte.
- 1818. [RILEY, C. V.] Effects of severe cold on insects. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 150.

A steady even if severe winter not prejudicial to insect life.

- 1819. [RILEY, C. V.] Revision of the *Lampyridæ*. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 150.
 - Notice of H. S. Gorham's work on the Lampyride and his request for additional material.
- 1820. [RILEY, C. V.] Apple-twig borer. Amer. Ent., June, 1880 | v. 3|, n. s., v. 1, p. 151.
 - Auswer to inquiry of A. S. H.; occurrence of Amphicerus bicaudatus on appletrees in Virginia.
- 1821. [RILEY, C. V.] Grape-vine flea-beetle. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 152–153.
 - Answer to letter of J. Nilis; ravages of Graptodera [= Haltica] chalybea on grape-vines in Pennsylvania.
- 1822. [RILEY, C. V.] Cynipid gall on oak twigs. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 153, fig. 56.
 - Answer to letter of J. A. Warder; figure of gall of *Cynips* (u. sp.?) from twig of *Quercus prinus* var. acuminata; comparison of this gall with that of C. [=Andricus] q.-punctata; the flies bred from these galls issue in spring and are all females, but probably have a bisexual form producing a different gall.
- 1823. [RILEY, C. V.] White-grub fungus. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 153.

 Answer to letter of H. S.; occurrence of Torrubia ravenelii at Iola, Kans.
- 1824. [RILEY, C. V.] Galerita janus. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 153, fig. 57.
 - Answer to inquiry of H. D. M. Fair; distribution, habitat, transformations, and seasons of Galerita janus; figure of larva of G. lecontei.
- 1825. [RILEY, C. V.] Not Fuller's rose-beetle. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 153-154, fig. 58.
 - Answer to inquiry of J. Stewart; habits and synonymy of Agonoderus comma [=pallipes]; figure of the same.
- 1826. [RILEY, C. V.] Not *Aletia* chrysalides. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 154.
 - Answer to letter of G. P. White; pupe of Agrotis saucia from cotton fields; food-habits of larva; ravages of Aletia argillacea [=xylina] at Brown Station, Ala., in August, 1879.
- 1827. [RILEY, C. V.] Mud-wasp and parasite. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 154, figs. 59-60.
 - Answer to inquiries of Mary Treat; nidification of several species of Odynerus; food-habits of O. birenimaculatus; figure of O. flavipes and of a nest of Odynerus sp.; figure of Cryptus junceus.
- 1828. [RILEY, C. V.] Monographs again. <Amer. Ent., June, 1880, [v. 3], n. s., v. 1, pp. 154-155.
 - Answer to letter of C. D. Marsh; references to works. Elaterida, Curculionida and Coccida.
- 1829. [RILEY, C. V.] Rearing wood-borers. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 155.
 - Answer to letter of W. H. Harrington; directions for rearing larvæ which bore in wood or under bark,

- 1830. RILEY, C. V. On a new tineid genus allied to *Pronuba*, Riley. <Amer. Ent., June, 1880, [v. 3], n. s., v. 1; pp. 155–156.
 - Description of *Prodoxus* n. g. [p. 155] of *Tineina* and of larva, pupa, and imagos of *P. decipiens* n. sp. [p. 155].
- 1831. RILEY, C. V. A parasite on *Prodoxus decipiens*. <Amer. Ent., June, 1880, [v. 3], n. s., v. 1, p. 156.

Description of larva and imago of Exothecus prodoxi n. sp.; habits of larva.

1832. RILEY, C. V. How to manage the cotton-worm: Suggestions to cotton planters. <Farmer's Review, 8 July, 1880. S.-b. No. 24, p. 68.

Means against Aletia xylina.

- 1833. [RILEY, C. V.] The cotton-worm investigation. <Selma [Ala.]

 Morning Times, 21 July, 1880, v. 55, No. 190, p. 3. Reprint:

 <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 197.
 - List of persons engaged in the cotton-worm investigation; statement of work to be done by each.
- 1834. [RILEY, C. V.] A foe to cottonwood. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 159–161, figs. 61–64. Extract: <Suppl. to Amer. Ent., July, 1880, p. 1.
 - Description of egg, larva, and image and figures of all stages of Melasoma [=Lina] scripta; description and figures of variations of the image; habits, ravages, seasons, food-plants, and means against the same; figure of M. [=L.] lapponica and of larva of M. [=L.] populi; description of these larvæ and of that of M. [=L.] tremula; food-plants and distribution of these species; acquisition of new habits by insects.
- 1835. [RILEY, C. V.] The northern army-worm. < Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 170–171, figs. 72–75.

Ravages of Leucania unipuncta in 1880; natural history, means against, description, and figures of all stages of the same.

- 1836. [RILEY, C. V.] The periodical Cicada. <Amer. Ent.. July, 1880 [v. 3], n. s., v. 1, pp. 172–173, fig. 76.
 - Broods of Cicada [= Tibicen] septendecim and C. [= T.] tredecim which appear in 1880.
- 1837. [RILEY, C. V.] Use of guano for grape Phylloxera. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 173. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 2.

Application of sulphide of carbon with infusorial earth or guano.

- 1838. [RILEY, C. V.] Fertilizers of alpine flowers. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 175. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 2.
 - Relative frequency, according to H. Müller, of the visits of insects to flowers in high alpine regions as compared with such visits at lower levels.
- 1839. [RILEY, C. V.] Carnivorous habits of caddis-worms. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 176.
 Notice of paper by G. C. Goody.

- 1840. [RILEY, C. V.] Development of the eyes and luminosity in the fire-flies. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 176.

 Abstract of H. S. Gorham's observations.
- 1841. [RILEY, C. V.] Grape Phylloxera not at the Cape. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 176. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 1.

Examination of roots of unhealthy vines by R. McLachlan and R. Trimen show that the disease of the vines is not caused by *Phylloxera vastatrix*.

- 1842. [RILEY, C. V.] [Bill providing for the extermination of insects.]

 <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 176–177.

 Text of and comments on a bill introduced into the California Assembly, providing for the extermination of insects.
- 1843. [RILEY, C. V.] [Appropriation for the U. S. Entomological Commission.] <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 177.

 Notice of the appropriation by Congress of \$25,000 for completing the work of the U. S. Entomological Commission.
- 1844. [RILEY, C. V.] Pronuba vs. Prodoxus. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 177-178.

 Comments on letter of V. T. Chambers.
- 1845. [RILEY, C. V.] Interesting cotton-worm notes from Vera Cruz, Mexico. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 179. Reprint with slight omission: <Suppl. to Amer. Ent., July, 1880, p. 3.
 - Letters from S. T. Trowbridge and R. de Zayas Enriquez, with notes on the irregular occurrence of Aletia argillacea [=xylina] in Vera Cruz.
- 1846. [Riley, C. V.] Clover-root borer. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 179-180, fig. 81. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 4. See: <Prairie Farmer, 31 July, 1880. S.-b. No. 45, p. 15.

Answer to letter of W. A. Henry; description and figures of larva, pupa, and imago of *Hylesinus trifolii*; ravages, habits, and means against the same.

- 1847. [RILEY, C. V.] Mud-wasp and spider egg-nest. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 180, figs. 82-84.
 - Answer to letter of Mary Treat; figures of cells and of imago of Eumenes fraterna, which stores its cells with larvae of Paleacrita vernata; figure and descriptions of egg-nests of Epeira sp.
- 1848. [RILEY, C. V.] Worm in joints of wheat. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 180–181, fig. 85. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 4.
 - Answer to letter of J. K. P. Wallace; ravages of an unknown, apparently hymenopterous, larva in wheat stalks; figure of larva and pupa of *Meromyza americana* and of stalks injured by the same.
- 1849. [RILEY, C. V.] Linden and ash destroyers. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 4.

- 1849. RILEY, C. V.—Continued.
 - Answer to letter of Shelby Reed; season of appearance of Odontota rubra on Tilia; larvæ of Hyphantria textor [= cunea] feeding on the same; Saperda calcarata boring in poplar, and an unknown larva (Neoclytus capræa?) boring in black-ash.
- 1850. [RILEY, C. V.] Larva boring along the axis of apple-twigs. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181.
 - Answer to letter of T. V. Munson; twigs of apple-trees bored by larvæ of Oberea sp.?
- 1851. [RILEY, C. V.] Aquatic larvæ. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181.
 - Answer to letter of S. A. Forbes; larvæ of Anax junius and Palingenia [= Hexagenia] bilineata named; the latter common in the stomach of fishes.
- 1852. [RILEY, C. V.] First appearance of cotton-worm in prairie belt. Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181. Reprint: <Suppl. to Amer. Ent., July, 1880, pp. 3-4.
 - Answer to letter of J. F. Bailey; larvæ of Aletia argillacea [=xylina] appear first on lands where the cotton is luxuriant.
- 1853. [RILEY, C. V.] Gyrinus larva; terrestrial insects in stomach of shad. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 182.
 - Answer to letter of S. A. Forbes; character of larva of *Gyrinus*; a mass of terrestrial insects including *Typhlocyba vitis*? a muscid, a Eurytomid, *Jassus* sp., *Triphleps insidiosus*, and other species from the stomach of Ohio shad.
- 1854. RILEY, C. V. Further remarks on the differences between *Pronuba* and *Prodoxus*. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 182.
 - Differences between Pronuba yuccasella and Prodoxus decipiens in the form, sculpture, and color of the terminal joint, and of the ovipositor.
- 1855. [Riley, C. V.] The grape-vine flea-beetle, *Graptodera chalybea* Illig. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 183–184, fig. 86.
 - Habits, seasons, ravages of, and means against Graptodera [= Haltica] chalybea; figures, injured leaf, and larvæ, cocoon, and imago of the same.
- 1856. RILEY, C. V. Further notes and observations on the armyworm. <Amer. Ent., 1880 [v. 3], n. s., v. 1: August, pp. 184, 185; September, pp. 214, 215. Reprint, with changes: <Sci. Amer., 4 September [v. 57], n. s., v. 43, p. 152. S.-b. No. 23, pp. 161–162.
 - Progress of knowledge upon the number of annual generations of *Leucania* unipuncta; stages of growth at which it hibernates; the destructive generation probably not the first of the season; explanation of the partial efficacy of the burning of fields in winter as a means against this insect; connection of wet and dry seasons with its increase; its natural habits; errors of A. Fitch.
- 1857. [RILEY, C. V.] Sprinklers and atomizers. <Amer. Ent., 1880 [v. 3], n. s., v. 1: August, pp. 185–189, figs. 87–98; September, pp. 211–214, figs. 111–117.
 - Extracts from pp. 56-57 and reprint of pp. 85-94 of Bull. No. 3, U.S. Entomological Commission. See No. 1736 for synopsis of contents.

- 1858. RILEY, C. V. The use of pyrethrum. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 193-195.
 - Record of experiments by W. A. Henry upon the effect of Pyrethrum powder on Halticidæ, Meloidæ, Pieris rapæ, Coreus [=Anasa] tristis, and Blattidæ; and of the fumes of burning pyrethrum on various insects.
- 1859. [Riley, C. V.] [Colorado potato-beetle in New Hampshire.] Amer.Ent., August, 1880 [v. 3], n. s., v. 1, p. 195. Ravages of Doryphora 10-lineata in Coos County, N. H., in 1879.
- 1860. [RILEY, C.V.] Retarded development in a blister-beetle. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 196.

 Final transformations of individuals from a single batch of eggs of *Epicauta vittata* occurring at the first, second, and third year after hatching.
- 1861. [RILEY, C. V.] .Ox-eye daisy as an insecticide. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 196.

 Experiments by W. S. Barnard upon the effect of alcoholic extracts of the flowers and stems of the ox-eye daisy on insects; "no evidence that they will prove of any practical value."
- 1862. [RILEY, C.V.] Directions for raising pyrethrum. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 197.
 Directions for sowing the seeds and cultivating the plants of pyrethrum.
- 1863. [RILEY, C. V.] State Entomologist for New York. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 197, 198.

 Notice of the appointment of J. A. Lintner as State Entomologist of New York.
- 1864. [RILEY, C. V.] Economic investigations in the South and West. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 198.

 Meeting of the U. S. Entomological Commission; plan of work to be done in 1880-1881; partial list of persons engaged in the work.
- 1865. [RILEY, C. V.] [Number of entomologists in Europe.] < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 198.

 Tabulation of the number of entomologists of the several countries of Europe.
- 1866. [RILEY, C. V.] [Catalogus coleopterorum by Gemminger and Harold.] <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 199. Commendation of the above work and of the supplementary list of *Elateridæ* by E. Candèze; need of co-operation for the completion of the whole catalogue.
- 1867. [RILEY, C.V.] Carnivorous propensity of plant-feeders. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 200.

 Larvæ of Plusia brassicæ devour those of Pieris rapæ and Pionea rimosalis in default of more natural food.
- 1868. [RILEY, C. V.] Beetles injuring cabbages and fuchsias. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 200, fig. 106.

 Answer to letter of G. T.; ravages, distribution, and figures of Epicarus imbricatus; description of eggs, ravages, food-plants, and means against Graptodera [= Haltica] carinata.
- 1869. [RILEY, C. V.] Spider and nest. < Amer. Ent., August, 1880, [v. 3], n. s., v. 1, p. 200.

 Approximate letter of Mrs. I. B. Harrison: distribution and distinctive charges.

Answer to letter of Mrs. J. B. Harrison; distribution and distinctive characters of Acrosoma stellatum,

- 1870. [RILEY, C. V.] Damage to wheat: Worm boring in the stalk. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 201.
 - Answer to letter of A. R. Frost; food-plants and ravages of larva and figure of larva and imago of Gortyna nitela.
- 1871. [RILEY, C. V.] Larvæ from stomach of blue-bird. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 201.
 - Answer to letter of S. A. Forbes; larva of Meracantha contracta and of Callimorpha sp. from stomach of blue-bird; habits and characters of the same.
- 1872. [RILEY, C. V.] Butterfly larva injurious to cotton squares. Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 201.
 - Answer to letter of B. F. Cooke; larva of *Thecla* (pæas?) feeding on leaves and bolls of cotton-plant; a species of *Microgaster* parasitic in the larva.
- 1873. [RILEY, C. V.] Cut-worms from stomach of robin. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 201.
 - Answer to letter of S. A. Forbes; larvæ of Agrotis (messoria?) from stomach of robin; Agrotis cochrani = A. messoria.
- 1874. [RILEY, C. V.] Large phosphorescent larva. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 201–202, fig. 108.
 - Answer to letter of S. F. Clarke; occurrence in Maryland and Missouri of a luminous larva (Mclanactes sp.?) [= Phengodes sp.]; figures of the larva, of its head and leg enlarged and of its probable parent; occurrence in the more northern States of a similar larva, probably that of Asaphes memnonius.
- 1875. RILEY. C. V. Worms injuring wheat. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 202.
 - Answer to letter of J. Monaghan; ravages of larva of Meromyza americana.
- 1876. RILEY, C.V. Ash-root borer: Supposed eggs of Odontota. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 202-203.
 - Answer to letter of Shelby Reed; habits and ravages of *Parandra brunnea*; *Pimpla* sp. parasitic upon its larva; probable manner in which the eggs of *Odontota rubra* are laid.
- 1877. [RILEY, C. V.] Army-worm notes and inquiries: Its work on clover. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 203. Answer to letter of L. T. Derousse; Lencania unipuncta frequently leaves clover-plants untouched while eating the grasses growing with the clover, but under some circumstances destroys young clover-plants.
- 1878. [RILEY, C. V.] Ichneumon from stomach of bluebird. <Δmer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 203.
 - Answer to letter of S. A. Forbes; Lampronota sp. eaten in quantity by bluebirds.
- 1879. [RILEY, C. V.] Leaf-miner on white oak. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 203.
 - Answer to letter of E. W. Claypole; description, habits, and ravages of Lithocolletis cincinnatiella.
- 1880. [RILEY, C. V.] Screw-worm: Its parentage in doubt. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 203.
 - Answer to letter of A. R. 'Kilpatrick; Lucilia macellaria supposed to be the parent of the "screw-worm"; food-habits of certain Muscide.

- 1881. [RILEY, C. V.] Silk culture: How to dispose of cocoons. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 203-204.
 - Answer to letter of C. F. Durksen; need of silk-filatures in the United States; present means of selling cocoons.
- 1882. [RILEY, C. V.] Best cotton-worm destroyer. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 204.
 - Answer to letter of J. G. Dauterive; reference to the most reliable poison and the best machines and methods for applying poison to kill Aletia argillacea [=xylina].
- 1883. [RILEY, C. V.] Twice-stabbed lady-bird. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 204, fig. 109.
 - Answer to letter of H. N. Patterson; Coccinellidae beneficial by feeding on Aphididae and Coccidae; figure of Chilocorus bivulnerus; significance of the presence of this beetle in large numbers on trees.
- 1884. [RILEY, C. V.] Pseudo-scorpion. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 204.
 - Answer to inquiry of C. H. S. Davis; habitats and food-habits of Chelifer cancroides and other Chernetide.
- 1885. [RILEY, C. V.] Bluebirds feeding on parasitic and predaceous insects. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 204–205.
 - Answer to letter of S. A. Forbes; bluebirds more destructive to parasitic and predaceous insects than the thrushes; percentage of such food found in the stomachs of the bluebird; larvæ of Leucania unipuncta, Telephorus bilineatus, and Nephelodes violans eaten by the same; probable hibernation of the first named; habits, hibernation, distribution, and popular name of the Nephelodes.
- 1886. RILEY, C. V. The cotton destroyers. <New Orleans Democrat, 21
 September, 1880, v. 5, No. 276, p. 8. S. b. No. 23, pp. 170-173.
 Reprint: <Selma [Ala.] Times, 29 September, 1880. <Southern Enterprise [Atlanta, Ga.], December, 1880, v. 5, pp. 77-82.
 S.-b. No. 23, pp. 184-189; No. 61, pp. 3-5; No. 63, pp. 57-59; 60-62. Reprint, with slight changes: <Proc. Amer. Assoc.
 Adv. Sci. for 1880, 14 October, 1881, v. 29, pp. 642-649. Separate: <Salem, July, 1881, pp. 26-33. Extract: <Sci. Amer., 16
 October, 1880 [v. 57], n. s., v. 43, p. 241. S.-b. No. 23, pp. 159-160. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, pp. 245-247. <Prairie Farmer, 30 October, 1880, v. 51, No. 44, p. 2. S.-b. No. 23, pp. 153. Abstract: <Farmer's Review, 7 October, 1880. S.-b. No. 23, pp. 162-163. <Journ. Appl. Sci., November, 1880, v. 11, pp. 170-171.
 - Facts and principles established by the U. S. Entomological Commission, applicable to the whole cotton belt, regarding the times and manner of first appearance of Aletia argillacea [=xylina] and Heliothis armigera upon the cotton-plant; habits of the larvæ and imagos and the best means of destroying the same; relative efficacy of several insecticides; methods of preparing and applying the same; importance of early poisoning; main object of the cotton-worm inquiry accomplished.

- 1887. [Riley, C. V.] The use of poisons to destroy insects. <Farmer's Review, 23 September, 1880, v. 5, p. 200. S.-b. No. 23, pp. 148-149; 165-166. Reprint: <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 244.
 - Advocacy of the use of Paris green and London purple as means against certain insects; danger of the use of the same against many insects; criticism of A. J. Cook's recommendation for their use against *Paria aterrima* and *Carpocapsa pomonella*; limitations within which they may be used.
- 1888. [Riley, C. V.] Supplementary instructions to agents of the United States Entomological Commission. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 218.
 - Call for experiments to ascertain the minimum quantity of poisons which can be used effectually against Aletia argillacea [=xylina]; effect of poisons on the several stages of this insect; food-plants, parasites, and enemies of the same; effect of yeast ferment upon it.
- 1889. RILEY, C. V. Dimorphism in locusts (Acrididæ). <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 219-220.
 - Review of paper by S. H. Scudder; certain forms described as species of *Pezotettix* are dimorphic forms of *Caloptenus*; extract from 8th Ann. Rept. State Ent. Mo., 1876, p. 115.
- 1890. [RILEY, C. V.] A scale insect on maple hitherto unobserved by American entomologists. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 220–221.
 - Notice of paper by Miss E. A. Smith, with emended reprint of a portion of the same; habits, colors, and behavior of young larvæ and transformations of the male of *Pseudococcus aceris*.
- 1891. [RILEY, C. V.] How flight in insects is directed. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 221.
 - Abstract of paper by Jousset de Bellesme; direction of flight not determined by the motion of the wings, but principally by the displacement of the center of gravity, resulting from the changes of position of other parts of the body.
- 1892. [RILEY, C. V.] Entomological work at the Department of Agriculture. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 222.

 Notice of the continuation of the appropriations by Congress for field-work and experiments in the entomological division of the U. S. Department of Agriculture.
- 1893. [RILEY, C. V.] Entomological papers read before the A. A. A. S. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 223.

 Titles of twenty-eight [28] papers read at the Boston meeting.
- 1894. [RILEY, C. V.] Entomologists at Boston. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 223.

 Notice of the 29th meeting of the A. A. A. S. at Boston, August-September,
- 1895. RILEY, C. V. Winged Phylloxera in California. Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 224–225.

1880.

Letter from J. S. Hyde, with remarks; extracts from papers by E. W. Hilgard; winged fertile females of *Phylloxera vastatrix* found in California; local evidence of the spread of this insect; means of eradicating the same.

- 1896. [RILEY, C. V.] Worm infesting meal sacks. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 229.
 - Answer to letter of J. Greenwood, jr.; description of larva and imago of Ephestia zeæ [=interpunctella]; food-habits of the larva.
- 1897. [RILEY, C. V.] Hesperid larva feeding on Canna. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 229.
 - Answer to letter of N. S. Reed; occurrence of Pamphila ethlius in Illinois and South Carolina; its larva injurious to Canna flaccida.
- 1898. [RILEY, C. V.] Grape-vine apple-gall. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 229, fig. 119.
 - Answer to letter of A. R. McCutchen; occurrence of *Cecidomyia* ? vitis-pomum and of *Calosoma scrutator* in Georgia; figures gall of the first named; foodhabits of the *Calosoma*.
- 1899. [Riley, C. V.] Apple-tree plant-lice in Oregon. Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 229-230.
 - Answer to letter of H. B. May; occurrence and ravages of Aphis mali? in Oregon; means against the same.
- 1900. [RILEY, C. V.] Phylloxera work. Wood-lice on grape-vine roots. Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 230.
 - Answer to letter of H. B. Trimble; occurrence of *Phylloxera vastatrix* at West Chester, Pa.; roots of grape-vines destroyed from unknown cause, perhaps by young of *Phyloscia* or *Porcellio*.
- 1901. RILEY, C. V. New hickory galls made by Phylloxera. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 230.
 - Description of *Phylloxera caryw-scissa* n. sp. and *P. caryw-avellana* n. sp. from Florida; larva of *Diplosis* found in the galls.
- 1902. RILEY, C. V. Food-habits of the longicorn beetles or wood-borers. <Amer. Ent., 1880 [v. 3], n. s., v. 1; October, pp. 237–239; November, pp. 270–271.
 - Food-plants and food-habits of the Prionida and Cerambycida of the United States.
- 1903. [RILEY, C. V.] Additional experiments with Pyrethrum. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 242.
 - Details of experiments made by H. G. Hubbard on the effect of very minute quantities of Pyrethrum powder on very young larvæ of Aleta argillacea [=xylina].
- 1904. [RILEY, C. V.] A new enemy to the strawberry. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, pp. 242-243, fig. 121.
 - Review of paper by A. J. Cook; extracts from the same; description and habits of larva and image of *Paria aterrima*; means against the image; habits of larva of *Colaspis flavida*; figure of the same; number of segments and spiracles in larvæ of Coleoptera.
- 1905. [RILEY, C. V.] A new enemy to corn: The long-horned *Diabrotica*. Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 247.
 - Notice of recent articles on the ravages of the larva of Diabrotica longicornis; food-habits and means against the same; food-habits of Drasterius amabilis.
- 1906. [RILEY, C. V.] Phylloxera congress in Spain. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 247.
 - Notice of a congress to be held in Spain to consider all topics connected with the ravages of *Phylloxera vastatrix*.

- 1907. [RILEY, C. V.] The grape Phylloxera not permanently destructive. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.
 - Recovery of grape-vines formerly injured by *Phylloxera vastatrix* in California and Missouri; belief that in Europe grape-vines will be grown again on the lands of late years ravaged.
- 1908. [RILEY, C. V.] Sale of silk-worm eggs. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.
 - \$6,000,000 worth of silk-worm eggs sent from Japan to France via San Francisco in each of the four years 1874-1877.
- 1909. [RILEY, C. V.] [Death of S. S. Haldeman.] < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.

 Obituary notice.
- 1910. [RILEY, C. V.] [Retirement of Mr. Fuller.] < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.
 - Announcement of the retirement of A. S. Fuller as assistant editor of the American Entomologist.
- 1911. [RILEY, C. V.] Insect enemies of growing rice. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 253.
 - Answer to letters of J. L. Leconte and J. Screven; ravages of and means against the "grub" [== Chalepus trachypygus] and the "maggot" [== Lissorhoptrus simplex] in rice fields in Georgia.
- 1912. [RILEY, C. V.] Blind-eyed *Smerinthus*. < Amer. Ent., October 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of Mrs. J. B. Harrison; occurrence at lamp-light in N. H. of Smerinthus excacatus; food-plant of the larva of the same.
- 1913. [RILEY, C. V.] White waxy secretion on stems of bitter-sweet. Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of J. A. Lintner; method of oviposition and characters of egg-mass of *Enchophyllum* [= *Enchenopa*] *binotata*; seasons, habits, and food-plants and description of the eggs of the same.
- 1914. [RILEY, C. V.] Prickly-ash larva: Tachinid eggs. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of R. W. Jones; larva of Papilio cresphontes on prickly ash; eggs of Tachina sp. on larva of Aletia argillacea [=xylina].
- 1915. [RILEY, C. V.] Worms on cabbage: Boll-worm feeding on leaf. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of R. W. Jones; food-habits of larva of Heliothis armigera and of Pionea rimosalis.
- 1916. [RILEY, C. V.] Buggy peas. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to letter of A. Berger; abundance of Bruchus pisi in Wisconsin; means against it.
- 1917. RILEY, C. V. Glow-worm. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254, fig. 123.
 - Answer to letter of J. J. Dean; occurrence of larva of *Photuris pennsylvanica* at Chatham, N. J.; figures larva and imago of the same: larva and imago of *Photinus pyralis*, also luminous; males and females of these species winged; female of *Lampyris noctiluca* wingless and more luminous than the male.

- 1918. [RILEY, C. V.] The use of fungus growths to destroy insects.
 Amer. Ent., November, 1880 [v. 3], n. s., v. 1, pp. 269–270.
 - Abstract of paper by A. N. Prentiss; the constant presence of spores of fungi in the air vitiates the value of most experiments on the application of the fungi to insects; insects rarely affected by the spores which are in the air while the plants are more so affected; yeast more injurious mechanically than infectiously.
- 1919. [RILEY, C. V.] New species of scale insects. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, pp. 275-276.

 Review of paper by W. H. Ashmead on the red scale.
- 1920. [RILEY, C. V.] Remedy for cabbage worms. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 276.

 Pyrethrum powder the most satisfactory means against all larvæ affecting

the leaves of cabbage.

- 1921. [RILEY, C. V.] Pyrethrum for the screw-worm. < Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 276.
 - Notice of paper by A. R. Kirkpatrick; directions for the use of pyrethrum powder against the "screw-worm" [=Lucilia macellaria]; objections to the use of other remedies.
- 1922. [RILEY, C. V.] Oviposition in the *Tortricidæ*. <Amer. Ent. November, 1880 [v. 3], n. s., v. 1, p. 276.

 Notice of and extract from paper by C. H. Fernald.
- 1923. [RILEY, C. V.] About *Phora* being merely a scavenger and not a true parasite. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 277.

Letter from C. R. Osten Sacken, with remarks; Phora aletia not a parasite; oviposition and habits of larva of the same.

- 1924. [RILEY, C. V.] Gall on Solidago leaves. < Amer. Ent., November, 1880 [v. 3], n. s.; v. 1, p. 278.
 - Answer to letter of H. Barnes; occurrence of galls of Cecidomyia carbonifera on leaves of Solidago nemoralis? at Mulberry Corners, Ohio.
- 1925. [RILEY, C. V.] Oak gall: Cynips q.-decidua Bass. < Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 278.
 - Answer to letter of J. Schenck; galls of Cynips q. decidua? found on leaves of Quercus muhlenbergii at Mount Carmel, Ill.
- 1926. [RILEY, C. V.] Insects from stomach of lark, robin, and sunfish. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 278.
 - Answer to letter of S. A. Forbes; Diplotaxis sordida from stomach of meadow-lark; egg of a reduviid from that of a robin; larva of an ephemerid (Polymitarcys alba?) from the stomach of a sunfish.
- 1927. [RILEY, C. V.] Supposed hibernating Aletia chrysalis. < Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 278.
 - Answer to letter of J. W. Davidson; pupa resembling that of Achatodes zew found in stalk of maize at Uniontown, Ala.
- 1928. RILEY, C. V. On the natural history of certain bee-flies (Bomby-liidae). <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp.

- 1928. RILEY, C. V.—Continued. 279–283, figs. 147–151. Review: <Ent. Mo. Mag., February, 1881, v. 17, pp. 206–207.
 - Advance print of pp. 262-267 and a general abstract of pp. 267-269 of the section entitled "Bee-fly larvæ, family Bombyliidæ," in 2d Rept. U. S. Entomological Commission. See No. 1959 for synopsis of contents.
- 1929. RILEY, C. V. On a new pyralid infesting the seed-pods of the trumpet-vine. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp. 286-288, figs. 152-153.
 - Description, with figures, of Clydonopteron n. g. [p. 287] and of larva, pupa, and image of C. tecomæ n. sp. [p. 288], and of the abode of this insect in pods of the trumpet-vine; habits of the insect.
- 1930. [RILEY, C. V.] Experiments with yeast-ferment on various insects. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp. 289, 290.
 - Reports by J. E. Willet and A. J. Cook upon their experiments in the application of beer and yeast to various insects; no infection communicated to the insects by these applications.
- 1931. RILEY, C. V. Notes on the imported elm leaf-beetle. < Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp. 291-292.
 - Answer to letter of J. L. Leconte; seasons, transformations, enemies, and ravages of and means against Galeruca xanthomelæna; hibernation of Chrysomeliāæ.
- 1932. [RILEY, C. V.] Synonyms of parasites' mistakes corrected. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 293. Review: <Ca. Ent., February, 1881, v. 13, pp. 31–33, fig. 3.
 - Review of paper by L. O. Howard; comparison of the alleged characters of Antigaster and Eupelmus to show grounds for the founding of the former; Didictyum synonymous with Hexaplasta; H. zigzag not a parasite of Aletia argillacea [=xylina], but of Phora aletia; habits of H. zigzag.
- 1933. [RILEY, C. V.] "A mystery in reference to *Pronuba yuccasella.*" <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 293.

 Critical review of paper by H. A. Hagen, who confounded *Prodoxus decipiens*
- 1934. [RILEY, C. V.] Mandible of Lithocolletis guttifinitella. < Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 294, fig. 138. Figure, with explanation, of the mandible of Lithocolletis guttifinitella.

with Pronuba yuccasella.

- 1935. [RILEY. C. V.] Excessive injury by a beetle in Russia. < Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 294.

 Ravages of Anisoplia austriaca in southern Russia; description of the same;
 - its habits and vernacular name.
- 1936. [RILEY, C. V.] Chemical change in the color of butterfly-wings. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 294.
 Note on paper by W. H. Edwards and J. M. Wilson.
- 1937. [RILEY, C. V.] Fungus foes. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 297.
 - Answer to inquiry of J. J. Brown; Cis fuscipes and all Cioidæ infest fungi growing on old trees and logs.

- 1938. [RILEY, C. V.] The twig-girdler. <Amer. Ent., December, 1880 [v 3], n. s., v. 1, p. 297, figs. 155–156.
 - Answer to letter of W. R. Maxwell; figures of larva, pupa, and imago of Oucideres cingulata and of twig injured by the imago; food-plants, habits, and transformations of this insect.
- 1939. [RILEY, C. V.] The bedeguar of the rose. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298, fig. 154.
 - Answer to letter of L. C. Bryan; account of the gall of Rhodites rosæ; figure of the same.
- 1940. [RILEY, C. V.] Minute borers in cherry, peach, and plum-trees. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298.
 - Answer to letters of J. L. Bennett and M. H. Boye; ravages, food-plants, and distribution of Scolytus rugulosus; food-habits of the species of Scolytus.
- 1941. [Riley, C. V.] Smilax injured by cut-worms. < Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298.
 - Answer to letter of G. Thommen; habits, ravages, and food-plants of Agrotis saucia.
- 1942. [RILEY, C. V.] Honey-producing oak-gall. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298.
 - Answer to letter of H. C. McCook; secretion of saccharine matter by galls growing on Quercus undulata; description of these galls, Cynips quercus-mellaria n. sp.; Myrmcocystus hortus-deorum [= melliger] collects the saccharine matter from these galls.
- 1943. [RILEY, C. V.] Dr. Hagen's mystery. <Ca. Ent., December, 1880, v. 12, pp. 263-264.
 - Review of paper by H. A. Hagen; Prodoxus decipiens confounded with Pronuba yuccasella.
- 1944. RILEY, C. V. Acorn-gall. <Trans. Acad. Sci. St. Louis, 1880, v. 4, p. 1 Proc.
 - Gall mistaken for abortive acorn, by G. B. Emerson, in his "Trees and shrubs of Massachusetts."
- 1945. RILEY, C. V. Silk-culture in the United States. Condensed account of the silk-worm and how to inaugurate a new source of wealth. <Western Farmer's Almanac for 1881, 1880, pp. 35-39, 4 figs. S.-b. No. 23, p. —.
 - Practicability and desirability of the culture of Scricaria mori in the United States; requisites to its success; superiority of S. mori over all other insects for silk-culture; brief illustrated natural history of the same; directions for silk-culture.
- 1946. RILEY, C. V. Legislation to control insects injurious to vegetation. <Farmer's Review, 20 January, 1881. S.-b. No. 23, p. 148. Reprint: <Amer. Nat., April [25 warch], 1881, v. 15, pp. 322–323. <Indiana Farmer, 16 April, 1881. S.-b. No. 23, p. 149. Notice: <Farmer's Review, 20 January, 1881. S.-b. No. 23, p. 170.
 - Appointment of C. H. Dwinelle, by the California State Horticultural Society, on a committee to consider what legislation is desirable to check the spread of noxious insects and to force land-owners to destroy the same; notice of the previous passage of such laws and of anticipated objections to them.

- 1947. RILEY, C. V. Larval habits of bee-flies, *Bombyliidæ*. < Amer. Nat., February [25 January], 1881, v. 15, pp. 143–145, figs. 1–3. Notice: < Western Stock Journ. and Farm, March, 1881, v. 11, p. 58.
 - Habits of Systachus and Triodites as parasites on eggs of Acridida; notice of paper by T. A. Chapman; figures larva, pupa, and imago of Systachus oreas and compares its larva and pupa with those of Bombylius major.
- 1948. RILEY, C. V. Experiments with pyrethrum: Safe remedies for cabbage-worms and potato-beetles. <Amer. Nat., February [25 January], 1881, v. 15, pp. 145-147.
 - Details of experiments made by A. J. Cook and W. R. Hubbert upon the effect of a dusting of pyrethrum powder upon larvæ of *Pieris rapæ*, larvæ and imagos of *Doryphora* 10-lineata, and upon *Eriosoma* [= *Pemphigus*] tessellata, Coreus [= Anasa] tristis, and flies and mosquitoes.
- 1949. RILEY, C. V. Insect enemies of the rice-plant. < Amer. Nat., February [25 January], 1881, v. 15, pp. 148-149.
 - Chalepus trachypygus feeds on roots of the rice-plant; conjectures as to other enemies of the rice-plant; Cecidomyia oryzæ injurious to the same in India.
- 1950. [RILEY, C. V.] The "yellow-fever fly." < Amer. Nat., February [25 January], 1881, v. 15, p. 150.
 - Review of paper by H. A. Hagen; food-habits of larvæ of Sciara sp.; occurrence of swarms of imagos of the same.
- 1951. [RILEY, C. V.] An aquatic Sphinx larva. Amer. Nat., February [25 January], 1881, v. 15, p. 151.

 Abstract of paper by H. A. Hagen.
- 1952. RILEY, C. V. Notes on the grape Phylloxera and on laws to prevent its introduction. <Amer. Nat., March [24 February], 1881, v. 15, pp. 238-241. Notice with extracts: <Pacific Rural Press, 23 April, 1881. S.-b. No. 42, p. 17.
 - Remarks upon letter of I. Bush; precautions to be adopted against the introduction of *Phylloxera vastatrix*; summary of the life-history of the same.
- 1953. RILEY, C. V. Hibernation of the cotton-worm moth: Ease with which mistakes are made. <Amer. Nat., March [24 February], 1881, v. 15, pp. 244-245, figs. 1-3.
 - Extract from letter of I. A. Wimbish; Leucania unipuncta mistaken for Aletia argillacea [=xylina]; characters of Aletia; figures ovipositor and eggs of L. unipuncta and imagos of both species.
- 1954. [RILEY, C. V.] On some interaction of organisms. < Amer. Nat., April [25 March], 1881, v. 15, pp. 323–324.
 - Review of paper by S. A. Forbes, with extracts; relative unimportance of 'special parasites as compared with predaceous animals of varied tastes in the limitation of the numbers of any species of animals; need of conservative action and exhaustive inquiry in the attempt to interfere with the order of nature.
- 1955. [RILEY, C. V.] Insect locomotion. <Amer. Nat., April [25 March], 1881, v. 15, p. 325.
 - Results of G. Carlet's studies on the order in which the feet are moved in the walking of *Hexapoda* and *Arachnida*.

1956. [Riley, C. V.] Plant-feeding habits of predaceous beetles. < Amer. Nat., April [25 March], 1881, v. 15, pp. 325-327.

Citation of evidence from numerous sources proving that certain Carabidæ and Coccinellida occasionally feed on plants, seeds, and spores.

1957. RILEY, C. V. Notes on Papilio philenor. < Amer. Nat., April, [25 March], 1881, v. 15, pp. 327-329, figs. 1-3.

Description of egg and newly-hatched larva of Papilio philenor; figure of larva, chrysalids, and imago of the same; food-plants, distribution, and its occurrence in swarms.

1958. [RILEY, C. V.] Entomological notes. < Amer. Nat., April [25] March], 1881, v. 15, pp. 330-331.

Notice of H. A. Hagen's paper on Simulium pictipes, with additional notes; abstracts and minor notices of other papers and items of news.

1959. [RILEY, C. V.] Second report of the United States Entomological Commission for the years 1878 and 1879, relating to the Rocky Mountain locust and the Western cricket, and treating of the best means of subduing the locust in its permanent breeding grounds, with a view of preventing its migrations into the more fertile portions of the trans-Mississippi country in pursuance of appropriations made by Congress for this purpose, with maps and illustrations. < Washington: 1880 [4 April, 1881], pp. 18+322+80, 10 figs., 17 pl., 9 maps.

TABLE OF CONTENTS.

LETTER OF TRANSMITTAL	IX IIIX
	AIII
CHAPTER I: By A. S. Packard and C. V. Riley. Additions to the chronology of locust ravages	1
The locust in 1878 in Minnesota and Iowa, 1; in Nebraska, Texas,	
and Indian Territory, 2; in Kansas, 2-3; in Dakota, 3-4; in	
Colorado, 4; in Wyoming, 4-5; in Utah and Idaho, 5-7; in	
eastern Oregon and Nevada, 7; in Montana, 7-9—Snmmary, 9—	
The locust in 1879 in Nebraska, Washington Territory, Dakota,	
and Texas, 10; in Colorado, 10-11; in Wyoming, 11-12; in	
Utah, 12-13; in Montana, 13-14.	
CHAPTER II: By C. Thomas.	
The relation of the locust and its ravages to agriculture and the	
settlement of the Territories	14
Character of the permanent region, 15—Importance of subduing	
the locusts in the permanent region, 15-16—Difficulties of burn-	
ing over, 16-18—Extermination impossible, but subjection pos-	
sible, 19-20—Settlement of the permanent region the best rem-	
edy, 19-20—Source of the more destructive swarms, 20—Rail-	
road needed in the permanent region, 21-Agricultural, 'not	
pastoral, population wanted, 22—Difficulties of constructing	
railroads in the permanent region, 22—Climatic character of the	
permanent region, 23—Disturbing and fighting the locusts in the	
permanent region, 24-25—Other proposed plans, 25—Plowing	
not feasible, 25-26—The locust question solved in the temporary	
region, 26—New definition of the temporary region, 27-28—The	
locust problem a national one, 29-31.	

CHAPTER III: By C. Thomas.

Facts concerning and laws governing the migrations of locusts in all countries

31

72

Most species of Aeridida not migratory, 31-Migratory disposition not caused by anatomical differences, 31-32-Writers on locust flights, 32-33—Earliest accounts of locust flights in Europe, 33-34-Locust flights in Europe in the middle ages, 34-37; in the seventeenth century, 37-38—The locust invasion of 1693 in Europe, 38-39-Locust flights in Europe in the eighteenth and nineteenth centuries, 40-41-Locusts in Asia and adjacent islands, 41; in Cyprns, 42; in Arabia and Syria, 43-44; in Persia, 44; in the Steppes, 45; in China, 46; in Manila, 46; in India, 46-47; in Australia, 47; in New Zealand, 47-Locust flights in Africa, 48; in northern Africa, 48-49; in Egypt, 49in Algiers, 50-51; in Abyssinia, 51; in central Africa, 51; in eastern Africa and the Canary Islands, 52-53-Locusts in South America, 53-54—Prevalence of locusts in deserts and dry regions, 54-55—Locust flights not governed by laws of periodicity, 55-56-Permanent breeding grounds of locusts, 56-57-Their geographical distribution, 58—Caloptenus spretus not in Mexico, 58— Migratory locusts in Yucatan and the West Indies, 58-59-In Nicaragua, 59—Geographical limits of Caloptenus spretus, 59; of Pachytylus migratorius, 59-60—Difficulty of determining species, 60-61—Definition of permanent home of migratory locusts, 62— Range of Pachytylus migratorius, 62-63; of Acridium peregrinum, 63-64—Comparison with the laws of geographical zoology, 65-66-Meeting point of three sub-regions, 67-Each species has its permanent breeding ground, 68—General character of the permanent breeding grounds, 68-69—Breeding ground of Caloptenus italicus, 69; of Pachytylus migratorius, 69-70; of Acridium peregrinum, 70; of Caloptenus spretus, 70-71.

CHAPTER IV: By C. Thomas.

Habits and characteristics of locusts in all countries within their areas of permanent distribution, so far as these relate to their movements

Caloptenus spretus always migratory, 72-73—Flights of C. spretus, 73-74—Invading swarms come from northwest, 74-75—Other opinions eliminated, 75-76—Swarms in 1875 and 1876, 76-77—Direction of returning swarms of C. spretus, 78; of other migratory locusts, 78-79—Direction of local flights, 79-81—Returning swarms do little injury, 81-82—Distance to which swarms may migrate, 82-84—Length of a single flight, 84-85—Swarms able to cross large bodies of water, 85-86—Swarms carried far by winds, 86-87—The mode of flight, 87—Position of the locust while flying, 87-88-Locusts flying with the wind, 88-89-Method of falling of swarms, 89-Formation of swarms, 89-91-Movements of swarms of C. spretus, as reported by Dr. Child, 91-93-Other reports, 94—Movements of swarms in other countries, 95-96— Swarms usually alight at night, 96-97-The height at which swarms move, 98-100-Different directions of swarms at the same time and place, 100—Fall of locusts into the sea, 100-102—Great flying power of locusts, 102—General causes of migration, 103— Excessive numbers, 103-104—Want of food, 104—Excessive heat,

109

156

160

163

	1959.	RILEY,	C.	V.]	-Co	ntinu	ed.
--	-------	--------	----	-----	-----	-------	-----

CHAPTER IV—Continued.

105—Dry condition of the atmosphere, 105-106—Arid condition of the mountain region, 106-107—Immediate causes of migration, 107-108.

CHAPTER V: By C. Thamas.

Influence of meteorological conditions on the development and migrations of locusts

Influence of heat and dryness, 109-110—Packard's table of locust years, 111-112-Table of rain-falls for 1860 to 1866, 112-Influence of the rainfall, especially in 1864, 112-113-Average monthly temperature for 1864 and 1866, 113-115—Rain-fall of the years 1863 to 1867, 116—Temperature of the years 1863 to 1867, 117—Extreme variations of monthly mean temperature from 1863 to 1867, 118-119-Monthly and annual mean temperature from 1872 to 1876, 119-121—Monthly and annual rain-fall from 1872 to 1878, 122-124-Extreme fluctuations between monthly mean temperature, 125-126—Greatest variations between monthly mean temperatures, 126-Variations of temperature between months of consecutive years, 127-128-Daily temperature at Fort Sully from 1872 to 1875, 129-134—Relation of temperature to the development of the eggs, 135-136-Professor Riley's experiments, 137-Professor Abbe's theory on the subject discussed, 137-140-Table showing temperature and direction and velocity of wind at Western stations, 141-154—Relation of temperature and winds to locust flights, 155.

CHAPTER VI: By A. S. Packard. '

The southern limits of the distribution of the Rocky Mountain locust. Few locusts permaneutly breeding in New Mexico and none in Arizona, 156—Caloptenus spretus probably not in southern New Mexico, central and southern Arizona, etc., 156—History of locust invasions in New Mexico from 1864 to 1879, 157–159—Southern range of the locust in Mew Mexico, 159—Changes in the map in the Report I, showing the distribution of the locust, 159–160.

CHAPTER VII: By A. S. Packard.

CHAPTER VIII: By A. S. Packard.

The western cricket.....

Difference in habits between the western cricket and the locust, 163—Occurrence of Anabrus purpurascens, 163–164; of A. simplex, 164—Mode of egg-laying, 164—Movements of an army of the western cricket, 164–165—Its ravages, 165–166—Its food, 166—Its enemies and parasites, 166—Its breeding habits, 166–167—Remedies, 167–168—Geographical distribution of the species of Anabrus, 168–169—Synopsis of the species of Anabrus and its allies, 169–170—External anatomy of Anabrus, 170; the head, 170–172; the thorax, 172–174; the abdomen, 174–175—Internal anatomy of Anabrus purpurascens, 175; the digestive system, 175–176; the nervous system, 176–177; the breathing apparatus, 177–178.

CHAPTER IX: By A. S. Packard.

CHAPTER X: By C. S. Minot.

Histology of the locust (Caloptenus) and the cricket (Anabrus).....

Study of histology of insects, 183–184—Explanations of Figs. 1 and 2, Plate II, 185–186—The cuticula, 186–188—The epidermis, 188–189—Sense organs, 189—Nervous system, 190—Tracheæ, 191–195—Air-sacs and spiracles, 195—Muscles, 195–196—Organs of circulation, 196—Connective tissue, 196–198—Ovary, 198–201—Anterior eæcum of the oviduct, 201—Uterus of locusts, 201–203—Male organs, 203—Testis, 203–204—Development of the spermatozoa, 204, 207—Vasa deferentia, 207—Ductus ejaculatorius, 207—Vesiculæ seminales, 207–208—Digestive canal, 208–210—Crop, 210–211—Proventriculus, 211–212—Stomach, 212–214—Diverticula, 214–215—Gastro-ileal folds, 215–217—Ileum, 217—Colon, 217–218—Rectum, 218–219—Summary on the digestive canal, 219–221—Malpighian vessels, 222.

CHAPTER XI: By A. S. Packard.

The brain of the locust..... The nervous system in general, 223-224—The brain of insects compared with that of vertebrates, 224-226-The brain of the adult locust, 226—Histological elements of the brain, 226-228—The sections of the brain, 228-230-Internal topography of the brain, 230—The central body, 230, 231—The mushroom bodies, 231-234— The optic lobes, 234--The optic ganglion, 234--The antennal or olfactory lobes, 235—The commissural lobes, 235—The brain of locusts compared with that of other insects, 235-236-Structure of the brain in the embryo locust, 236—The brain of the embryo locust compared with the first thoracic ganglion, 238—The brain in the second embryonic stage, 238-239-Structure of the subcesophageal ganglion, 239—The brain of the freshly-hatched larva of Caloptenus spretus, 239; of the third larval stage of C. bivittatus, 239; of the second or last pupal stage of C. spretus, 240-241— Bibliography of the internal structure of the brain of crustacea

CHAPTER XII: By C. V. Riley and C. Thomas.

and insects, 241-242.

Locust ravages in California

Position of the genus Camnula in the Œdipodini, 242-243—Description of the genus Camnula, 243; of Œdipoda pellucida, 243; of Œdipoda atrox, 243-244; of Camnula pellucida, 244-246—The locust in California in 1878, 246; in Sierra Valley, 247; in the vicinity of Loyalton, 247-248; of Sierraville, 249—History of locust devastation in California, 250—Enemies of the Californian locust, 251—Protective measures, 251-252—Life history of Camnula atrox, 252-253—The red or locust mite, 253—Damage done by the locust in Sierra Valley, 254-255—The locust in California in 1879, 255-257—Description of Œdipoda obliterata, 257-259—Remarks on Cratypedes Putnami, 259.

CHAPTER XIII: By C. V. Riley.

Further facts about the natural enemies of the locust.

Blister-beetle larvæ feeding on the eggs of the Californian locust, 259-260—Retardation in the development of blister-beetles, as shown in *Epicauta vittata*, 260—Philosophy of such retardation, 260-261—Eggs of *Chauliognathus pennsylvanicus*, 261—Habits of the young larva, 261-262—Egg-laying of asilid-flies, and particularly of *Mallophora orcina*, 262—Bee-fly larvæ common among the

183

223

242

259

CHAPTER XIII—Continued.

eggs of Camnula pellucida, 263—Habits of bee-fly larvæ, 263–264—Observations on the larval habits of Bombylii, 264–265—Larval habits of Argyramæba, 265–266—Abundance of bec-flies and blister-beetles in the Western country, connected with the abundance of locusts, 266—Life history of Systæchus oreas, 266–267—Description of its larva, 267; of the pupa, 267–268; of the imago, 268—Description of Systæchus oreas, 268—Larva of Triodites mus, 268–269—Imago of Triodites mus, 269—Dr. T. A. Chapman's observations on Bombylius major, 269—Hair-worms and red-mites abundant upon locusts in California, 270—Chalcid-fly parasitic on locusts, 270—Synonymy of the locust-egg parasite, 270—Diggerwasps killing locusts, 270.

CHAPTER XIV: By C. V. Riley.

Conrses that may be adopted by the General Government to lessen locust injury.....

271

Importance of destroying the locusts in their native breedinggrounds, 271-Recapitulation of means suggested in first report, 271-Importation of English rooks, 271-Importance of burning over the permanent breeding-grounds, 272—Breeding-grounds occupy comparatively small areas in the permanent region, 273— Protection from invading swarms, 273-Locust warnings through the Signal Bureau, 273-Diverting swarms by means of smoke, 274—Co-operation of governments and governmental institutions, 274—Apathy apt to result from periods of immunity from locust invasions, 274-How the Government can aid, 275-Surface characteristics of the permanent region and the proportion of burnable land, 275—Number of square miles in the permanent region, 276—The plains area east of the mountains, 276—Its vegetation, 276-Burnable land practically indentical with grazing-land, 277—The plains area in the British possessions, 278—In the United States, 279—The mountain area, 280—Timber-lands in the mountain area dependent upon latitude and altitude, 280-Northern section of the mountain area, 280, 281—River valleys in Montana, 281-Valley of the Yellowstone and its tributaries, 281—Of the Madison River, 282—Of the Jefferson and Missouri Rivers, 283—The Wind River and the Bighorn Mountains, 283— The Green River Basin, 283—Valley of the Snake River and its tributaries, 286, 289—Southern section of the mountain area, 289—In southern Wyoming, 290-291—In Colorado, 291—The San Luis Valley, 292-Mountain area in New Mexico, 292-The platean area, 293—Its extent, 293—The Roan Plateau, 293—The Uinta Valley, 293—The Grand River Valley, 294—The Great Sage Plain, 294—The San Juan River Valley, 295—The Colorado River Platean, 205—The Great Basin area and the Wasatch Mountains, 296-Valleys of the Bear River and its tributaries, 296-The Wasatch Range and its valleys, 298—Mountain Ranges in Nevada, 299-300—The Mojave Desert, 300—Preventive measures in the plains region, 300—Caloptenus spretus breeds, especially in British America and Montana, 300-Fertile grass-land in the Territories, 300—Its extent, 302—Encouragement to settlement, 302— Modification of climate by settlement and cultivation, 302—Professor Thomas' communication to Governor Pillsbury, 303—The settlement of Dakota will benefit Minnesota, 304-Forest planting

CHAPTER XIV-Continued.

on the coteau of the prairies, 305-Preserving of lakes, ponds, and swamps in Minnesota, 306—Encouragements to railroads, 307— Irrigation, 307—Extent of the arid region, 307-308—Only a small portion of the land irrigable, 308-Agriculture in the arid West and in the Mississippi Delta require assistance of the National Government, 308-Prevention of floods by means of reservoirs, 309—Cost of irrigation, 310—Influence of irrigation on the locust question, 310-Mr. H. Gannett's report on irrigation in the arid region, 310-Waste of water in irrigating, 311-Amount of irrigable land, 311—Amount of water necessary for irrigation, 312-313—Irrigable land in Colorado, Utah, California, &c., 313— Arable land in the Territories, 314—Amount of land in actual cultivation, 314—Irrigation by means of artesian wells, 315— Methods of irrigation, 315-316—Uselessness of reservoirs and great irrigating canals in the arid region, 316—How irrigation should be regulated, 316—Preservation of forests; tree-planting, 317— Judicious burning in spring, 317-Permanent establishment of locust signals and warnings, 318-Co-operation with the Dominion Government, 319-Preventive measures in the mountain and plateau areas, 319-Breeding-grounds of the locust in the mountain area, 319-Influence of settlement, 320-Replanting of forests, 320-Importance of locust signals to the farmer in Colorado and Utah, 321—Very destructive locust invasions not likely to recur in the future, 321—The locust has ceased to be an object of dread, 322.

APPENDIX 1:	
Miscellaneous data and replics to Circular No. 1	[3]
Circular No. 1, [3]—Nebraska data, [4]—Data for Minnesota, [17]—	
Data for Iowa, [20]—Data for Dakota, [21].	
APPENDIX II: By S. H. Scudder.	
List of the Orthoptera collected by Dr. A. S. Packard, jr., in the	
western United States in the summer of 1877	[23]
Bradynotes opimus, Scudd., n. sp., [24]—Pezotettix pacificus,	
Scudd., n. sp., [24]—Gomphocerus shastanus, Scudd., n. sp., [25]—	
Circotettix maculatus, Scudd., n. sp., [26]—Trimerotropis latifas-	
ciata, Scudd., n. sp., [26]—T. similis, Scudd., n. sp., [27]—T. caru-	
leipes, Scudd., n. sp., [27]—Psinidia wallula, Scudd., n. sp., [27].	
APPENDIX III:	
Report of John Marten	[29]
Observations of John Marten, special agent, on locusts in Iowa,	
[29]—In Minnesota, [30]—In Dakota, [30]—In Nebraska, [31].	
APPENDIX IV: By B. P. Mann and C. Thomas.	
Bibliography of some of the literature concerning destructive lo- custs	[33]
Introductory and explanatory remarks, [33]—Bibliography on the	
destructive locusts of Europe, Asia, and Africa, [33]—Supplementary list to the bibliography on locusts in foreign countries,	
[50]—Additions to bibliography on the locusts in America, [55].	
APPENDIX V:	
Data concerning locust in Texas	[57]
Notes furnished by Gen. A. J. Myer, U. S. A., Chief Signal Officer,	1

on locusts in Texas, chiefly in the year 1877, [57]—Locust data

from Indian Territory in 1877, [61].

1959.	RILEY, C. V.]—Continued.	
	APPENDIX VI:	
	On the flight of locusts	[63]
	Translation, by F. P. Spofford, of a memoir by Gaetano de Lucre-	
	tus, [63]—Translation of d'Azara's paper on locust invasions in	
	Spain from 1754 to 1757, by F. P. Spofford, [66].	
1	APPENDIX VII:	
	Notes of a journey made to Utah and Idaho in the summer of 1878,	
	by A. S. Packard, jr	[69]
	Observations on locusts in Nebraska and Colorado, [69]—In Wy-	
	oming, [69]—In Utah, [69]—In Idaho, [70]—Observations on	
	the journey returning East, [71].	
4	APPENDIX VIII:	
	Yersin's rescarches on the functions of the nervous system of the ar-	
	ticulate animals	[73]
	Translation, by Dr. A. S. Packard, jr., of Saussure's abstract from	
	Yersin's Recherches sur les fonctions du système nerveux dans	
	les animaux articulés, [73].	

- 1960. RILEY, C. V. Notes on North American Microgasters, with descriptions of new species. <Trans. Acad. Sci. St. Louis, 16
 April 1881, v. 4, pp. 295-315, 9 figs. Separate: <[St. Louis, Mo.], 6 April, 1881, 20 pp., 9 figs.
 - Characteristics and habits of the *Microgasters*; description of larva of *Apanteles aletiæ*; the cocoons and their construction; effect of the parasites upon the longevity of the host; secondary parasites; habits and descriptions of several new species and varieties, for a list of which see the *Systematic Index*; figures *A. aletiæ*, healthy and parasitized larvæ of *Chærocampa pampinatrix* [= *Ampelophaga myron*]; also mass of *Microgaster* cocoons, and the formation of the cocoon by the larva.
- 1961. [RILEY, C. V.] Exuviation in flight. < Amer. Nat., May [16 April], 1881, v. 15, p. 395.
 - Criticism of R. McLachlan's observations; process of casting subimaginal skin in *Polymitarcys alba*.
- 1962. RILEY, C. V. The rascal leaf-crumpler in Georgia. <Amer. Nat., May [16 April], 1881, v. 15, p. 400.

Occurrence and ravages of Acrobasis nebulo [=indiginella] in Bryan County, Georgia.

1963. [RILEY, C. V.] Vertical vs. horizontal insect boxes. < Amer. Nat., May [16 April], 1881, v. 15, p. 401.

Review of pamphlet by A. Preudhomme de Borre; reply of G. H. Horn to Preudhomme de Borre.

1964. [RILEY, C. V.] Insects affecting the China tree. < Amer. Nat., May [16 April], 1881, v. 15, pp. 401-402.

General immunity of the China tree from the attacks of insects and its consequent value as a shade tree; occurrence of a *Lecanium* and *Ceroplastes* n. sp. upon it in Alabama; injuries to its leaves by *Atta fervens* in Texas.

1965. [RILEY, C. V.] Galls on *Eucalyptus*. <Amer. Nat., May [16 April], 1881, v. 15, p. 402.

Critical review of paper by R. McLachlan; the supposed dipterous galls probably cecidomyidous; the supposed lepidopterous galls probably not lepidopterous, but inhabited by a lepidopterous inquiline.

- 1966. [RILEY, C. V.] North American Anthomyidæ. <Amer. Nat., May [16 April], 1881, v. 15, p. 402.

 Notice and summary of paper by H. A. Hagen.
- 1967. [RILEY, C. V.] Galls and gall-insects. <Amer. Nat., May [16 April], 1881, v. 15, pp. 402–403.
 - Review of paper by H. F. Bassett, with additional notes on the gall of Cynips [= Andricus] quercus-californica found on Quercus douglasii and infested by Ozognathus cornutus; the Cynips produced from this gall all females; earlier accounts of the gall and of the habits of O. cornutus.
- 1968. RILEY, C. V. Scale insect on raspberry. <Farmer's Review, 21 April, 1881, v. 6, p. 243. S.-b. No. 23, p. 150. Reprint: <Amer. Nat., June [19 May], 1881, v. 15, p. 487.
 - Remarks on communication from R. B. Fulton; occurrence and ravages of a scale (Diaspis harrisii?) [= Chionaspis furfurus] on black-cap raspberry in Mississippi; mode of dissemination and means of extermination of Coccidæ.
- 1969. RILEY, C. V. Descriptions of some new *Tortricidw*, leaf-rollers. <Trans. Acad. Sci. St. Louis, May, 1882, v. 4, pp. 316–324. Separate: <[St. Louis, Mo.], 28 April, 1881, 9 pp.
 - Description of two (2) new genera and thirteen (13) new species of *Tortricidæ*; see the *Systematic Index* for a list of the same; mention of the food-plants of some of the species.
- 1970. RILEY, C. V. Larval habits of bee-flies. <Amer. Nat., June [19 May], 1881, v. 15, pp. 438-447, pl. 6. Separate: <pp. 438-447. Adapted from the 2d Rept. U. S. Ent. Commission, pp. 262-269; see No. 1959 for synopsis of contents.
- 1971. [RILEY, C. V.] The periodical Cicada alias "seventeen-year locust." <Amer. Nat., June [19 May], 1881, v. 15, pp. 479-482, fig. 1. Correction: <Ibid., July [22 June], 1881, p. 578.
 - Extract from 1st Rept. State Ent. Mo., with additional notes; figures eggs, pupæ, and imago of *Cicada* [= *Tibicen*] *septendecim*, and of twig punctured by the imago for the deposition of her eggs.
- 2972. RILEY, C. V. A new species of oak coccid mistaken for a gall. <Amer. Nat., June [19 May], 1881, v. 15, p. 482.
 - Description of Kermes galliformis n. sp. [p. 482] occurring on Quercus palustris in the sonthern and central United States; the coccid infested by Euclemensia bassettella.
- 1973. [RILEY, C. V.] The "water-weevil" of the rice-plant. <Amer. Nat., June [19 May], 1881, v. 15, pp. 482–483. Extract: <Ann. Rept. [U.S.] Commissioner Agric. for 1881 and 1882, pp. 130–133. Note on the injury done to rice plants by the larvæ of Chalepus trachypygus
 - Note on the injury done to rice plants by the larvæ of *Chalepus trachypygus* and by other larvæ, probably *Lissorhoptrus simplex*; letter from J. Screven; seasons, habits, and ravages of the last-mentioned species.
- 1974. [RILEY, C. V.] The impregnated egg of *Phylloxera vastatrix*. <Amer. Nat., June [19 May], 1881, v. 15, pp. 483, 484. Reprint: <Amer. Wine and Grape Grower, 1 July, 1881, v. 3, p. 104. S.-b. No. 42, pp. 12-13.
 - Verification by P. Graell of author's conclusion that the impregnated egg of *Phylloxera vastatrix* would generally be found to hatch in the same season in which it was laid.

- 1975. RILEY, C. V. Works on North American Microlepidoptera. Amer. Nat., June [19 May], 1881, v. 15, pp. 484–486.
 - Review of works of T. de Grey [Lord Walsingham]; discussion of new genera; list of *Tortricidæ* of economic interest common to Europe and North America; additions to the synonomy of *Tortricidæ*.
- 1976. [RILEY, C. V.] Moths mistaken for *Aletia*. <Amer. Nat., June [19 May], 1881, v. 15, pp. 486-487.
 - Platyhypena [= Hypena] scabra and Phoberia atomaris mistaken for Aletia argillacea [= xylina]; hibernating habits of the first species.
- 1977. [RILEY, C. V.] Specific value of Apatura alicia Edw. < Amer. Nat., June [19 May], 1881, v. 15, p. 487. Critical review of paper by W. H. Edwards.
- 1978. RILEY, C. V. Antigaster vs. Eupelmus. <Ca. Ent., May, 1881, v. 13, p. 114.
 - Validity of L. O. Howard's reasons for considering Antigaster a synonym of Eupelmus.
- 1979. RILEY, C.V. The periodical Cicada alias "seventeen-year locust." <Farmer's Review, 16 June, 1881, v. 6, p. 370. S.-b. No. 42, pp. 18-20.
 - Extract from 1st Rept. State Ent. Mo., with additional notes and request for further information in regard to the chronology and geographical distribution of the broads of Cicada [= Tibicen] septendecim and C. [= T.] tredecim which appear in 1881.
- 1980. [RILEY, C. V.] Dimorphism in *Cynipidw*. <Amer. Nat., July [22 June], 1881, v. 15, p. 566.
 - Claim of author to have made the first record at least in North America of proof of dimorphism in Cynipidw; notice of writings of B. D. Walsh, H. F. Bassett, and H. Adler on the subject; list of species of Cynipidw in which the occurrence of dimorphic forms has been proven, and of closely allied species in which no alternate generation seems to occur.
- 1981. [RILEY, C. V.] Blepharoceridæ. <Amer. Nat., July [22 June], 1881, v. 15, pp. 567-568.
 - Account of various investigations into the natural history of Blepharoceridæ; description of larvæ and pupæ of these flies.
- 1982. [RILEY, C. V.] Braula cæca not particularly injurious to the honey-bee. <Amer Nat., July [22 June], 1881, v. 15, p. 568.

 Notice of paper of J. Fedarb; ravages of Braula cæca in hives generally overestimated at present and formerly unknown.
- 1983. [RILEY, C. V.] Economic entomology in England. <Amer. Nat., July [22 June], 1881, v. 15, p. 568.

 Notice of Miss E. A. Ormerod's report for 1880.
- 1984. [RILEY, C. V.] The cultivation of pyrethrum and manufacture of the powder. <Amer. Nat., 1881, v. 15, July [22 June], pp. 569-572; September [23 August], pp. 744-746; October [23 September], pp. 817-819. See: <Amer. Wine and Grape Grower, December, 1881, v. 4, p. 22. S.-b. No. 51, p. 149.

• 'h •

Partial history of the use of pyrethrum powder as an insecticide; directions for the cultivation of the plants and for the production and use of the powder.

- 1985. [RILEY, C. V.] Hudson Bay Lepidoptera. <Amer. Nat., July [22 June], 1881, v. 15, pp. 572, 573.
 - Review of paper of J. J. Weir; reprint of the list of Lepidoptera and of the remarks on the explanation of the relations of the fauna of Hudson's Bay to that of Europe; probability that the species identical with those of Europe are recent immigrants.
- 1986. [RILEY, C. V.] Trade in insects. <Amer. Nat., July [22 June], 1881, v. 15, p. 573.
 - General decline in prices of insects within sixty years or more; statement of some prices obtained for Coleoptera recently.
- 1987. [RILEY, C. V.] Ants injurious in Arizona. <Amer. Nat., July [22 June], 1881, v. 15, pp. 573, 574.

Statement by H. H. Rusby of ravages of Formicide, which occur in vast colonies.

- 1988. [RILEY, C. V.] Covering of egg-puncture mistaken for *Dorthesia*. <Amer. Nat., July [22 June], 1881, v. 15, p. 574.
 - Waxy material covering the egg-punctures of Enchophyllum [= Enchenopa] binotata, labeled Dorthesia viburni and D. celastri in collection of A. Fitch.
- 1989. [RILEY, C. V.] [Dolerus unicolor.] < Amer. Nat., July [22 June], 1881, v. 15, p. 574.
 - Doubts the statement of H. Keenan that the imagos of *Dolerus unicolor* injure the fruit buds of pear-trees.
- 1990. [RILEY, C. V.] Supposed army-worm in New York and other Eastern States. < Amer. Nat., July [22 June], 1881, v. 15, pp. 574-577.
 - Ravages of Nephelodes violans and Crambus vulgivagellus in New Jersey, Long Island, and northern New York; natural history, vernacular names, and description of the larva of the former species; previous accounts of this larva.
- 1991. [RILEY, C. V.] Migration of butterflies. <Amer. Nat., July [22 June], 1881, v. 15, p. 577.
 - Report by J. H. Mellichamp of the eastward flight of thousands of *Pieris monuste* in small groups of two, three, or more individuals, on 1 and 2 June, 1881, over Bluffton, S. C.; description of larva and pupa; food-plants of larva.
- 1992. [RILEY, C. V.] Classification of the mites. < Amer. Nat., July [22 June], 1881, v. 15, pp. 577-578.
 - Abstract of letter of G. Haller; obscrvations on the appendages and systematic position of the Acarina.
- 1993. [RILEY, C. V.] Carrying out the law. <Amer. Nat., July [22 June], 1881, v. 15, p. 578.
 - A fine of £5 imposed upon a man in England for importing living *Doryphora* decemlineata into that country.
- 1994. RILEY, C. V. Locusts and locusts. <N. Y. Tribune, 22 June, 1881. S.-b. No. 32, pp. 55-56; No. 42, pp. 7-11.
 - Cicada [= Tibicen] septendecim and C. [T.] tredecim compared with Caloptenus spretus to explain and correct the confusion of the species by those who call them all "locusts"; differences in the habits and characters of these

- 1994. RILEY, C. V.—Continued.
 - insects, their periodicity and distribution; no fear that *C. spretus* will commit great ravages this year; recommendation of a system of observations and warnings by the United States Signal Service to guard against unexpected inroads of *C. spretus*.
- 1995. RILEY, C. V. The caterpillar nuisance. <Evening Star [Washington, D. C.], 24 June, 1881, v. 57, No. 8802, p. 3. S.-b. No. 23, p. 152; No. 42, p. 13. Reprint: <Amer. Nat., September [23 August], 1881, v. 15, pp. 747-748, 1 fig.
 - Natural history of *Hyphantria textor* [=cunea]; means against it; methods of applying poisoned liquids to trees.
- 1996. RILEY, C. V. Directions for cultivating pyrethrum for insect powder. < Gardener's Mo. and Hortic., June, 1881, v. 23, pp. 172-173. S.-b. No. 23, p. 153.
 - Directions for raising Pyrethrum cinerariæfolium from seed; climate not favorable to the growth of P roseum; preparation of these plants for the destruction of insects.
- 1997. RILEY, C. V. Cotton-worms and Cicadas. Prof. Stelle's logic! <Selma [Ala.] Times, 19 July, 1881. S.-b. No. 24, pp. 44-45; No. 39, p. 123.
 - Criticism of statements of J. P. Stelle; impossibility of exterminating Aletia xylina; waves of destructiveness; Cicada [= Tibicen] tredecim abundant in sections of Alabama, as predicted.
- 1998. RILEY, C. V. The Rocky Mountain locust alias Western grass-hopper. <Amer. Agric., July, 1881, v. 40, pp. 283-284, 6 figs. S.-b. No. 23, p. 142.
 - Summary of facts concerning the egg-laying development, habits, food-plants, destructive powers, migration, flight, and enemies of and means against Caloptenus spretus; figures of the eggs of the insect in all stages, of the manner of oviposition, and of pans for the destruction of the locusts; prospects of future injury.
- 1999. RILEY, C. V. Lepidopterological notes. <Papilio, July, 1881, v. 1, pp. 106-110.
 - Advance print of extracts from Bull. No. 6, U. S. Entomological Commission, pp. 56-58, 78, 82-83. See No. 2026 for synopsis of contents.
- 2000. RILEY, C. V. Further notes on the pollination of Yucca and on *Pronuba* and *Prodoxus*. <Proc. Amer. Assoc. Adv. Sci. for 1880, 14 October, 1881, v. 29, pp. 617-639, 16 figs. Separate: <[Salem, Mass, July, 1881], 23 pp., 16 figs.
 - Recapitulation of published observations on Pronuba yuccasella and Prodoxus decipiens, with additional observations and references to literature; habits and functions of the two species; descriptions and figures of the generic characters of Pronuba, Prodoxus, and Hyponomeuta, and of the specific characters of Pronuba yuccasella, P. maculata n. sp., Prodoxus decipiens, P. intermedius n. sp., P. marginatus n. sp., P. cinereus n. sp., P. anescens n. sp., Hyponomeuta malinella, and H. multipunctella, especially the structure of the ovipositor and the male characters, and the venation of the wings; discussion of the structure of the ovipositor in Lepidoptera characterization of the new family Prodoxida to comprise Pronuba and Prodoxus; generic and specific synonymy; reply to criticism of T. Meehan; list of insects frequenting Yucca; synonymical list of the described Prodoxia.

- 2001. RILEY, C. V. Additional notes on the army worm, Leucania unipuncta. <Proc. Amer. Assoc. Adv. Sci. for 1880 [14 October], 1881, v. 29, pp. 640-642. Separate: <Salem, Mass., July, 1881, pp. 24-26. Abstract: <Boston Daily Advertiser, 2 September, 1880, v. 136, p. 4.
 - Number of annual generations of *Leucania unipuncta*; stages at which hibernation takes place; occasional immigration of moths into uninfested districts; preferred breeding-places; connection of wet and dry seasons with the abundance of these insects.
- 2002. RILEY, C. V. The hitherto unknown life-habits of two genera of bee-flies, *Bombyliidæ*. <Proc. Amer. Assoc. Adv. Sci. for 1880 [14 October], 1881, v. 29, p. 649. Separate: <Salem, Mass., July, 1881, p. 33.
 - Food-animals and food-habits of Systachus oreas and Triodites mus; parallelism in the life-history of Bombyliida and Meloida; connection of the abundance of insects of these families in the western United States with the prevalence of Acridida there; retardation of development in the early stages of these parasites explained as a beneficial characteristic.
- 2003. RILEY, C. V. A remarkable case of retarded development. <Sci. Amer., 20 August, 1881 [v. 59], n. s., v. 45, p. 116. S.-b. No. 23, p. —.
 - Report of a case in which eggs of Caloptenus spretus remained undeveloped for a period of four and one-half years after being laid; remarks upon the problems involved in the explanation of retarded development.
- **2004.** [RILEY, C. V.] Blepharoceridæ. <Amer. Nat., September [23 August], 1881, v. 15, p. 748.
 - Discovery by J. Q. Adams of pupe and images of Blepharoceridæ at Watertown, N. Y.
- 2005. [RILEY, C. V.] Remarkable case of retarded development, <Amer. Nat., September [23 August], 1881, v. 15, pp. 748-749.
 - Eggs of Caloptenus spretus, buried about 25 cm. (10 inches) under ground, remained unhatched and alive for four and one-half years, and hatched upon being exhumed.
- 2006. [RILEY, C. V.] Promotion of silk-culture in California. < Amer. Nat., September [23 August], 1881, v. 15, p. 749.
 - Answer to letter of Mrs. T. H. Hittell; popularization of silk-culture in California; preparation of the fiber for the market.
- 2007. [RILEY, C. V.] Locust flights in Dakota. >Amer. Nat., September [23 August], 1881, v. 15, pp. 749-750.
 - Flights of Caloptenus spretus in Dakota on 7th and 16th July, 1881; large numbers of Diplax rubicundula accompanying the locusts.
- 2008. [RILEY, C. V.] The Hessian fly. <Amer. Nat., September [23 August], 1881, v. 15, p. 750.
 - Extensive damages done by Cecidomyia destructor in Illinois and Missouri; its abundance in the western prairie States in 1881.

- 2009. [RILEY, C. V.] The genuine army-worm in the West. < Amer. Nat., September [23 August], 1881, v. 15, p. 750.
 - Occurrence and ravages of Leucania unipuncta in Illinois and Indiana in 1881; these larvæ either from eggs of hibernated moths or of a second brood; other insects mistaken for this in other States; periods of the Leucania and influence of the seasons upon it.
- 2010. [RILEY, C. V.] A new imported enemy to clover. <Amer. Nat., September [23 August], 1881, v. 15, pp. 750-751.

Clover injured by Phytonomus punctatus at Barrington, N. Y., in July, 1881.

- 2011. [RILEY, C. V.] Another enemy of the rice-plant. <Amer. Nat., September [23 August], 1881, v. 15, p. 751.
 - Rice-plants injured by the larva of Laphygma frugiperda in Georgia in the summer of 1881.
- 2012. [RILEY, C. V.] Canker-worms. < Amer. Nat., September [23 August], 1881, v. 15, p. 751.

Excessive ravages of Palcacrita vernata in Tazewell County, Illinois, in 1881.

- 2013. [RILEY, C. V.] Lepidopterological notes. < Amer. Nat., September [23 August], 1881, v. 15, pp. 751-752.
 - Advance extract from Bull. No. 6, U. S. Entomological Commission, pp. 55-56. See No. 2026 for synopsis of contents.
- 2014. RILEY, C. V. The periodical Cicada. <Amer. Agric., August, 1881, v. 40, p. 132, 5 figs. S.-b. No. 23, p. 141.
 - Brief sketch of the natural history of $Cicada \ [= Tibicen]$ septendecim and C. [= T.] tredecim, with figures of all stages and of punctured twigs; the reputed stinging by these insects probably done by Stizus graudis $[= Sphecius \ speciosus]$; figure of the Stizus; recommends the substitution of the name "periodical Cicada" in place of the name "locust."
- 2015. RILEY, C. V. Entomological notes. <Farmer's Review, 22 September, 1881. S.-b. No. 42, pp. 54-55. Reprint: <Amer. Nat., [3] December, 1881, v. 15, pp. 1012-1013. See: <Sci. Amer., 11 February, 1882. S.-b. No. 38, p. 1½.
 - Abstract and critical review of paper by M. Saint-André; attempted explanation of the immunity of grape-vines in sandy soil from the ravages of *Phylloxera vastatrix*.
- 2016. RILEY, C. V. Notes on Hydrophilus triangularis. <Amer. Nat., October [23 September], 1881, v. 15, pp. 814-817, figs. 1-2.
 - Description and figures of egg-case and figures of eggs and of male imago, with structural details of *Hydrophilus triangularis*; characters and life-history of the larva and pupa; figures of eggs, egg-case, larva, and pupa of *H. piceus* of Europe.
- 2017. [RILEY, C. V.] Migration of plant-lice from one plant to another. <Amer. Nat., October [23 September], 1881, v. 15, pp. 819–820.
 - Exposition of J. Lichtenstein's theory that most Aphidida, especially gall-making Pemphigini, live upon two different plants in passing through their cycle of development; cites instances in which the host-plants are of different families.
- 2018. [RILEY, C. V.] The chinch-bug. <Amer. Nat., October [23 September], 1881, v. 15, pp. 820-821.
 - The abundance of *Blissus leucopterus* is connected with the moisture of the season; extraordinary noxiousness of this insect in the dry season of 1881 in various parts of the United States.

- 2019. [RILEY, C. V.] Phylloxera laws. < Amer. Nat., October [23 September], 1881, v. 15, p. 821.
 - Summary of the existing laws regulating the traffic in plants, with a view of preventing the introduction of *Phylloxera vastatrix* into different countries.
- 2020. [RILEY, C. V.] One half the vine area of France affected by Phylloxera. <Amer. Nat., October [23 September], 1881, v. 15, p. 821.
 - Statement that nearly one-fourth of the area of vineyards in France is destroyed by *Phylloxera vastatrix* and as much more attacked; submersion the only sure remedy, and the use of American resisting stocks the only available preventive.
- 2021. [RILEY, C. V.] London purple and Paris green. < Amer. Nat., October [23 September], 1881, v. 15, p. 821.
 - London purple more than twice as efficacious and less expensive than Paris greeu as an insecticide; it needs to be well ground; composition of London purple.
- 2022. [RILEY, C. V.] Entomologist for the Pacific coast. < Amer. Nat., October [23 September], 1881, v. 15, pp. 821–822.
 - Notice of attempts made to have a State entomologist appointed in California.
- 2023. [RILEY, C. V.] Dilar in North America. <Amer. Nat., October [23 September], 1881, v. 15, p. 822.

 Notice of paper by R. McLachlan.
- **2024.** [RILEY, C. V.] Locusts in Nevada. < Amer. Nat., October [23 September], 1881, v. 15, p. 822.
 - Statement from Reno [Nev.] Journal that the countless locusts which hatched in western Nevada in the spring of 1881 flew toward the Sicrra Nevada without doing any injury.
- 2025. [RILEY, C. V.] Odor in butterflies. <Amer. Nat., October [23 September], 1881, v. 15, p. 822.

 Notice of paper by Miss M. E. Martfeldt.
- 2026. RILEY, C. V. General index and supplement to the nine reports on the insects of Missouri. <Bull. No. 6, U.S. Ent. Commission [24 March], 1881, 178 pp.
 - Order of matter: (1) Introduction. (2) Table of contents. (3) Corrections. (4) Notes and additions. (5) Descriptions of new species and varieties. (6) List of descriptions of adolescent states. (7) List of descriptions, mostly amplified, of species not new. (8) List of illustrations by reports. (9) Classified list of illustrations. (10) General index. (11) Index to foodplants.
- 2027. [RILEY, C. V.] The permanent subsection of entomology at the recent meeting of the A. A. A. S. < Amer. Nat., 1881, v. 15, November [28 October], pp. 909-912; [3] December, pp. 1008-1011.
 - Notice of the meeting at Cincinnati, Ohio, August, 1881, with abstracts and notices of papers read.
- 2028. RILEY, C. V. The new imported clover enemy. <Amer. Nat., November [28 October], 1881, v. 15, pp. 912-914.
 - Occurrence and habits of *Phytonomus punctatus* in New York; description of the egg and larva of the same; food-plants of the European species of *Phytonomus*.

- 2029. RILEY, C. V. Crambus vulgivagellus. <Amer. Nat., November [28 October], 1881, v. 15, pp. 914-915.
 - Excessive abundance of this species in the eastern United States in 1881; description of the egg; method of oviposition.
- 2030. RILEY, C. V. Larval habits of *Sphenophori* that attack corn. Amer.Nat., November [28 October], 1881, v. 15, pp. 915-916.

 Several species of *Sphenophorus* injurions to maize-plants in different parts of the United States; habits and ravages of *S. robustus*.
- 2031. [RILEY, C. V.] Effect of drought on the Hessian fly. < Amer. Nat., November [28 October], 1881, v. 15, p. 916. See: < N. E. Farmer, 10 December, 1881. S.-b. No. 45, p. 39. < Farmer's Review, 15 December, 1881. S.-b. No. 45, p. 99.

Hot and dry weather dries up and kills Cecidomyia destructor and its parasites.

- 2032. [RILEY, C. V.] Simulium from Lake Superior. <Amer. Nat., November [28 October], 1881, v. 15, p. 916.
 - Notice of paper by H. A. Hagen; larvæ and pupæ from Lake Superior similar to those of S. pictipes, but the images from the same locality are different.
- 2033. [RILEY, C. V.] Coleopterous cave fauna of Kentucky. < Amer. Nat., November [28 October], 1881, v. 15, pp. 916, 917.
 - Notice of H. G. Hubbard's investigations; Adelops contains but one species, which seems to be equally common in all caves in the State; Ancphthalmus contains several species, and more may be expected to be found.
- 2034. [RILEY, C. V.] Hemipterological studies. <Amer. Nat., November [28 October], 1881, v. 15, p. 917.

 Notice of paper of V. Signoret.
- 2035. [RILEY, C. V.] Entomology in Buffalo, N. Y. <Amer. Nat., November [28 October], 1881, v. 15, p. 917.

 Notice of Vol. 4, No. 1, of the Bulletin of the Buffalo Society of Natural Science; list of entomological papers therein.
- 2036. [RILEY, C. V.] Lampyridæ. <Amer. Nat., November [28 October], 1881, v. 15, p. 917.
 Notice of paper by J. L. Leconte.
- 2037. [RILEY, C. V.] Severe cold and hibernating apple-worms. <Amer. Nat., November [28 October], 1881, v. 15, p. 917.
 - According to A. J. Cook larvæ of Carpocapsa pomonella were killed in their cocoons where exposed to the severe cold of the winter of 1880-1881.
- 2038. RILEY, C. V. [Address delivered 4 November, 1881, at the cotton convention held in Atlanta, Ga., 2-4 November, 1881.] <[U. S. Department of Agriculture.] Address of Hon. George B. Loring... and other proceedings of the cotton convention, Washington, 1881, pp. 19-35. Reprint: <Atlanta Constitution, 5 November, 1881. S.-b. No. 24, p. 34.
 - Beneficial and injurious influence of insects; methods of counteracting the same; ravages and natural history of and search for means against Aletia argillacea [=xylina]; improved methods and contrivances for the application of poisons to plants.

- 2039. RILEY, C. V. The chinch bug. <Amer. Agric., November, 1881, v. 40, p. 476, figs. 1-3. S.-b. No. 42, pp. 21-23.
 - Ravages, food-plants, and natural history of Blissus leucopterus; descriptions and figures of its several stages and of dimorphic form of the imago; effect of wet weather upon it.
- 2040. [RILEY, C. V.] Retarded development in insects. <Amer. Nat., [3] December, 1881, v. 15, pp. 1007-1008. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1881, [13 October], 1882, v. 30, pp. 270, 271. Notice: <Gardener's Chronicle, 27 May, 1882, v. 17, pp. 708-709. S.-b. No. 42, p. 31.
 - Eggs of Caloptenus spretus retained their vitality four and one-half years under abnormal environment and then hatched on exposure to normal conditions; speculations on the cause of the phenomena of retardation of development.
- 2041. [RILEY, C. V.] Preparation of Diptera. Amer. Nat., [3] December, 1881, v. 15, p. 1008.
 Notice of paper of J. Mik.
- 2042. [RILEY, C. V.] Another herbivorous ground-beetle. <Amer., Nat., [3] December, 1881, v. 15, p. 1011.
 - Anisodactylus confusus injuring strawberry-plants in California.
- 2043. [RILEY, C. V.] A disastrous sheep parasite. <Amer. Nat., [3] December, 1881, v. 15, p. 1011.

 A parasite (*Trichodectes ovis?*) doing great injury to sheep in Illinois.
- 2044. [RILEY, C. V.] Locusts in the West. <Amer. Nat., [3] December, 1881, v. 15, p. 1013.
 - Caloptenus spretus scarce in Colorado and Kansas; Camnula pellucida abundant on the Pacific coast; destructiveness of locusts in many parts of South America and of Pachytylus migratorius in Turkey; means adopted against the latter.
- 2045. [RILEY, C. V.] Structure of the claw in Psocina. <Amer. Nat., [3] December, 1881, v. 15, pp. 1013-1014.

 Notice of paper of H. A. Hagen.
- 2046. [RILEY, C. V.] Insect collection for sale. <Amer. Nat., [3] December, 1881, v. 15, p. 1014.

 Notice of the collection of Coleoptera left by C. Trabrandt.
- 2047. RILEY, C. V. Peach-tree bark-borer. Important note from Prof. C. V. Riley. <Rural New Yorker, 24 December, 1881, v. 40, p. 866. S. b. No. 42, p. 13.
 - Habits and ravages of Phlaeotribus liminaris; means against bark-borers.
- 2048. RILEY, C. V. The chinch-bug. <Amer. Agric., December, 1881, v. 40, p. 515, figs. 1-4. S.-b. No. 42, pp. 23-26.
 - Figures Anthocoris [= Triphleps] insidiosus and Harpactor [= Milyas] cinctus as enemies of Blissus leucopterus and Nysius destructor [= angustatus] and Piesma cinerea as likely to be confounded with it; mentions other enemies of the chinch-bug; means against it; importance of irrigation.
- 2049. RILEY, C. V. On the oviposition of *Prodoxus decipiens*. <Amer. Nat., January, 1882 [30 December, 1881], v. 16, pp. 62-63. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1881 [13 October],

- 2049. RILEY, C. V.—Continued. 1882, v. 30, p. 272. Separate: <[Salem, Mass., February, 1882], p. 3.
 - Time and manner of oviposition of *Prodoxus decipiens* in stem of *Yucca filamentosa*.
- 2050. [RILEY, C. V.] Clover insects. <Amer. Nat., January, 1882 [30 December, 1881], v. 16, p. 63.
 Notice of paper of J. A. Lintner.
- 2051. RILEY, C. V. Horn's classification of the *Carabidæ*. < Amer. Nat., January, 1882 [30 December, 1881], v. 16, pp. 63-64. Notice of paper of G. H. Horn.
- 2052. [RILEY, C. V.] The butterfly-trees of Monterey again. Amer Nat., January, 1882 [30 December, 1881], v. 16, p. 64.

 Swarming and migrating habits of Danais archippus; hibernation of the
- 2053. [RILEY, C. V.] Interest felt in economic entomology in California. <Amer. Nat., January, 1882 [30 December, 1881], v. 16, p. 65.
 - Notice of a call issued by the Board of State Horticultural Commissioners of California for a State convention to consider horticultural subjects, including means against insects.
- 2054. [RILEY, C. V.] Obituary. <Amer. Nat., January, 1882 [30 December, 1881], v. 16, p. 65.

 Biographical notice of J. D. Putnam; announcement of the death of G. V. Mniszech.
- 2055. RILEY, C. V. New insects injurious to agriculture. <Amer. Nat., February [25 January], 1882, v. 16, pp. 151–152. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1881, [13 October], 1882, v. 30, pp. 272–273.
 - The occurrence of previously unnoticed noxious insects due to, 1, the introduction of species from abroad; 2, previously existing unnoticed species; and 3, the acquisition of new habits by previously known innoxious species; in the last case the acquisition of new habits may be accompanied by the acquisition of new descriptional character, thus constituting a new species; new species thus become developed within brief periods of time.
- 2056. [RILEY, C. V.] New entomological periodicals. <Amer. Nat., February [25 January], 1882, v. 16, pp. 152–153. Notice of several prospective periodicals.
- 2057. [RILEY, C. V.] Locust probabilities for 1882. <Amer. Nat., February [25 January], 1882, v. 16, p. 153. Favorable prospects of immunity from ravages of Caloptenus spretus in 1882.
- 2058. [RILEY, C. V.] Entomological notes. <Amer. Nat., February [25 January], 1882, v. 16, p. 153.
 - Notice of J. J. Weir's paper on the "Lepidoptera of the Outer Hebrides," etc.
- 2059. [RILEY, C. V.] Bibliography of gall literature. <Amer. Nat., March [24 February], 1882, v. 16, pp. 246-247.
 Notice of the record of F. Thomas.

- 2060. RILEY, C. V. A new depredator infesting wheat-stalks. < Amer. Nat., March [24 February], 1882, v. 16, pp. 247-248, fig. 1.
 - Isosoma allynii French is a species of Eupelmus probably parasitic on some of the wheat-stalk feeders and on some species of Chlorops; habits and figures of larva and description of female imago of Isosoma tritici [p. 247]; comparisons between I. tritici and I. hordei; alleged habits of I. lineare of Europe; relative conspicuousness of the "humeral spot" in European, American, and Australian species of Isosoma. See No. 2063.
- 2061. RILEY, C. V. Further notes on the imported clover-leaf weevil (*Phytonomus punctatus*). <Amer. Nat., March [24 February], 1882; v. 16, pp. 248, 249.
 - Identity of *Phytonomus opimus* Lec. with *P. punctatus* Fab.; instances in which injurious insects have been overlooked for a long time; probability that plant-feeding Coleoptera imported from Europe will not spread far from the Atlantic coast; some species, injurious in Europe, are thus far innoxious in America; hibernation of and means against *P. punctatus*.
- 2062. [RILEY, C. V.] Silk-worm eggs; prices and where obtained. <Amer. Nat., March [24 February], 1882, v. 16, pp. 249–250.
 - Sources from which eggs of Sericaria mori can be obtained in the United States for purposes of silk-culture; prices at which eggs are sold by Crozier & Co.
- 2063. RILEY, C. V. The wheat *Isosoma*. A new depredator infesting wheat-stalks. <Rural New Yorker, 4 March, 1882. S.-b. No. 42, pp. 55-56. See: <Sci. Amer. Suppl., 17 June, 1882. S.-b. No. 37, p. 57.
 - Description of female imago of *Isosoma tritici*; habits, ravages, and distribution of and means against the same; comparison with related species; a probable parasite of it, described as *Isosoma allynii* by G. H. French, belongs to the genus *Eupelmus*. See No. 2060.
- 2064. RILEY, C. V. Possible food-plants for the cotton-worm. < Amer. Nat., April [22 March, 1882], v. 16, pp. 327-329. Reprint: < Rept. [U. S.] Com. Agric. for 1881 and 1882, 1882 [January, 1883], pp. 164-166. Separate: < pp. 164-166.
 - Food-plants of Aletia xylina. See No. 2119 for synopsis of contents.
- 2065. [RILEY, C.V.] Mode of feeding on the larva of *Dytiscus*. < Amer. Nat., April [22 March], 1882, v. 16, p. 330.
 - Notice of paper of E. Burgess; structure of the mouth of the larva of *Dytiscus*; manner in which this larva feeds.
- 2066. [RILEY, C. V.] Entomological notes. <Amer. Nat., April [22 March], 1882, v. 16, p. 330.
 - Brief notices of several articles recently published with items of news.
- 2067. [RILEY, C. V.] Riley's researches. Orange insects and the cottonworm; how the orange insect operates and the cure for its ravages; the hibernation of the cotton-worm; settlement of a mooted question. <Florida Daily Times, 29 March, 1882. S.-b. No. 42, pp. 28–29.
 - Report of interview with C. V. Riley; discovery by W. H. Ashmead that "orange rust" is caused by a mite; habits of and means against the same;

- 2067. RILEY, C. V.—Continued.
 - successful application by H. G. Hubbard of an emulsion of kcrosene and milk as a means against this mite and against scale-insects; proof of the hibernation of Alctia argillacca [=xylina] in the extreme south of the United States; means against the Alctia.
- 2068. RILEY, C. V. Prof. C. V. Riley and the Yucca moth. < Gardener's Mo. and Hortic., March, 1882, v. 24, p. 92. S.-b. No. 23, pp. 198-199.
 - Critical review and reply to editorial of T. Meehan.
- 2069. RILEY, C. V. The *Noctuidæ* in the Missouri entomological reports. < Papilio, March, 1882, v. 2, pp. 41–44. Separate: <N. Y., 1882, 4 pp.
 - Critical review of paper of A. R. Grote; discussion of the synonymy of the *Noctuidæ* in the Missouri entomological reports.
- 2070. RILEY, C. V. Silk-culture in the United States. <Sci. Amer., 1 April, 1882 [v. 60], n. s., v. 46, p. 193. S.-b. No. 42, pp. 93-96.
 - Statement of the dangers to be avoided and the obstacles to be overcome for the successful establishment of silk-culture in the United States.
- 2071. RILEY, C. V. Little known facts about well known animals. A lecture delivered in the National Museum, Washington, D. C., April 8, 1882. <Washington, April, 1882, 32 pp., figs. See: <National Farmer, 13 July, 1882. S.-b. No. 42, p. 15. <Prairie Farmer, 22 July, 1882. S.-b. No. 46, p. 99.
 - The entomological subjects include brief and popular illustrated accounts of the life-histories and transformations of the house-fly (Musca domestica) and mosquito (Culex pipuen s); the development of Trombidium from Astoma; parasitism; truth of the theory of evolution; reasons for the prevalent ignorance of natural history; interest and value of a knowledge of natural history.
- 2072. RILEY, C. V. Lichtenstein's theory as to dimorphic asexual females. <Amer. Nat., May [24 April], 1882, v. 16, p. 409.
 - Criticism of J. Lichtenstein's theory that winged female Aphidide and asexual female Cynipide are larve, and their eggs pupe; errors in the comparison of these forms with the hypermetamorphotic stages of Mcloide.
- 2073. [RILEY, C. V.] Naphthaline cones for the protection of insect collections. <Amer. Nat., May [24 April], 1882, v. 16, pp. 409-410.
 - Use of and objections to the use of naphthaline cones; formula of a preferable protective mixture.
- 2074. [RILEY, C. V.] Injurious insects in California. <Amer. Nat., May [24 April], 1882, v. 16, p. 410.
 - Notice of treatise of Matthew Cooke.
- 2075. [RILEY, C. V.] Sarcophaga lineata destructive to locusts in the Dardanelles. <Amer. Nat., May [24 April], 1882, v. 16, pp. 410-411.
 - Notice of communications, by F. Calvert, regarding the parasitic habits of Sarcophaga lineata, which preys upon Œdipoda cruciata; egg-pods of the same locust destroyed by larvæ of Callostoma fascipenuis.

- 2076. [RILEY, C. V.] Parasitic Diptera. <Amer. Nat., May [24 April], 1882, v. 16, p. 411.
 - Dilophus parasitic in larvæ of Chætoptria hypericana; the Bibionidæ have hitherto been known only as vegetable feeders in the larval state.
- 2077. RILEY, C. V. The cotton-worm. <Times-Democrat [New Orleans], 7 May 1882, v. —, p. 4. S.-b. No. 42, pp. 29–31; 112–113; No. 63, pp. 47–48. Reprint: <Rept. [U. S.] Com. Agric. for 1881 and 1882, 1882 [January, 1883], pp. 157–159. Separate: <p><pp. 97–99.</p>

See No. 2119 for synopsis of contents.

- 2078. RILEY, C. V. The house-fly. <Prairie Farmer, 13 May, 1882. S.-b. No. 39, p. 122. See: <Colman's Rural World, 15 June, 1882. S.-b. No. 68, p. 194. <Times-Democrat [New Orleans], 24 June, 1882. S.-b. No. 42, p. 31. <N. E. Farmer, 5 August, 1882.
 - Wide distribution of Musca domestica; identity of M. harpyia with the species; description of eggs; habits, transformations, and hibernation.
- 2079. RILEY, C. V. Habits of *Cybocephalus*. <Amer. Nat., June [20 May], 1882, v. 16, p. 514.
 - Cybocephalus nigritulus feeds on scales of Chionaspis pinifolii on Pinus elliottii in South Carolina, and C. californicus on a scale on apple-trees in California.
- 2080. [RILEY, C. V.] One effect of the Mississippi floods. < Amer. Nat., June [20 May], 1882, v. 16, pp. 514, 515.
 - Probability that Ligyrus rugiceps will have been drowned out of the region submerged for months by the Mississippi River; liability of the importation of new foes with new seed-plants.
- 2081. [RILEY, C. V.] Doryphora decemlineata in England. < Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - A living specimen carried from North America to London, England, in a barrel of potatoes.
- 2082. [RILEY, C. V.] Dr. Dimmock's inaugural dissertation. < Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - Review of G. Dimmock's paper on "The anatomy of the mouth-parts and of the sucking apparatus of some Diptera."
- 2083. [RILEY, C. V.] The triungulin of *Meloida*. <Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - The triungulin of Meloe proscarabaus was described by J. L. Frisch in 1727.
- 2084. [RILEY, C. V.] Fossil tineids. <Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - Notice of paper of V. T. Chambers; occurrence of the supposed burrow of a tineid larva in a fossil leaf of Acer sp.
- 2085. [RILEY, C. V.] Classification of North American Coleoptera. <Amer. Nat., June [20 May], 1882, v. 16, pp. 515-516.
 - Notice of the approaching completion of the classification of the Coleoptera of North America by J. L. Leconte and G. H. Horn.

- 2086. RILEY, C. V. Hibernation of the army-worm. < Amer. Nat., June [20 May], 1882, v. 16, p. 516.
 - Confirmation of views in regard to the hibernation of the larvæ of *Leucania unipuncta*; places of oviposition; prospects of extensive injury by these insects in the more northern States in 1882.
- 2087. RILEY, C. V. Chinch-bug and army-worm prospects. < Rural New Yorker, 27 May, 1882. S.-b. No. 61, p. 27.
 - Blissus leucopterus abundant in dry weather; Leucania unipuncta in wet weather; the latter hibernates principally as a larva; its eggs secreted in old grass and stubble; means against L. unipuncta.
- 2088. RILEY, C. V. Successful management of the insects most destructive to the orange. <Sci. Amer., 27 May, 1882 [v. 60], n. s., v. 46, pp. 335-336, 5 figs.
 - Treats of Coccida injurious to the orange; means against the same; use of kerosene emulsion; figures stages of Mytilaspis pomicorticis [=pomorum], M. gloveri, and M. citricola.
- 2089. RILEY, C. V. The utilization of ants in horticulture. <Nature, 8 June, 1882, v. 26, p. 126. Reprint: <Gardener's Chronicle, 17 June, 1882, v. 17, p. 805. S.-b. No. 42, p. 32.
 - Abstract of paper of C. J. Macgowan; capture and sale of two species of ants which build nests in trees; colonization of these ants in orange orchards to destroy injurious insects.
- 2090. RILEY, C. V. The army-worm vs. the clover hay-worm. Remedies for the army-worm. <Rural New Yorker, 10 June, 1882.
 S.-b. No. 47, pp. 158-159; No. 61, pp. 32-33. Reprint: <Nat. Farmer, 22 June, 1882.
 S.-b. No. 37, p. 123. <Lancaster Farmer, July, 1882.
 S.-b. No. 46, p. 104. <Home and Farm, 1 July, 1882.
 S.-b. No. 37, p. 122.
 - Abundance of Leucania unipuncta and Asopia costalis in Alabama and Tennessee; confusion of the army-worm with the clover hay-worm; prevalence of enemies of and means against the former.
- 2091. RILEY, C. V. Repelling insects by malodorants. <Amer. Nat., July [22 June], 1882, v. 16, p. 596.
 - Critical review of paper of J. A. Lintner; odorous substances repel insects more by their toxic properties than by their odor; failure of attempts by the use of strongly smelling substances to prevent oviposition; sight, touch, and taste generally more important in insect economy than smell.
- 2092. [RILEY, C. V.] Habits of *Bittacus apterus*. <Amer. Nat., July [22 June], 1882, v. 16, pp. 596–597.
 - Notice of paper of C. R. Osten Sacken; locomotion and food-habits of Bittacus apterus in California.
- 2093. RILEY, C. V. Habits of Coscinoptera dominicana. <Amer. Nat. July [22 June], 1882, v. 16, p. 598.
 - Discovery by F. H. King that Coscinoptera dominicana is inquilinous in ants' nests in its earlier stages; similar habits of related species.
- 2094. [RILEY, C. V.] Sun-spots and insect life. <Amer. Nat., July [22 June], 1882, v. 16, pp. 598–599.
 - Review of paper of A. H. Swinton; dependence of certain insect phenomena on the periods of sun-spots, through the determination by the latter of meteorological conditions.

- 2095. RILEY, C. V. The silk-worm. No. 1-3. < National Farmer, 13, 20, 27 July, 1882. S.-b. No. 35, p. 15; No. 48, pp. 32, 49.
 - Nature of the silk-worm; stages, enemies, diseases, races; wintering and hatching of eggs; feeding and rearing of larvæ; the cocoons; spinning and gathering, choking chrysalids; egg-laying, reproduction, reeling, foodplants.
- 2096. RILEY, C. V. Change of habit; two new enemies of the eggplant. <Amer. Nat., August [28 July], 1882, v. 16, pp. 678-679.
 - Sudden acquisition by Doryphora juncta and by Cassida texana of the habits of feeding on Solanum melongena, these species having been found previously on S. carolinense and S. elwagnifolium respectively; occurrence of C. texana on S. carolinense; distribution of the two insects.
- 2097. RILEY, C. V. Notes on *Microgasters*. <Amer. Nat., August [28 July], 1882, v. 16, pp. 679-680.
 - Critical review of paper of A. S. Packard; synomymical notes on Packard's species.
- 2098. [RILEY, C. V.] Are honey-bees carnivorous? <Amer. Nat., August [28 July], 1882, v. 16, p. 681.
 - Observations of Fritz Müller on the carnivorous habits of Brazilian Apidæ.
- 2099. [RILEY, C. V.] The "overflow bugs" in California. < Amer. Nat., August [28 July], 1882, v. 16, pp. 681-682.
 - Extract from letter of Mrs. A. E. Bush, with introductory remarks; occurrence of *Platynus maculicollis* in such abundance as to become a nuisance.
- 2100. [RILEY, C. V.] Insects and drouth. < Amer. Nat., September [24 August], 1882, v. 16, p. 745.
 - Occurrence of excessive drouth in the spring and early summer of 1880 in New England; statement by S. Lockwood of the exceptional abundance of several insects in New Jersey in that year.
- 2101. RILEY, C. V. Probable sound organs in sphingid pupæ. < Amer. Nat., September [24 August], 1882, v. 16, pp. 745-746.
 - Occurrence of a peculiar structure on the abdominal joints of the pupæ of certain Sphingidæ; genera in which this structure is observed; probable connection of this structure with the function of producing sound as observed in Sphinx atropos.
- 2102. RILEY, C. V. Is *Cyrtoneura* a parasite or a scavenger? < Amer. Nat., September [24 August], 1882, v. 16, pp. 746-747.
 - Cyrtoneura stabulans bred from pupæ of Aletia argillacea [=xylina] usual food of this species; doubt whether the species is a parasite or is only a scavenger in decayed pupæ; Phora aletia merely a scavenger; great liability of pupæ of Aletia to decay
- 2103. [RILEY, C. V.] Habits of Polycaon confertus Lec. < Amer. Nat., September [24 August], 1882, v. 16, p. 747.
 - Polycaon confertus bores in twigs of apple- and pear-trees and grape-vines; the larvæ probably live in the dead and dry wood of forest-trees.
- 2104. RILEY, C. V. Dinoderus pusillus as a museum pest. < Amer. Nat., September [24 August], 1882, v. 16, p. 747.
 - Dinoderus pusillus feeds on cork and paper lining in an insect box; they occur usually in drugs and other stored and dry vegetal products.

- 2105. RILEY, C. V. Myrmecophilous Coleoptera. <Amer. Nat., September [24 August], 1882, v. 16, pp. 747-748.
 - Larvæ and images of Euphoria hirtipes live in hills of Formica rufa; pupæ of Hymenorus rufipes in nests of Formica fusca and of H. obscurus in nests of another species of ant; known myrmecophilous habits of Cetonia, Cremastochilus, Euparia castanea (in nests of Solenopsis xyloni [= geminata]), Tenebrionidæ, and Anthicus.
- 2106. [RILEY, C. V.] Discontinuance of publication. <Amer. Nat., September [24 August], 1882, v. 16, p. 748.

Announcement of the discontinuance of publication of "Revue Coléoptérologique;" critical review of the same.

- 2107. RILEY, C. V. Buffalo tree-hopper injurious to potatoes. < Amer. Nat., October [28 September], 1882, v. 15, p. 823. Habits, food-plants, and ravages of Ceresa bubalus.
- 2108. [RILEY, C. V.] Moths attracted by falling water. < Amer.
 Nat., October [28 September], 1882, v. 16, p. 826.
 Notice of paper of J. S. Gardner; gleaming water-falls in Iceland as attractive to moths as artificial light would be.
- 2109. [RILEY, C. V.] A new museum pest. <Amer. Nat., October [28 September], 1882, v. 16, p. 826. Reprint: <Psyche, September-October, 1882 [1 March, 1884], v. 3, p. 408.

 *Perimegatoma variegatum as a pest in collections of insects.
- 2110. [RILEY, C. V.] Fleas feeding on lepidopterous larvæ. Amer.
 Nat., October [28 September], 1882, v. 16, p. 826.
 Notice of paper of C. J. Boden; abundance of fleas (*Pulex sp.*) where few or no warm-blooded animals occur may be explained by the feeding of the fleas on insects.
- 2111. [RILEY, C. V.] [Gasteracantha cancer.] < Gonzales [Tex.] Inquirer, 30 September, 1882. S.-b. No. 42, p. 15.

 Habits of Gasteracantha cancer; the male unknown.
- 2112. RILEY, C. V. Cicada septendecim. < Gardener's Mo. and Hortic., September, 1882, v. 24, pp. 274-275. S.-b. No. 38, p. 42; No. 39, p. 6; No. 42, pp. 2-4.
 - Orthography of the names C. [= Tibicen] septendecim and C. [= T.] tredecim; natural relations of the same; indistinguishability of the species of certain genera; dimorphic forms more numerous than usually recognized; $Massospora\ cicadina\ parasitic\ on\ Cicada.$
- 2113. RILEY, C. V. Remarkable felting caused by a beetle. <Rural New-Yorker, 14 October, 1882, v. 41, pp. 699-700. S.-b. No. 42, p. 16. Reprint, with changes: <Amer. Nat. [2] December, 1882, v. 16, pp. 1018-1019.
 - Description of the felting of the interior of a pillow-ticking with fragments of feathers formed by the ravages of Attagenus megatoma within a feather pillow.
- 2114. RILEY, C. V. The buckeye-leaf stem-borer. < Amer. Nat., November [28 October], 1882, v. 16, pp. 913-914. See: < Sci. Amer. Suppl., 16 December, 1882. S.-b. No. 47, p. 145.
 - Sericoris instrutana †Claypole = S. [= Steganoptycha] claypoleana n. sp.; habits and food-plants of this species and of Proteoteras asculana; distinctions between these two species.

- 2115. [RILEY, C. V.] Efficacy of chalcid egg-parasites. <Amer. Nat., November [28 October], 1882, v. 16, pp. 914-915. See: <Prairie Farmer, 2 December, 1882. S.-b. No. 60, p. 123.
 - Beneficial influence and occasional vast abundance of egg-parasitic Chalcididæ and especially of Trichogramma pretiosa and of Telenomus sp.
- 2116. [RILEY, C. V.] On the biology of Gonatopus pilosus Thoms. <Amer. Nat., November [28 October], 1882, v. 16, p. 915.
 - Notice of paper of J. Mik; parasitism of Gonatopus pedestris on Athysanus maritima, and of G. pilosus on Deltocephalus xanthoneurus; transformations and habits of G. pilosus; probable parasitism of a Gonatopus (G. contortulus?) on Amphiscepa bivittata.
- 2117. RILEY, C. V. Species of Otiorhynchidæ injurious to cultivated plants. <Amer. Nat., November [28 October], 1882, v. 16, pp. 915-916. See: <Sci. Amer. Suppl., 16 December, 1882. S.-b. No. 47, p. 145.
 - Aramigus fulleri the only species of North American Otiorhynchidæ whose development and earlier stages are known; food-plants of several species; discovery by G. P. Peffer of the injuries of Anametis grisea upon roots of apple- and pear-trees.
- 2118. [RILEY, C. V.] Bombyliid larvæ destroying locust eggs in Asia Minor. <Amer. Nat., November [28 October], 1882, v. 16, pp. 916-917.
 - Notice of and extract from communications of F. Calvert; parasitic habits of Callostoma fascipennis; similarity of habits of Bombyliidæ in Asia and in North America; probability that the larvæ of Cantharis vesicatoria and of other Meloidæ will be found to feed on eggs of Aerididæ.

1

2119. RILEY, C. V. Report of the Entomologist. <Ann. Rept. [U. S.]
Commissioner Agric. for 1881 and 1882, 1882 [January, 1883],
pp. 61-214, 20 pl. Separate: <Washington, 1 December, 1882,
pp. 8+167, 20 pl.

CONTENTS.

Plan of the report, 1—Insufficiency of former appropriations to the Entomological Division, 1—Reorganization of the Division, 1—Great increase in the correspondence of the Division, 2—Preparation of special bulletins, 2—Periodical bulletins desirable, 2—Special reports in preparation, 2—Work of the United States Entomological Commission, 3—The United States National Museum in connection with the Entomological Division, 3—Assistants, agents, and observers of the Division, 3—The illustrations to the report, 4.

EXTRACTS FROM CORRESPONDENCE

Miscellaneous correspondence on various insects injurious to vegetation in 1881-'82, 4.

Distribution of eggs by the Department, 7—Mr. L. S. Crozier's opinion as to the value of American silk products, 7—Mr. E. Fasnach on the importance of a home market for silk products, 8—Associations, 8—Ladics' Association at Spring Hill, Ala., 8—Women's Silk Culture Association of California, 8—Women's Silk Culture Association of the United States, 8—American Silk Exchange in New York, 8—Sales of eggs and cocoons in the

16

27

29

46

2119. RILEY, C. V.—Continued.

SILK CULTURE-Continued.

United States, 9—Various reports from correspondents of the division, 9—Experience in 1882 at the Department in raising worms fed on osage orange or mulberry, 11—New mulberry tree from southern Russia, 13—Business ventures in the United States in connection with silk-producing industry, 13—Salé of eggs, 13—Second edition of the manual of silk culture, 14—Preface thereof, 14—Reasons why the growing interest in silk culture in the United States should be checked at the present time, 15—Import duty on raw silk demanded, 16.

PYRETHRUM: ITS USE AS AN INSECTICIDE Circular in reference to Pyrethrum, 16—Distribution of seed by the Department, 16-History of Pyrethrum in Asia and Dalmatia, 17—Experience in growing Pyrethrum roseum in America, 17— Soil best adapted for raising the plant, 18—The Dalmatian plant in California, 18-Experience with raising the plant at Washington, 18-Preparation of the plant for use, 18-Gathering and drying the flowers, 18-Pulverizing, 19-Keeping the powder, 19-Advantages and disadvantages of the powder as an insecticide, 19—How Pyrethrum affects different insects, 19—Modes of application, 19-Pyrethrum as dry powder, 19-Pyrethrum applied in fumes, 20-Alcoholic extract, 20-The powder stirred up in water, 20—Tea or decoction, 21—Reports from persons to whom Pyrethrum seed was sent, 21—Success in raising the plant at Washington, 24-Miss M. E. Murtfeldt's experiments with powder on various insects, 25.

The army worm, Leucania unipuncta..... Habits and natural history, 30—History of investigation in America, 30—Where the eggs are laid, 30—When the eggs are laid, 31— Mode of oviposition, 31—Prolificacy of the moth, 32—Duration of the egg state, 32-Habits of the young worms, 32-Guenée's characterization of Lencania larvæ, 32-Marching of the worms abnormal, 33-The marching worm not a distinct race, 33-Duration of worm life, 34-Remedies, 34-Burning old grass, 34-Predictions of army-worm years unreliable, 35-Mcteorological influences on the species, 35-Ditching and fencing as remedies, 36—Use of arsenical poisons, 36—Kerosene and coal tar, 36— Heavy rolling not to be recommended, 37—"Drawing the rope" of little avail against the worm, 37-Mr. L. O. Howard's report on army-worm injury, 1881, in Illinois and Indiana, 37—Reports from correspondents, spring of 1882, 39-Rev. Samuel Lockwood's account of the invasion of 1880, in New Jersey, 44.

Characterization of the species, 46—Comparative destructiveness of the three species treated of, 47—Periods of migration, growth, and incubation, 47—Scale insects easily assailable only during a brief period of their development, 48—The hatching process and its maximum periods, 48—Great vitality of the eggs, 49—The work of enemies and parasites, 49—Lady-bugs, lace-wings, and lepidopterous enemies, 48—Importance of mites as external

2119. RILEY, C. V.—Continued.

SCALE INSECTS OF THE ORANGE—Continued.

enemies of scale insects, 49—Hymenopterous parasites the most efficient destroyers, 49—Sudden outbursts of the pcst, 50—Destruction of the scales by their own undue increase, 51—Injury inflicted on young trees, and faulty methods of destroying the scales, 51—Recapitulation, 51—Remedies, 52—Danger in applying undiluted volatile oils, 52—Kerosene, when properly applied, not injurious to the tree, 52—Process of preparing kerosene butter, 52—Nature of kerosene butter, 53—How it should be applied, 54—The kerosene butter a sure remedy, 54—How it affects the scales and the eggs, 54—The aquapult force-pump very convenient for applying kerosene emulsions, 54—Effect of kerosene emulsion on the tree, 55—Kerosene emulsions experimented with, 55—Whale oil soap as remedy, 56—Oil of creosote, its effect on scale insects and on the trees, 56—Saponaceous compounds of creosote, 57—Creosote inferior to kerosene as insecticide, 57—Various other substances as remedies, 57—Tables of experiments, 60—Note on more recent discoveries in producing kerosene emulsions, by C. V. Riley, 67.

cide, 57—Various other substances as remedies, 57—Tables of	
experiments, 60—Note on more recent discoveries in producing	
kerosene emulsions, by C. V. Riley, 67.	
Insects affecting the rice-plant	6
Rice production in the United States in 1879, 67.	
The rice-grub, Chalepus trachypygus	68
Mr. L. O. Howard's observations on the occurrence and habits of	
the insect, 68—It can easily be kept in check, 68—Structural char-	
acters and geographical distribution of the genus Chalepus, 69—	
Description of imago and larva, 69.	
The water-weevil, Lissorhoptrus simplex	70
Colonel Screven on the specific identity of "the maggot" and "the	
water-weevil," 70-Mr. L. O. Howard's report on the occurrence	
of larva and beetle in rice-fields, 70—Habits of the beetle, 71—	
Remarkable shape of the larva, 71—Characters of the genus Lis-	
sorhoptrus, 72—Description of the imago, 72—Description of the	
larva, 72.	
The rice-stalk borer, Chilo oryzæellus	7
Mr. Howard's account of the work of the larva and the damage	
caused by it, 73—Enemies, 74—Preventive measures, 74—Difficul-	
ties in the specific determination of the insect, 74—Characteris-	
tics of the genus Chilo, 75—Description of the moth, 75—Descrip-	
tion of larva and pupa, 75.	
White blast	70
Colonel Screven's account of the disease, 76—Mr. Howard's observa-	
tions on the insects found on the diseased plants, 77—The blast	
possibly caused by insect work, 77.	
Other insects injurious to growing rice	78
The "grass-worm" and its injury, 78—The "Inbber grasshopper,"	
78—Acridium obscurum and various Heteroptera, 78.	
Insects affecting corn or maize	78
The corn-bill bug, Sphenophorus robustus	78
Species of Sphenophorous injurious to agriculture, 78—Nature of	
damage done by the beetles, 79—Mr. Howard's report on the work	
of the larva and its development, 79—Preventive measures, 80—	
Burning the stubble in winter time to be recommended, 80—Gen-	

eric characters of Sphenophorus and its natural groups, 80—Diagnosis of the imago, 81—Description of the larva, 81—Larva of

	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	295
2119.	RILEY. C. V.—Continued.	
	INSECTS AFFECTING CORN OR MAIZE—Continued.	
	Rhodobænus 13-punctatus and Rhynchophorus zimmermanni, 82—	
	Description of the pupa, 82.	
	The smaller corn-stalk borer, Pempelia lignosella	82
	First appearance of the pest, 82—Geographical distribution in the	0.2
	United States, 83—Habits and natural history, 83—Work of the	
	larva, 83—Transformation, 83—Habits of the moth, 83—Prevent	
	ive measures, 84—Color variations of the moth, 84—Mouth parts	
	of the moth, 84—Description of the larva, 84—Description of the	•
	pupa, 85.	
	The boll-worm, alias corn-worm, Heliothis armigera	85
	Injury to corn in 1881, 85—Food-plants, 85—Identity of the corn-	00
	worm and boll-worm, 85—Number of broods in the northern	
	States, 86—Nature of damage done to corn, 86—The worm also	
	feeding on hard corn, 87—Corn-feeding broods in the southern	
	States, 87—Damage done to tomatoes, 88—Food-plants of the	
	families Solanaceæ and Leguminosæ, 88-Cucurbitaceous and	
	malvaceous food-plants, 89—Various other food-plants, 89—List	
	of food-plants probably still incomplete, 89—Carnivorous habit	
	of the worm, 89—Report upon Heliothis armigera, by Judge Law-	
	rence Johnson, 90.	
	THE COTTON-WORM, Aletia xylina	92
	Address by C. V. Riley before the Atlanta cotton convention, 93—	
	Past work on the practical side of the cotton-worm question, 93—	
	Obstacles and difficulties of the investigation, 93—Natural his-	
	tory, 94—The insect in its different stages, 94—Time of develop-	
	ment and first appearance of the worms, 94—Poisons best applied	
	from beneath, 95—Improvements in machinery for applying poi-	
	sons, 95—Nozzles of various forms, 96—Machinery for poisoning	
	from below, 96.	0~
	Protection from injury in regions overflowed by the Mississippi	97
	Poisons should be applied to the under surface of the leaves, 98—	
	Paris green and London purple, 98—Sprinkling and sifting, 98—	
	Device for mixing poisons and diluents, 98—Arsenic and arseniate of soda, 98—Early poisoning, 99.	
	Poisoning devices—machine for spraying from below	99
	The skid, 99—Device for thorough mixing and filling the barrel,	70
	100—Details of barrel, pump, and stirrer, 100—Inserting and ex-	
	tricating the stirrer-bar, 100—Fork of the descending pipe, 100—	
	The nozzles, 102.	
	Damage in 1881	102
	Loss of cotton by worms by States and counties, 102 — Loss of	
	cotton by worms in 1881, tabulated, 104.	
	Possible food plants of the cotton-worm	105
	No other food-plant than cotton yet found, 104—Eggs of Anomis	
	erosa found on Urena lobata, 104—Eggs and young larvæ of Anomis	
	erosa distinguished from those of Aletia xylina, 105—Plants upon	
	which eggs related to Aletia were found in the Department her-	
	barium, 105-Localities for malvaceous plants, with possible	
	food-plants of Aletia indicated, 105.	
	The question of hibernation settled	106

Theories of hibernation, 106—Eggs deposited early in March, 106— The early brood of worms, 106-Where the moth hibernates,

106-Value of fall and winter preventive work, 106.

MISCELLANEOUS INSECTS	10
The Urena anomis, Anomis erosa	10
Habits and natural history; resemblance to the cotton-worm, 107-	
Distribution, 107—Its eggs mistaken for those of Aletia, 108—Dis-	
tinguishing characters, 108—Habits of larvæ, 108—Characters of	
moth, 109—Descriptive, 109.	
The clover leaf-beetle, Phytonomus punctatus	11
Habits of the genns Phytonomus, 111—Ph. punctatus au Enropean	
insect, 111—Food-plants of European species, 111—History of Ph.	
punctatus in N. A., 111—Specific identity of Ph. punctatus and Ph.	
opimus, 112—When first noticed as destructive, 112—Damage in	
Yates County, N. Y., in 1881, 112—Mr. Schwarz's observations in	
Jnne, 1882, 113—Life history of the species, 113—The eggs, 113—	
The newly-hatched larvæ, 113—Method of locomotion of larvæ,	
114—The cocoon, 114—Method of spinning, 115—Its method of	
spinning differs from that alleged of the European Ph. rumicis,	
115—Principal damage done in the beetle state, 115—Feeds on	
white and red clover, 115—Length of life in the different states,	
115—Number of annual broods, 116—A second brood exceptional,	
116—Remedies, 117—Will it spread?, 117—Natural enemies, 117—	
Description of earlier states, 118. The vagabond Crambus, Crambus vulgivagellus	119
History of its injury and identification, 119—Correspondence with	11:
Professor Lintner, 119—Abundance of the moths in the eastern	
States in 1881, 119—Habits and natural history, 119—The eggs,	
120—The larva and its tube, 120—Habits of the moth, 120—Dam-	
age done by the worms, 120—Parasites, 121—Similarity of habit	
in an Enropean species, 121—Remedies, 121—Descriptive, 121—	,
Specimens from Vancouver's Island compared, 121—Descriptions	
of earlier states, 122 - Bibliographical list, 122.	
The wheat Isosoma, Isosoma tritici	123
Past history and habits, 123—When first studied, 123—Professor	
Thomas finds it in Illiuois and Dr. Packard in Virginia, 123—Its	
work differs from that of the joint-worm, 123—Its injuries in	
Missouri, 124—Professor French's observations, 124—Comparisons	
with the joint-worm and other allied species, 124—Errors cor-	
rected, 125—Its relationship to the European I. linearc; habits	
of latter, 125—Number of broods, 125—Remedies, 125—Parasites,	
126—Descriptive, 126—Bibliographical list, 126.	104
The sorghum web-worm, Nola sorghiella.	127
Its injuries, 127—Letter from J. P. Stelle, 127—Habits, 128—Method	
of work, 128—Systematic position, 128—Descriptive, 128—Earlier stages, 129.	
The catalpa sphinx, Ceratomia catalpa	129
Value of the catalpa tree, 129—Its usual exemption from insect	120
attacks, 129—Past history of Sphinx catalpa, 129—Its larvæ used	
as fish bait, 130—Its distribution, 130—Extracts from correspond-	
ence concerning the larva, 130—Characters and natural history:	
The eggs, 131—The larvæ, 131—The pupa, 131—Description of	
the moth, 131—Number of broods, 132—Remedies, 132—Descrip-	
tive, 132—Pecnliarity of structure of pnpa, 133.	
The osage orange sphinx, Ceratomia hageni	133
Value of the osage orange, 133—Its comparative exemption from	
insect attack, 133—Rarity of the osage orange sphinx, 133—Its	
generic place, 133—Characters of the species, 134—Its affinities,	

211	19. RILEY, C. V.—Continued.	
	REPORT ON MISCELLANEOUS INSECTS, BY PROF. J. HENRY COMSTOCK.	135
	The apple-maggot, Trypeta pomonella	135
	Its importance compared with the codlin moth, 135—Method of work, 135—Food-plants, 135—Its spread, 136—Extracts from cor-	
,	respondence, 136—Early apples principally infested, 136—Tech-	
	nical descriptions of different states, 137—Remedies, 138—Manner	
	of distinguishing it from codlin-moth larva, 138.	
	The vine-loving pomace-fly, Drosophila ampelophila	13 8
	Reasons for popular name, 139—Reasons for treating of the spe-	
	cies, 139—Where found, 139—Rapidity of multiplication, 139— Destructive to grapes, 139—Technical descriptions of different	
	stages, 140—Remedies, 141.	
	The pretty pomace-fly, Drosophila amana	141
	Technical descriptions of different stages, 141—Remedies, 142.	
	The ocellate leaf-gall of the red-maple, Sciara ocellaris	142
	Distribution, 142—Description of gall, 142—Description of larva,	
	143—Its cocoon, 143—Number of generations, 143—Gall described as cecidomyid by Osten-Sacken, 143—Another gall-making	
	Sciara, 143—Description of adult male, 144.	
	Ladybirds, Coccinellida	144
	General remarks on habits, 144—The ashy-gray ladybird, 144—De-	
	scription of larva, 144—Description of pupa, 145—The adult, 145—	
	The blood-red ladybird, 145—Description of pupa, 145—The lady- bird of the cactus, 145—Descriptions of larva and pnpa, 145—The	
	ambiguous Hippodamia, 146—Descriptions of larva and pupa,	
	146—Other species observed at Los Angeles, Cal., 146.	
	Methods of destroying scale insects	146
	Note by the Entomologist, dissenting from the conclusions of this	
	article, 146—Letter from S. F. Chapin, 147—Letter from Matthew Cooke, 148—Method of spraying trees in groves, 148—Mr. Chapin's	
	contrivance, 148.	
	Lac insects, Carteria.	149
	General remarks, 149—The genus Carteria of Signoret, 49—Carteria	
	lacca (Kerr); general appearance, 149—Preparation for study, 150—Technical description, 150—Remarks on Mr. Carter's descrip-	
	tion, 151—Carteria larrew n. sp., 151—Remarks on the creosote	
	plant, 151—Mr. J. M. Stillman's paper, 151—Differs from C. lacca,	
	151—Technical description of C. larrew, 152—Carteria mexicana n.	
	sp., 152—Where found, 152—General appearance, 152—Technical	
	description, 152. A new wax insect, Cerococcus quercus	153
	General description, 153—Can it be utilized?, 153—Cerococcus new	100
	genus, 153—Cerococcus quercus n. sp., 153—Female sac, 153—Fe-	
	male, 154—Male sac, 154.	
	Note on the structure of mealy bugs	154
	The anal and genital openings distinct, 154—Two pairs of openings homologous with the honey tubes of Aphididæ, 154.	
21:	20. [RILEY, C.V.] A new rice stalk-borer: Genus-grinding. <at< td=""><td>ner.</td></at<>	ner.
	Nat., [2] December, 1882, v. 16, pp. 1014-1015.	
	Extract from Rept. [U. S.] Commis. Agric. for 1881 and 1882, pp. 134	
	with additional remarks; Chilo oryzwellus = Diphryx prolatella; Dip Grote founded on a mutilated specimen, with mistake of maxillary	
	labial palpi.	

- 2121. [RILEY, C.V.] A butterfly larva injurious to pine-trees. < Amer. Nat., [2] December, 1882, v. 16, pp. 1015–1016.
 - Habits and ravages of *Pieris menapia* on yellow-pine and tamarack in Washington Territory.
- 2122. [RILEY, C. V.] The army-worm in 1882. <Amer. Nat., [2] December, 1882, v. 16, p. 1017.
 - Occurrence of Leucania unipuncta in great abundance and with disastrous effects, especially in southern United States in 1882.
- 2123. [RILEY, C. V.] The wheat-stalk worm on the Pacific coast. <Amer. Nat., [2] December, 1882, v. 16, pp. 1017–1018.
 - Isosoma tritici injuring wheat-stalks in Washington Territory; reference to prior notices of this insect.
- 2124. [RILEY, C. V.] Deserved honor. <Amer. Nat., [2] December, 1882, v. 16, p. 1018.
 - Notice of the appointment of Eleanor A. Ormerod to be consulting entomologist to the Royal Agricultural Society of Great Britain.
- 2125. [RILEY, C. V.] Important work on *Cynipidæ*. <Amer. Nat., [2] December, 1882, v. 16, p. 1018.
 - Notice of G. Mayr's "Die europäischen Arten der gallenbewohnenden Cynipiden."
- 2126. RILEY, C. V. Emulsions of petroleum and their value as insecticides. <Rural New Yorker, 9 December, 1882, v. 41, pp. 833, 834. S.-b. No. 42, pp. 17-18.
 - Results of experiments made upon methods of using petroleum as an insecticide without injury to plants; soap and milk emulsions the most available; methods of preparing the same.
- 2127. RILEY, C. V. The bean-weevil. <Rural New Yorker, 9 December, 1882, v. 41, p. 835. S.-b. No. 42, p. 18.
 - Answer to inquiry of R. J. B.; habits of and means against Bruchus fabæ.
- 2128. RILEY, C. V. The "cluster-fly." < Prairie Farmer, 23 December, 1882, v. 54, p. 7. S.-b. No. 42, pp. 16–17. Reprint: < Amer. Nat., [5] January, 1883, v. 17, pp. 82–83.
 - Habits and synonymy of *Pollenia rudis*; notice of other accounts of the swarming of Diptera. See No. 2174.
- 2129. RILEY, C. V. Darwin's work in entomology. <Proc. Biol. Soc. Wash., 1882, v. 1, pp. 70–80.
 - Analysis of the interest shown by C. R. Darwin in entomology and of his contributions to the same.
- 2130. RILEY, C. V. The cotton-worm. < Western Farmer's Almanac for 1883, 1882, p. 40. S. b. No. 42, p. 71.
 - Various theories hitherto held in regard to the hibernation of Aletia argillacea [=xylina]; proof of its hibernation; seasons and conditions of development of the first broad of the year, precautionary measures to be adopted.
- 2131. RILEY, C. V. Pyretbrum, an important insecticide. <Western Farmer's Almanac for 1883, 1882, pp. 41–42. S.-b. No. 42, p. 49. See: <Prairie Farmer, 27 January, 1883. S.-b. No. 42, pp. 56–58.
 - Condensed account of the history of the use of pyrethrum flowers as an insecticide; cultivation of the plants and preparation of the powder; methods of its application; experiments in the cultivation of the plants.

- 2132. [RILEY, C. V.] New lists of North American Lepidoptera. <Amer. Nat., [5] January, 1883, v. 17, pp. 80–82.
 - Reviews of lists of Brooklyn Entomological Society, of C. H. Fernald and A. R. Grote.
- 2133. [RILEY, C. V.] Naphthaline cones. <Amer. Nat., [5] January, 1883, v. 17, pp. 83, 84.
 - The cones stain the paper lining of boxes; they seem to destroy mites and Psoci very soon, but have little effect on Dermestidæ.
- 2134. RILEY, C. V. Emulsions of petroleum as insecticides. <Sci. Amer., 6 January, 1883 [v. 62], n. s., v. 48, p. 3. S.-b., No. 42, pp. 4-7.
 - Notice of experiments made, under author's direction, in 1882, in the use of emulsions of kerosene oil; report of H. G. Hubbard upon experiments made by him; eritical review of S. F. Chapin's "Seale insects on deciduous and ornamental trees;" effect of pure kerosene, of emulsions, and of lye upon trees.
- 2135. RILEY, C. V. Entomological notes. <Rural New Yorker, 13 January, 1883. S.-b. No. 42, p. 78. See: <Amer. Nat., 1883, v. 17, pp. 198–199.
 - 1. A new enemy to wax-beans; extract from letter of G. H. Stone, on the food-plants, habits, and ravages of *Epilachna corrupta*; distribution of the same. 2. Spread of the 12-punctured asparagus beetle; increasing noxiousness of *Crioceris* 12-punctata recorded from near Baltimore, Md., by O. Lugger; comparative description of this species with *C. asparagi.* 3. An internal mite in fowls; presence of *Cytoleichus sarcoptoides* in lungs and other parts of diseased chickens.
- 2136. [RILEY, C. V.] The "lignified snake of Brazil." < Evening Star [Washington, D. C.], 20 January, 1883, v. 61, p. 2. S.-b. No. 42, pp. 59-60. Reprint: < Sci. Amer. Suppl., 17 February, 1883. See: < Science, 23 February, 1883, v. 1, p. 84.
 - Discussion of a specimen of problematical character, supposed to be the burrow of a larva under bark; notice of writings upon the subject; frequency with which the true nature of natural objects is mistaken; letter from J. H. Hutchins accompanying a gall of Cecidomyia vitis-pomum mistaken for a hybrid fruit.
- 2137. RILEY, C. V. Utilization of ants in horticulture. <Sci. Amer., 27 January, 1883 [v. 62], n. s., v. 48, p. 49. S.-b., No. 42, pp. 65-66.
 - Abstracts of papers of C. J. Maegowan and H. C. McCook, with additional notes; the introduction of ants might involve objectionable consequences; probability that they would not be of service against *Coccida*.
- 2138. RILEY, C. V. Natural sugaring. <Amer. Nat., February [31 January], 1883, v. 17, pp. 197–198. Reprint: <Country Gentl., 31 May, 1883, v. 48. S.-b. No. 42a, p. 297.
 - Lachnus platanicola n. sp. [p. 198], abundant in 1882 on syeamore trees; description of the species; attraction of great numbers of insects to its saccharine exudations, and growth of Funago salicina upon these exudations.
- 2139. [RILEY, C. V.] Trogoderma tarsale as a museum pest. < Amer. Nat., February [31 January], 1883, v. 17, p. 199.
 - Notice of paper of F. H. Snow; remarks on the abundance and ravages of Trogoderma tarsale; habits of its larva.

2140. [RILEY, C. V.] Phylloxera in California. <Amer. Nat., February [31 January], 1883, v. 17, pp. 199-200.

Phylloxera vastatrix in California is most injurious in moist soils.

2141. RILEY, C. V. The hibernation of Aletia xylina [Say] in the United States a settled fact. <Sci. Amer., 3 February, 1883, v. 48, p. 68. S.-b. No. 42, pp. 66-67. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1882, 1883, v. 31, pp. 468-469. Separate: <Salem, Mass., July, 1883, pp. 468-469. Abstract: <Amer. Nat., April [15 March], 1883, v. 17, pp. 420-421. <Nature, 28 December, 1882, v. 27, p. 214.

Proof of the hibernation of Aletia xylina as a moth and of the perpetual existence of the species in Florida.

- 2142. [RILEY, C. V.] Fostering the study of economic entomology. <Rural New-Yorker, 12 February, 1883. S.-b. No. 42, p. 58. Reprint: <Amer. Nat., April [15 March], 1883, v. 17, p. 420. Efforts of the French for the promotion of economic eutomology.
- 2143. [RILEY, C. V.] Diseases of the chinch-bug. <Rural New-Yorker, 17 February, 1883. S.-b. No. 42, p. 58.

 Note concerning paper of S. A. Forbes on two fungus parasites of Blissus leucopterus.
- 2144. R[ILEY], C. V. [Instinct of Cicada [= Tibicen] septendecim.]

 Amer. Nat., March [21 February], 1883, v. 17, p. 322.

 Remarks on note of E. W. Claypole; sense of direction in insects.
- 2145. [RILEY, C. V.] Food-habits of Megilla maculata. <Amer. Nat., March [21 February], 1883, v. 17, pp. 322-323.

 Summary of S. A. Forbes' observations upon the food of Megilla maculata; results of other observations on this subject.
- 2146. [RILEY, C. V.] Clothes-moths observed in the United States. <Amer. Nat., March [21 February], 1883, v. 17, p. 323.

 Notice of paper by C. H. Fernald.
- 2147. RILEY, C. V. Entomological notes. <Rural New Yorker, 27 February, 1883. S.-b. No. 42, p. 58. See: <Amer. Nat., April [15 March], 1883, v. 17, pp. 419-420.
 - Abstract of the rules of the international convention at Berne for the prevention of *Phylloxera* ravages; their adoption by Belgium.
- 2148. RILEY, U. V. Notice of an illustrated essay on the *Noctuidæ* of North America. <Bull. Brooklyn Ent. Soc., February, 1883, v. 5, pp. 77-79. Separate: <[Brooklyn, N. Y., 1883], 4 pp. Critical review of A. R. Grote's essay.
- 2149. RILEY, C. V. Concerning canker-worms. <Indiana Farmer, 3
 March, 1883. S.-b. No. 56, p. 69. <Prairie Farmer, 3 March,
 1883. S.-b. No. 56, p. 69. <Pacific Rural Press, 10 March,
 1883. S.-b. No. 54, p. 15. <Lancaster Farmer, March, 1883.
 S.-b. No. 56, p. 72. <Gardener's Mo. and Hortic., April, 1883,
 v. 25. S.-b.No. 42a, p. 631.

Description of Paleacrita vernata and Anisopteryx pometaria; ravages of the former; request for information; method of observation.

- 2150. RILEY, C. V. Dipterous enemies of the *Phylloxera vastatrix*. <Ca. Ent., February [9 March], 1883, v. 15, p. 39.
 - Critical review of paper of T. W. Fyles; the characters given of *Diplosis grassator* are insufficient to distinguish the species; galls of *Phylloxera vastatrix* inhabited by *Leucopis phylloxera* Riley MS.; comparison of the early stages of these two Diptera.
- 2151. [RILEY, C. V.] The food relations of the Carabidæ and Coccinellidæ. <Amer. Nat., April [15 March], 1883, v. 17, pp. 417-419.

 Summary of S. A. Forbes's observations on the food relations of the Carabidæ and Coccinellidæ.
- 2152. [RILEY, C. V.] Relations of the *Carabida* and *Coccinellida* to birds. <Amer. Nat., April [15 March], 1883, v. 17, p. 419.

 Reprint of S. A. Forbes's summary, with note; *Coccinellida* not caten by birds; *Carabida* eaten in proportion as they have phytophagous habits.
- 2153. [RILEY, C. V.] Viviparity in a moth. <Amer. Nat., April [15 March], 1883, v. 17, p. 420.

 Notes Fritz Müller's discovery of a moth seen to deposit living larvæ.
- 2154. [RILEY, C. V.] Damage to silver-plate by insects. <Amer. Nat., April [15 March], 1883, v. 17, p. 420.

 Holes in silver-plate said to have been made by Niptus hololeucus.
- 2155. RILEY, C. V. Possible food-plants of the cotton-worm. Amer.
 Nat., April [15 March], 1883, v. 17, pp. 421–422.
 - Notice of paper of J. S. Bailey; occurrence of newly issued images of *Aletia* xylina at Karuer, N. Y., 7 and 8 October, 1882, proving that the larva of this insect feeds upon some genus of plants other than Gossypium.
- 2156. [RILEY, C. V.] Agrotis messoria Harr. vs. Agrotis scandens Riley. <Amer. Nat., April [15 March], 1883, v. 17, p. 422, 2 figs.
 - Critical review of views of A. R. Grote; Agrotis lycarum, A. repentis, and A. cochranii = A. messoria; A. scandens is a distinct species; comparison of the imagos; figures larvæ and imagos of the two species.
- 2157. [RILEY, C. V.] An internal mite in fowls. <Amer. Nat., April [15 March], 1883, v. 17, pp. 422–423.
 - Lungs, bronchia, and linings of thoracic and abdominal cavities of a sick chicken covered with *Cytoleichus sarcoptoides* Mégnin; habitat of the same in fowls in Europe; diseases caused by it.
- 2158. [RILEY, C. V.] Prevalence of the screw-worm in Central America. <Amer. Nat., April [15 March], 18-3, v. 17, p. 423.

 Extract from letter of L. C. Zeledon on the abundance and rayages of Lucilia.
 - Extract from letter of J. C. Zeledon on the abundance and ravages of *Lucilia* macellaria and related flies in Costa Rica.
- 2159. R[ILEY], C. V. Dried leaves as food for lepidopterous larvæ. <Amer. Nat., April [15 March], 1883, v. 17, pp. 423-424.
 - Review of paper of A. H. Mundt; larvæ successfully fed upon fresh leaves transported from a distance under pressure; chopping the leaves would possibly permit of more rapid curing and more convenient packing.
- 2160. [RILEY, C. V.] Lepidopterological notes. <Amer. Nat., April [15 March], 1883, v. 17, p. 424.
 - Duplication of descriptions of the early stages of Lepidoptera to be avoided; notice of Mrs. C. H. Fernald's list of *Noctuide* taken in Orono, Me.

- 2161. [RILEY, C. V.] Obituary. <Amer. Nat., April [15 March], 1883, v. 17, p. 424.
 - Obituary notices of G. W. Belfrage and F. W. Mæklin.
- 2162. R[ILEY], C. V. Mosquitoes vs. malaria. <Sci. Amer., 14 April, 1883, v. 48, pp. 224–225. S.-b. No. 42, pp. 63–64; No. 67, p. 8. Abstract: <Amer. Nat., May [18 April], 1883, v. 17, p. 549.
 - Criticism of the views of Dr. A. F. A. King in support of the thesis that malarial disease is the result of inoculation of the body with malarial poison by the bites of insects; citation of twenty correspondencies in the conditions affecting the prevalence of mosquitoes [Culicidæ] and malarial disease.
- 2163. RILEY, C. V. Jumping seeds and galls. <Sci. Amer., 14 April, 1883, v. 48, p. 228, fig. S.-b. No. 42, pp. 61–63; No. 67, p. 2.
 - Figures of larva, pupa, and imago of Carpocapsa saltitans, with figures and description of seeds inhabited by the larva of this moth, and description of the plant bearing these seeds; vernacular names of plant and insect; movements imparted to the seeds by the insects and by Cynips q.-saltatorius to the galls of the same.
- 2164. RILEY, C. V. Reports of experiments, chiefly with kerosene, upon the insects injuriously affecting the orange-tree and the cotton-plant, made under the direction of the entomologist. Subartaledam. No. 1, Div. Ent. U. S. Dept. Agric., [17 April], 1883, 62. pp.

CONTENTS.

Letter of submittal	3
Introduction	5
Miscellaueous notes on orange insects. By H. G. Hubbard	9
Experiments upon scale insects affecting the orange. By Jos. Voyle	19
Report of observations and experiments. By J. C. Neal	31
Report of observations and experiments on the cotton-worm (Aletia	
xylina). By R. W. Jones	47
Report upon the cotton-worm, boll-worm, and other insects. By Law-	
rence Johnson	53
Index	59

2165. RILEY, C. V. Reports of observations on the Rocky Mountain locust and the chinch-bug, together with extracts from the correspondence of the division on miscellaneous insects. <Bull. No. 2, Div. Ent. U. S. Dept. Agric., [17 April], 1883, 36. pp. Second edition, 16 September, 1883.

'CONTENTS.

Letter of submittal	3
Iutroduction	5
Report of observations in the Northwest on the Rocky Mountain locust,	
by Lawrence Bruner	7
Experiments on chinch-bugs. By S. A. Forbes	23
Extracts from correspondence	27
Iudex	35
	-

2166. [RILEY, C. V.] Insects as food for man. <Amer. Nat., May [18 April], 1883, v. 17, pp. 546-547.

Summary of Max Buchner's observations on the insects used as food by the Bantus tribe of negroes.

- 2167. RILEY, C. V. Number of molts and length of larval life as influenced by food. <Amer. Nat., May [18 April], 1883, v. 17, pp. 547-548.
 - Variability in habits and characters of insects; periods and number of molts observed in larvæ of *Tenebrio molitor*, *T. obscurus* and *Trogoderma tarsale*; conclusion that insufficient food retards development and occasions frequent molting.
- 2168. [RILEY, C. V.] Entomological notes. <Amer. Nat., May [18 April], 1883, v. 17, pp. 549-550.

 Brief notes on recent publications.
- **2169.** RILEY, C. V. Larval stages and habits of the bee-fly *Hirmoneura*. Science, 27 April, 1883, v. 1, pp. 332–334, figs. 1–3. S.-b. No. 42, p. 52.
 - Summary of the life-history of *Hirmoneura obscura* condensed from Handlirsch, and from Brauer; figures the several stages; correspondences of the structure and early history of the larva; predictions in reference to the larvæ of *Bombyliidæ*.
- 2170. RILEY, C. V. The capitalizing of specific names. < Papilio, March [April], 1883, v. 3, p. 62.
- Inquiry as to the use and purpose of capitalizing specific names. See No. 2257. 2171. RILEY, C. V. Observations on the fertilization of Yucca and on
- structural and anatomical peculiarities in Pronuba and Prodoxus. <Gardener's Mo. and Hortic. April, 1883, v. 25, pp. 118–119. S.-b. No. 51, p. 122½. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1882, 1883, v. 31, pp. 467–468. Separate: <Salem. Mass., July, 1883, pp. 467–468. Abstract: <Amer. Nat., February [31 January], 1883, v. 17, p. 197.
 - Description of the manner in which *Pronuba yuccasclla* gathers the pollen in flowers of *Yucca*; the work of this species necessary for the fertilization of the capsular species of *Yucca*, the irregularity of whose fruit is due to its punctures; description of the egg and of the manner of oviposition of this species.
- 2172. RILEY, C. V. The potato-stalk borer. <Rural New Yorker, 12 May 1883. S.-b. No. 42a, p. 213.
 - Answer to inquiry; life-history and means against Baridius [= Trichobaris] trinotata.
- 2173. RILEY, C. V. Jumping seeds and galls. <Proc. U. S. Nat. Mus., 12 May, 1883, v. 5, pp. 632–635, fig. Separate: <[Washington, 1884], pp. 632–635, fig.
 - Description of seeds of "arrow-weed"; their motions caused by larvæ of Carpocapsa saltitans; figures of the seeds and of the larva, pupa, and imago of the Carpocapsa; habits and seasons of the same; description of the plant bearing these seeds; character and motions of the gall of Cynips q.-saltatorius.
- **2174.** RILEY, C. V. Note on cluster-flies. <Proc U. S. Nat. Mus., 12 May, 1883, v. 5, pp. 636–637.
 - Habits of *Pollenia rudis*; notice of recorded cases of swarming among the Diptera. See No. 2128,

- 2175. [RILEY, C. V.] The new classification of the Coleoptera of North America. <Amer. Nat., June [17 May], 1883, v. 17, pp. 660-661.
 - Notice of work of J. L. Lecontc and G. H. Horn.

are most endangered.

- 2176. RILEY, C.V. A pretty and unique gall-making tortricid. < Amer. Nat., June [17 May], 1883, v. 17, p. 661, fig. 1.
 - Description and figure of imago of *Grapholitha ninana* n. sp. reared from galls on stems of *Acacia filicina* in Arizona.
- 2177. [RILEY, C. V.] Simulium feeding on other insects. < Amer. Nat., June [17 May], 1883, v. 17, pp. 661-662. Comment on note of H. A. Hagen.
- 2178. [RILEY, C. V.] Synopsis of the N. A. Heliothinæ. <Amer. Nat., June [17 May], 1883, v. 17, pp. 662-663.

 Review of paper of J. B. Smith; nature of the generic characters of Noctuidæ.
- 2179 [RILEY, C. V.] Death of Professor Zeller. <Amer. Nat., June [17 May], 1883, v. 17, p. 663.
 Obituary notice of P. C. Zeller.
- 2180. [RILEY, C. V.] Protection of insect collections. <Amer. Nat., June [17 May], 1883, v. 17, pp. 663-664.

 Power of the larvæ of Dermestidæ to endure the effects of certain insecticides; prerequisites more important than insecticides; seasons when collections
- 2181. [RILEY, C. V.] Cocoon of Telea polyphemus. <Amer. Nat., June [17 May], 1883, v. 17, p. 664.
 - Answer to inquiry of H. Morris; cocoon of *Telea polyphemus* usually falls to the ground with the leaves, yet quite frequently it is attached to twigs and does not fall to the ground.
- 2182. [RILEY, C. V.] The sucking organs of bees, wasps, and flies. Amer. Nat., June [17 May], 1883, v. 17, pp. 664-665.

 Notice of paper of K. Kraepelin.
- 2183. [RILEY, C. V.] The "pine moth of Nantucket." < Amer. Nat., June [17 May], 1883, v. 17, pp. 665-666.
 Critical review of paper of S. H. Scudder.
- 2184. [RILEY, C. V.] Entomological notes. <Amer. Nat., June [17 May], 1883, v. 17, pp. 666-667.

 Brief notes on recent publications.
- 2185. RILEY, C. V. Elephantiasis or Filaria disease. <Science, 18
 May, 1883, v. 1, pp. 419-421, fig. S.-b. No. 42, p. 51, fig.
 - Criticism of the views of Dr. A. F. A. King; notice of the writings of P. Manson and others on the connection of *Culex musquito* with the life-history of *Filaria sauguinis-hominis* and on the production of elephantiasis and related diseases by *Filaria*.
- 2186. RILEY, C. V. Nemestrinida. Science, 8 June, 1883, v. 1, p. 513.

 Note on papers concerning Nemestrinida; supplementary to No. 2169.
- 2187. RILEY, C. V. The corn-root Diabrotica. <Rural New-Yorker, 9
 June, 1883. S.-b. No. 54, p. 42.
 Increasing distribution of Diabrotica longicornis; means against it.

- 2188. RILEY, C. V. Elm-leaf beetle. <Sci. Amer., 16 June, 1883, v. 48. S.-b. No. 42a, p. 265.
 - Reply to inquiries concerning Galeruca xanthomelana.
- 2189. RILEY, C. V. A unique and beautiful noctuid. <Amer. Nat., July [20 June], 1883, v. 17, pp. 788-790, fig.
 - Figure of imago of Cirrhophanus triangulifer; description of its generic characters; its affinities, synonyms, and probable habits.
- 2190. [RILEY, C. V.] Insects affecting stored rice. < Amer. Nat., July [20 June], 1883, v. 17, p. 790.
 - Lists of insects, chiefly Coleoptera, found in a lot of damaged rice from the Chinese centennial exhibit; two species are carnivorous.
- 2191. RILEY, C. V. Hypermetamorphoses of the Meloidæ. <Amer. Nat., July [20 June], 1883, v. 17, pp. 790–791.
 - Proposal of simpler and more natural terms to designate the stages of development of larvæ of Meloidæ.
- 2192. [RILEY, C. V.] Entomological notes. Amer. Nat., July [20 June], 1883, v. 17, pp. 792–793.

 Comments on recent entomological publications; items of news.
- 2193. RILEY, C. V. The grape Phylloxera in France. <Science, 22 June, 1883, v. 1, pp. 576-578.
- Review and criticism of the report of the Commission supérieure du Phylloxera. 2194. RILEY, C. V. Fig insects. <Science, 29 June, 1883, v. 1, p. 599.
- Review of S. S. Saunders's views on fig insects.

 2195. RILEY, C. V. Egg-punctures on raspberry- and grape-vines, etc.
 - Rural New-Yorker, 30 June, 1883, v. 42, p. —. S.-b. No. 42, p. 56.
 Answer to inquiry of T. H. G.; oviposition of Œcanthus niveus in stems of raspberry- and grape-vines, and of Ceresa bubalus in twigs of apple-trees;
 - raspberry- and grape-vines, and of *Ceresa bubalus* in twigs of apple-trees; habits and ravages of and means against the former; the latter seldom very injurious.
- **2196.** RILEY, C. V. Silk culture in the United States. < Rural New-Yorker, 14 July, 1883, v. 42, p. —. S.-b. No. 42a, p. 531.
 - Revival of interest in silk culture in the United States; experience of the year; relation of silk culture to import duties; warning against too high expectations as to profits; notes on manuals of silk culture.
- 2197. RILEY, C. V. Economic entomology of Iowa. Sci. Amer., 14
 July, 1883, v. 49, p. —. S.-b. No. 53, p. 147.
 - Review of work done in Iowa, especially of papers of J. N. Dixon, H. Osborn, and A. B. Walton.
- 2198. [RILEY, C. V.] Insect plagues. <Boston Herald, 22 July, 1883, p. —. S.-b. No. 42, pp. 53-54; No. 67, p. 11. Extract: <Mirror and Farmer, 26 July, 1883, v. 35. S.-b. No. 42, pp. 126-127.
 - Newspaper interview; seasons, habits, ravages of and means against Caloptenus atlanis, Nematus erichsonii, and Orgyia leucostigma.
- 2199. RILEY, C. V. Report by C. V. Riley. <Proc. U. S. Nat. Mus., 27 July, 1883, v. 6, pp. 104–105. Reprint: <Sci. Amer. Suppl., 13 October, 1883, v. 16, p. 6486.
 - Note to F. Humbert's Lucilia macellaria infesting man; references to other mentions of it; its distribution and means against it.
 - 20 ENT

- 2200. RILEY, C. V. Emulsions of petroleum and their value as insecticides. <Proc. Amer. Assoc. Adv. Sci. for 1882, 1883, v. 31, pp. 469-470. Separate: <Salem, Mass., July, 1883, pp. 469-470. Reprint: <Kansas City Rev. of Science and Industry, November, 1883, v. 7, pp. 447-448. S.-b. No. 42, p. 124. Extract: <Sci. Amer., 19 November, 1883, v. 49, p. 294. S.-b. No. 51, p. 153.
 - Description of modes of making emulsions of petroleum for use against insects.
- 2201. [RILEY, C. V.] The old, old question of species. <Amer. Nat., September [15 August], 1883, v. 17, p. 975.
 - Comments upon the discussion between H. A. Hagen and W. H. Edwards; views of both parties extreme; views as to the true nature of species.
- 2202. [RILEY, C. V.] Myrmecophila. Amer. Nat., September [15 August], 1883, v. 17, pp. 975-976.

 Record of recent captures of Myrmecophila in Oregon and District of Colum-

bia; record of former captures in the United States; habits of the genus.

- 2203. [RILEY, C. V.] Salt-water insects used as food. <Amer. Nat., September [15 August], 1883, v. 17, pp. 976-977.
 - Occurrence of Ephydra (hians?) in Lake Tetscoco, Mexico; E. gracilis found in Great Salt Lake, Utah, and E. californica in lakes in California; account by W. H. Brewer of the manner in which the last-named species is collected and used for food by the Indians living near Mono Lake.
- 2204. [RILEY, C. V.] Food-plants of Samia cynthia. <Amer. Nat., September [15 August], 1883, v. 17, p. 977.
 - Review of paper of H. H. Birney; list of plants on which Samia [== Attacus] cynthia has hitherto been found feeding; some of these are the favorite food-plants of Callosamia [= Attacus] promethea.
- 2205. [RILEY, C. V.] Bitten by an aphid? <Amer. Nat., September [15 August], 1883, v. 17, p. 977.
 - Letter of S. Swan, with answer; Siphonophora [= Nectarophora] rudbeckiæ common on Solidago and Rudbeckia; the biting was probably caused by antsor some other insect that escaped notice at the time.
- 2206. RILEY, C. V. Steganoptycha claypoleana. <Amer. Nat., September [15 August], 1883, v. 17, p. 978. Reprint: <Papilio, September-December, 1883, v. 3, p. 191.
 - Comparative description of Steganoptycha claypoleana with Proteoteras asculanum; habits of the former.
- 2207. R[ILEY], C. V. Extermination and restriction of Phylloxera in Switzerland. <Rural New-Yorker, 25 August, 1883, v. 42. S.-b. No. 49, p. 74.

Notice of report of Valery-Mayet.

- 2208. RILEY, C. V. Hackberry psyllid galls. <Ca. Ent., August [5 September], 1883, v. 15, pp. 157–159, figs. 6, 7.
 - Critical review of paper of T. W. Fyles; Phylloxera vastatrix has many parasites and Celtis is attacked by many species of gall-insects; characters of Pachypsylla n. g. [p. 157]; figures of galls of P. celtidis-venusta and P. c.-mamma; derivation and orthography of the generic term Celtis.

- 2209. RILEY, C.V. Some recent discoveries in reference to Phylloxera. <Science, 7 September, 1883, v. 2, p. 336. S.-b. No. 42, pp. 68, 69. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1883, 1884, v. 32, p. 320. Separate: <Salem, Mass., July, 1884, p. 320. <Amer. Nat., December [28 November], 1883, v. 17, p. 1288.
 - Summary of the cycle of development of the genus *Phylloxera*; character of the gall of *P. spinosa* and location of the impregnated egg of the species.
- 2210. RILEY, C. V. The *Psyllida* of the United States. <Science, 7 September, 1883, v. 2, p. 337. S.-b. No. 42, pp. 67-68. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1883, 1884, v. 32, p. 319. Separate: <Salem, Mass., July, 1884, p. 319.
 - Characteristics and economic importance of *Psyllida*; status of the present knowledge of this family in the United States; list of new genera and species; food-plants of the same; characteristics of the eggs and young of *Psyllida*.
- 2211. RILEY, C. V. Improved method of spraying trees for protection against insects. <Science, 14 September, 1883, v. 2, p. 378. S.-b. No. 42, p. 68. Reprint: <Proc. Amer. Asoc. Adv. Sci. for 1883, 1884, v. 32, pp. 466–467. Separate: <Salem, Mass., July, 1884, pp. 466, 467.

Description of apparatus, especially of "cyclone nozzle" and adjustable hose; mention of insecticide substances.

Note.—Nos. 2209-2211 were issued as a separate, Salem, 1884.

- 2212. [RILEY, C. V.] Entomology at Minneapolis. <Amer. Nat., 1883, v. 17: October [17 September], pp. 1068–1070; November [19 October], pp. 1169–1174.
 - Minutes of the meetings of the entomologists at the session of the American Association for the Advancement of Science; reorganization of the entomological club; election of officers and change in the constitution; abstracts of papers read.
- 2213. RILEY, C. V. Notes on Pædisca scudderiana. <Amer. Nat., October [17 September], 1883, v. 17, pp. 1069–1070.

 Habits and variation of Pædisca scudderiana; difference between the gall of

this insect and that of Gelechia gallæsolidaginis.

2214. RILEY, C. V. A myrmecophilous lepidopteron. <Amer. Nat.,
October [17 September], 1883, v. 17, p. 1070.
Larva of Helia americalis found in nests of Formica rufa; this species the only

one of the Lepidoptera known to develop in ants' nests.

- 2215. [RILEY, C. V.] Enemies of the egg-plant. <Amer. Nat., October [17 September], 1883, v. 17, p. 1070.
 - Extract from a letter of A. Œmler, proving that the occurrence of Cassida texana and Doryphora juncta on Solanum melongena is not accidental or temporary.
- 2216. [RILEY, C. V.] The periodical Cicada in southeastern Massachusetts. <Amer. Nat., October [17 September], 1883, v. 17, p. 1071.
 - Note to paper of C. E. Bessey; Cicada [= Tibicen] septendecim at Martha's Vineyard, Mass., in June, 1883, the precursor to septendecim Brood XXI which will appear in North Carolina and Virginia as well as Martha's Vineyard in 1884; accelerated specimens of the same brood received from Loudoun County, Va., recently.

- 2217. RILEY, C. V. Habits of Murmidius. <Amer. Nat., October [17 September], 1883, v. 17, p. 1071.
 - List of insects found in a lot of damaged rice from South America; occurrence of Murmidius ovalis in vast numbers in this rice; its probable food-habits; description of its cocoon; list of families of beetles some of whose larvæ spin cocoons; habitat of Mychocerus.
- 2218. [RILEY, C. V.] Obituary. <Amer. Nat., October [17 September], 1883, v. 17, p. 1072.
 Notices of V. T. Chambers and Townend Glover.
- 2219. [RILEY, C. V.] Entomological notes. <Amer. Nat., October [17 September], 1883, v. 17, pp. 1072–1073.

 Notes on recent entomological publications; items of news.
- 2220. [RILEY, C. V.] Economic notes. < Amer. Nat., October [17 September], 1883, v. 17, pp. 1073-1074.

Comments on the ravages of several injurious insects during the past season.

- 2221. RILEY, C. V. A parasite of the cabbage-worm. < Rural New-Yorker, 6 October, 1883, v. 42. S.-b. No. 42, p. 58.

 Answer to letter of J. H. B.; parasitism of Pteromalus puparum in larvæ and pupæ of Pieris rapæ.
- 2222. RILEY, C. V. The handmaid moth. <Rural New Yorker, 13 October, 1883, v. 42. S.-b. No. 42, p. 77.
 - Answer to inquiry of H. B. S.; description of larva, pupa, and image of *Datana ministra* from hickory- and walnut-trees, and of a phytophagic variety of the larva from apple and other trees; habits of the larvæ; the larvæ unusually abundant in 1883.
- 2223. RILEY, C. V. Remarks on Arzama obliquata. <Amer. Nat., November [19 October], 1883, v. 17, p. 1169.

Description of the egg-mass of Arzama obliquata; colors and habitat of the larva; variations of the imago; number of annual broods.

- 2224. [RILEY, C. V.] Rare monstrosities. <Amer. Nat., November [19 October], 1883, v. 17, p. 1175.
 - Notice of monstrosities recorded in Melanippe montanata and Zygana minos.
- 2225. [RILEY, C. V.] The nervous system of insects. <Amer. Nat., November [19 October], 1883, v. 17, pp. 1175-1176. Summary of the observations of Ed. Brandt.
- 2226. [RILEY, C. V.] Hymenorus rufipes as a myrmecophilous species. <Amer. Nat., November [19 October], 1883, v. 17, p. 1176.

 Imagos of Hymenorus rufipes raised from larvæ found in nests of Formica fusca;

character of the nests of the Formica; food-habits of the Hymenorus unknown.

- 2227. [RILEY, C. V.] Recent publications. <Amer. Nat., November [19 October], 1883, v. 17, p. 1177.

 Notice of J. H. Comstock's work on Diaspina and of other recent publications.
- 2228. [RILEY, C. V.] Entomological notes. <Amer. Nat., November [19 October], 1883, v. 17, pp. 1177-1179.

Review of "General Index of the Entomological Reports of the Province of Ontario;" recent publications and items of news.

99

101

2229. RILEY, C. V. The potato-stalk borer. <Rural New-Yorker, 20 October, 1883. S.-b. No. 42, pp. 78-79.

Answer to letter of S. C. R.; description, habits of, and means against Gortyna nitela; means against Paria aterrima, Graphops pubescens, and other larvæ of Chrysomelidæ injurious to the roots of strawberry-plants.

- 2230. [RILEY, C. V.] Recent advances in horticultural entomology. <Rural New-Yorker, 20 October, 1883, v. 42. S.-b. No. 42, pp. 79–81. Reprint: <Proc. 19th Sess. Amer. Pomol. Soc., 1884, p. 45. <Trans. Wise. State Hortic. Soc., 1886, v. 17, p. —.
 - Report of address delivered; discussion of measures recommended for adoption to prevent the ravages of insects injurious to horticulture, especially of Carpocapsa pomonella and Conotrachelus nenuphar; correction of statements in regard to the oviposition of Saperda bivittata [= candida] and? Bembex marginata; advance in knowledge of the life-history of Aphidida and in the development of machinery for the application of poison sprays to plants; relative value of the principal insecticides.
- 2231. RILEY, C. V. On a gall-making genus of *Apioninæ*. <Bull. Brooklyn Ent. Soc., October, 1883, v. 6, pp. 61–62. Separate: <Brooklyn, N. Y., 1883, 2 pp.

List of North American gall-making Coleoptera; description of *Podapion* n. g. [p. 62] and of the gall and imago of *P. gallicola* n. sp., found on twigs of *Pinus inops*; probable life-habits, inquilines, and parasite of this species.

2232. RILEY, C. V. Report of the Entomologist. <Ann. Rept. [U. S.] Commissioner Agric. for 1883, pp. 99–180, 13 pl. Separate: <Washington, 31 October, 1883, pp. 5+pp. 99–180+pp. 7, 13 pl.

TABLE OF CONTENTS.

Scope and limitations, 99—Cabbage insects, 99—Lesser locust, 99—Protection of frnit and shade trees, 99—Office force, assistants and agents, 99—Observations and report on insects injurions to forest-trees, by Dr. A. S. Packard, jr., 99—Special report on insects affecting the orange, by H. G. Hnbbard, 99—Work of agents, 100—Monographs in preparation, 100—Commission sent to Brazil to study certain insects, 100—Publication of third report of the United States Entomological Commission, 100—Publication of special bulletins, 100—Correspondence and work of the division, 101—The illustrations to the report, 101.

Legislation, 101—California State Board of Silk Culture, 101—Preminms offered in California, 101—Home culture, 102—Guaranty of egg supply, 102—Woman's Silk Culture Association of California, 102—California Silk Culture Association, 102—Reports from correspondents of the division, 102–104—Issne of "Silk Culture Directory," 104—Establishment of school in New Jersey, 105—Organization of Sonthern Silk Industrial Association, 105—A new machine patented, 105—Report on the future of silk culture in the United States, by United States Consul Peixotto, of Lyons, France, 105—Silk culture by the Mennonites, 106—Work at the Department, 106.

0000	Description of the state of the	
2232.	RILEY, C. V.—Continued.	
	Cabbage worms	107
	Interest in and importance of the subject, 107—Accurate estimate	
	of loss impossible, 107.	400
	Imported cabbage-worm, Pieris rapæ.	108
	History of its spread, 108—Ravages, 109—Characters, 110—Habits,	
	110—Food-plants, 111—Seasons of appearance, 111—Natural ene-	
	mies, 111.	
	Southern cabbage-butterfly, Pieris protodice.	114
	Range, 114—Ravages, 114—Characters, 114—Food-plants, 115—	
	Parasites, 115.	
	The potherb-butterfly, Pieris oleracea	115
	Range, 115—Characters, 116—Habits, 116—Number of broods, 116—	
	Food-plants, 117—Enemies, 117.	
•	The larger cabbage-butterfly, Pieris monuste	117
	Range, 117—Characters, 117—Food-plants, 118—Parasites, 118—	
	Description of early states, 118.	110
	The cabbage plusia, Plusia brassice.	119
	Its range, 119—Food-plants, 119—Habits and natural history, 119—	
	Close relation to an European species, 120—Natural euemies,	
	120—Botrytis rileyi, a new species of fungus, 121—Remedies, 121.	100
	The cabbage mamestra, Mamestra trifolii	123
	Its habits and natural history, 123—Remedies, 124—Description,	
	124.	104
	The zebra cabbage-worm Mamestra picta.	124
	History, 124—Description, 125—Habits, 125—Broods, 125—Reme-	
	dies, 125.	100
	The cabbage pionea Pionea rimosalis	126
	Its past history, 126—Notes by Prof. Cyrus Thomas, 126—Habits	
	and natural history, 127—Parasites, 127—Remedies, 127—De-	
	scriptive, 128. The cauliflower botis, Botis repetitalis	128
		120
•	Past history, 128—Rauge, 128—Descriptive, 129.	129
	The cabbage plutella, Plutella cruciferarum.	129
	Its past history, 129—Habits and natural history, 130.	101
	REMEDIES FOR CABBAGE-WORMS.	131
1	Hot water, 131—Pyrethrum, 131—Kerosene emulsions, 131—Other	
	substances, 131—Paris green, London purple, and white helle- bore, 132—Preventive measures, 132—Report of Capt. R. S.	
	Lacey, 133—Report of Col. Wright Rives, 134—Poisoning de-	
	vices, 136. Causes of destruction of evergreen forests in New England	
	AND NEW YORK, BY A. S. PACKARD, JR	138
	The larch saw-fly, Nematus ericksonii	138
	History of its ravages, 138—Its devastations in Maine, 139—Its	100
	ravages in New Hampshire, 141—Its appearance in Massachu-	
	setts, 141—In northern New York, 142—History of the species	
	and its habits, 142—Description of egg, larva, cocoon, and imago,	
	145—Remedies, 146—Parasites, 146.	
	The spruce-bud tortrix, Tortrix fumiferana	146
	Its ravages in Maine, 146—Habits and transformations, 147—De-	140
	scription of different stages, 148.	
	The spruce nematus, Nematus integer	149
	Range, 149—Description of different stages, 150.	
	Trained, 110 Donott Prior of alterior of the good 100.	

2232.	RILEY, C. V.—Continued.	
	Causes of destruction of evergreen forests, etc.—Continued.	
	The hemlock gelechia, Gelechia abietisella	150
	Habits, 151—Description of larva, pupa, and moth, 151.	
	EXPERIMENTS ON SCALE INSECTS, WITH PRACTICAL SUGGESTIONS, BY	
	H. G. HUBBARD	152
,	Report of progress in experiments, 152—Soap emulsions, 152—Kerosene and soap emulsions, formula, 152—Receipt for making emulsions of kerosene and soap, 152—Introduction and use of the cyclone nozzle, 152—Effects of kerosene emulsions upon plants, 153—Effects of kerosene emulsions varied by change of weather, 153—Most favorable season for applying kerosene emulsions, 153—Application of liquid insecticides, 153—The cyclone nozzle, 154—Cost of kerosene wash, 154.	
	Recent experiments	155
	Potash compared with soda as an insecticide, 155—Details of ex-	100
	periments with potash, 155-156—Experiments with soda, 156.	150
	Introduction and spread of scale insects Great witality of healt lies 156. Importation on live trees 157	156
	Great vitality of bark-lice, 156—Importation on live trees, 157— Precautionary measures; infection from nursery stock, 157—Pro-	
	tection afforded by hedges and forest trees, 158.	
	MISCELLANEOUS INSECTS	159
	The imported elm-leaf beetle, Galeruca xanthomelana	159
	Range, 159—An importation from Europe, 160—Habits and natu-	
•	ral history, 160—Remedies, 161—Natural enemies, 163—More re-	
	cent experiments at the Department, 164-Past history of the	
	elms on Department grounds, 164—Condition and characteristics of the grove in 1882 and 1883, 164—Extent of injury in 1882	
	and 1883, 164—Preferences of the elm-beetle for certain varieties	
	and species of clms, 165—Effects of arsenical poisons on insect	
	and plant, 165—Preventive effects of the poison, when best applied, 166—Treatment with London purple, 166—Preparation of the poison, 166—Effects of the mixture, 167—Treatment with Paris green, 167—Mechanical means of applying the poison, 168—The eddy-chamber nozzle, 168—Hose and bamboo combination.	
	tion, 169. The legger migratory legget Calentonic atlanic	170
	The lesser migratory locust, Caloptenus atlanis. Its ravages in the Merrimac Valley, 170—Historical, 170—Characters, 172-Range and life-history, 173—Natural enemies, 174—Remedies, 175—Destruction of eggs, 175—Destruction of young, 175—Coal oil, 175—Machines, 175–177—Coal tar, 177—Catching or bagging, 178—Protection of vegetable gardens, 179—Necessity for co-operation, 180.	170
2233.	RILEY, C. V. The imported orchard Scolytus (Scolytus rugula	osus
		8b.
	' -	+
0004	Answer to inquiry; S. rugulosus bores in twigs of pear; means against i	
2234.	RILEY, C. V. Osage orange vs. mulberry for the silk-wo <sci. 17="" 1883,="" 305.="" 49,="" amer.,="" b.="" no.<="" november,="" p.="" s.="" td="" v.=""><td></td></sci.>	
	p. 91. Reprint: < Amer. Nat., 5 January, 1884, v. 18, pp.	

42, p. 118.

Comparative value of leaves of Maclura aurantiaca and of Morus as food for *Sericaria mori; critical review of V. dcs Lauriers's conclusions.

79. < Prairie Farmer, 22 March, 1884, v. 56, p. 187. S.-b. No.

2235. RILEY, C. V. Entomological notes of the year. < Prairie Farmer, 24 November, 1883, v. 55. S.-b. No. 42, pp. 86-87.

Progress of experiments for the destruction of scale insects, Coccidæ; introduction of Aspidiotus rapax into California on apples from New Zealand; strawberries injured by Capsus oblineatus [= Lygus pratensis] and a myriapod in Illinois; occurrence of Leucania unipuncta in several places, and of Cecidomyia destructor in Illinois; extension of culture of Pyrethrum cinerariæfolium in California; occurrence of Anarsia lineatella on strawberry-plants in Illinois; of Doryphora juncta and Cassida texana on Solanum melongena in Georgia; of Ceresa bubalus on potato-plants in Pennsylvania; of Epilachna corrupta on wax-beans, etc.

2236. RILEY, C. V. A satisfactory remedy for melon bugs, flea-beetles, etc. <Rural New-Yorker, 3 November, 1883, v. 42. S.-b. No. 42, pp. 77–78.

Quinn's method of sprinkling the vines with a mixture of tobacco water and soft soap and then powdering with lime is probably the best general preventive against Diabrotica vittata and Halticida; description of J. M. Nicholson's siphon arrangement by which to keep the vines constantly moist with the liquid.

2237. RILEY, C. V. The chinch-bug in New York. <Science, 9 November, 1883, v. 2, p. 621. Extract: <Sci. Amer., 22 December, 1883, v. 49, p. 384. S.-b. No. 42, pp. 122–123.

Present outbreak of Blissus leucopterus in New York the result of an increase due to favorable conditions rather than an invasion.

2238. R[ILEY], C. V. Insects in relation to agriculture. <Stoddart's Encylopædia Americana, 1883, v. 1, pp. 135–142, figs. 1–29.

Chapter 9 of article "Agriculture." Brief accounts with numerous illustrations of the insects named below, and of means against them, with cross-references to accounts of other insects in other portions of the work. The headings and subjects of the sub-chapters are as follows:

INSECTS INJURIOUS TO FRUIT AND FRUIT TREES:

Apple curculio (The), Anthonomus quadrigibbus Say, p. 135, fig. 1-Apple-magget (The), or "railroad-worm," Trypeta pomonella Walsh, p. 135—White-marked tussock-moth (The), Orgyia leucostigma Sm. and Abb., pp. 135-136, fig. 2—Apple-tree tent-caterpillar (The), Clisiocampa americana Harr., p. 136, figs. 3-4-Fall webb-worm (The), Hyphantria textor Harr. [= cunea], p. 136, fig. 5—Oyster-shell bark-louse of the apple (The), Mytilaspis pomicorticis Riley [= pomorum], pp. 136-137, fig. 6-Round-headed appletree borer (The), Saperda bivittata Say [= candida], p. 137, fig. 7— Flat-headed apple-tree borer (The), Chrysobothris femorata Fabr., p. 137—Spring canker-worm (The), Paleacrita vernata Peck, p. 137, figs. 8-9—Fall canker-worm (The), Anisopteryx pometaria Harris, p. 137, figs. 10-11—Peach-tree borer (The), Ægeria [= Sannina] exitiosa Say, pp. 137-138, fig. 12 — Currant-stalk borer (The), Ægeria tipuliformis L., p. 138—Imported currant-worm (The), Nematus ventricosus Klug. [= ribesii], p. 138, figs. 13-14—Native currant-worm (The), Pristiphora grossularia Walsh, p. 138-Snowy tree-cricket (The), Ecanthus niveus Harr., p, 138, figs. **1**5-16.

2238. R[ILEY], C. V.—Continued.

INSECTS INJURIOUS TO CEREALS AND FORAGE-CROPS:

White-grub (The), Lachnosterna fusca Froh., pp. 138-139, fig. 17—Clover-seed midge (The), Cecidomyia leguminicola Lintner, p. 139—Joint-worm (The), Isosoma hordei Harr., p. 139, fig. 18—Wheat midge (The), Diplosis tritici Kirby, p. 139—Cut-worms, family Noctuidae, genera Agrotis, Mamestra, Hadena, and Prodenia, p. 139—Wire-worms, family Elateridae, p. 139.

INSECTS INJURIOUS TO GARDEN VEGETABLES:

Imported cabbage-worm (Thc), Pieris rapæ Schrank, p. 139, figs. 19-20—Southern cabbage butterfly (The), Pieris protodice Boisd., p. 140—Potherb butterfly (The), Pieris oleracea Boisd., p. 140—Cabbage plusia (The), Plusia brassicæ Riley, p. 140, fig. 21—Harlequin cabbage-bug (The), Murgantia histrionica Hahn, p. 140, fig. 22—Pea-weevil (The), Bruchus pisi Linn., p. 140, fig. 23—Bean-weevil (The), Bruchus fabæ Riley, p. 140, fig. 24—Blister-beetles, meloid genera Macrobasis, Epicauta, etc., p. 140, fig. 25—Striped cucumber-beetle (The), Diabrotica vittata Fabr., pp. 140-141, figs. 26-27.

INSECTS INJURING MISCELLANEOUS FIELD-CROPS:

Cotton-boll worm (The) or corn-ear worm, Heliothis armigera, Hübn., p. 141, fig. 28—Tobacco-worm (The), Sphinx [= Protoparce] carolina L., p. 141.

INSECTS INJURIOUS TO THE VINE.

INSECTS INJURIOUS TO LIVE-STOCK:

Bot-fly of cattle (The), Hypoderma bovis Latr., p. 141—Sheep bot-fly (The), Estrus ovis Linn., p. 141, fig. 29—Horse bot-fly (The), Gastrophilus equi Fabr., pp. 141–142.

LIST OF PRINCIPAL AMERICAN WRITERS AND WRITINGS UPON INSECTS
IN RELATION TO AGRICULTURE

142

- 2239. R[ILEY], C. V. Army-worm, Leucania unipuncta, Haw. <Stoddart's Encylopædia Americana, 1883, v. 1, pp. 317-318, 2 figs. Distribution, scasons, habits, and food-plants of and means against Leucania unipuncta; references to the more important articles on this insect; figures of larva and imago.
- 2240. RILEY, C. V. An epidemic disease of *Caloptenus differentialis*. Amer. Nat., December [28 November], 1883, v. 17, p. 1287.

 Note to communication of H. Osborn; the *Entomoghthora calopteni* a result rather than the cause of disease.
- 2241. RILEY, C. V. The growth of insect eggs. < Amer. Nat., December [28 November], 1883, v. 17, p. 1289.

Notice of paper of J. A. Osborne; swelling of the eggs of Phaneroptera [= Scudderia] curvicauda probably connected with embryological development.

- 2242. [RILEY, C. V.] Protective device employed by a glaucopid caterpillar. <Amer. Nat., December [28 November], 1883, v. 17, p. 1289.
 - Notice of paper of F. Müller; general use of shed hairs by larvæ of Arctiidæ in the construction of their cocoons; description of method in which the larva of Eunomia eagrus arranges its shed hairs to form a protection for the pupa.
- 2243. [RILEY, C. V.] Saw-fly larvæ on the quince. <Amer. Nat., December [28 November], 1883, v. 17, p. 1289.

Notice of paper of J. A. Lintner; food-plants of Selandria [=Eriocampa] cerasi.

- 2244. [RILEY, C. V.] Entomology in New York. < Amer. Nat., December [28 November], 1883, v. 17, pp. 1289–1291.
 - Review of J. A. Lintner's first annual report as State entomologist; list of subjects treated in the report.
- 2245. [RILEY, C. V.] Fruit insects in California. < Amer. Nat., December [28 November], 1883, v. 17, p. 1291.
 Review of M. Cooke's "Injurious Insects of the Orchard."
- 2246. [RILEY, C.V.] Death of Dr. J. L. Leconte. < Amer. Nat., December [28 November], 1883, v. 17, p. 1291.

 Notes the loss to entomologists occasioned by the death of J. L. Leconte.
- 2247. [RILEY, C. V.] Entomological notes. < Amer. Nat., December [28 November,] 1883, v. 17, pp. 1291–1292.

 Brief notices of published articles with items of news.
- 2248. [RILEY, C. V.] Economic notes. <Amer. Nat., December [28 November], 1883, v. 17, p. 1292.

 Effect of Phylloxera laws in Europe; use of lime as a means against Macro-
 - Effect of Phylloxera laws in Europe; use of lime as a means against Macrodactylus; award of the Le Brun prize for the most valuable improvement relating to the silk industry.
- 2249. RILEY, C. V. [Raspberry canes punctured by *Orchelimum glaberrimum*.] <Fruit Recorder, Purdy, 1 December, 1883, v. 15. S.-b. No. 42, p. 92.
 - Description of and means against Orchelimum glaberrimum.
- 2250. [RILEY, C. V.] The Phylloxera in sandy soil. <Rural New-Yorker, 1 December, 1883, v. 42. S.-b. No. 42, p. 97. Reprint: <Amer. Nat., January, 1884 [29 December, 1883], v. 18, p. 78.
 - Condition of the grape crop in France; favorable account of the use of American grape-stocks; *Phylloxera vastatrix* harmless in sandy soils.
- 2251. RILEY, C. V. Bacterial disease of the imported cabbage-worm. <Sci. Amer., 1 December, 1883, v. 49, p. 337. S.·b. No. 42, pp. 91–92. Reprint: <Amer. Nat., January, 1884 [29 December, 1883], v. 18, p. 80.
 - Notice of observations of S. A. Forbes on the death of larvæ of *Pieris rapæ* from infection by *Bacterium*; previous mention of this disease.
- 2252. RILEY, C. V. The chinch-bug in New York State. <Sci. Amer., 8 December, 1883, v. 49. p. 359. Reprint: <Amer. Nat., 5 January, 1884, v. 18, pp. 79–80.
 - Critical review of papers of J. A. Lintner; the occurrence of *Blissus leucopterus* in New York in unusual abundance in 1882 and 1883 is not warrant for great alarm. See No. 2271.
- 2253. RILEY, C. V. Reports of observations and experiments in the practical work of the division, made under the direction of the entomologist. <Bull. No. 3, Div. Ent. U. S. Dept. Agric., [8 December], 1883, 75 pp., 3 pl.

CONTENTS.

Preface	7
Further notes on the army worm (Leucania unipuncta)	9
Experiments with pyrethrum	16

CONTENTS—Continued.

Notes on forest-tree insects. By A. S. Packard, jr	24
The cotton-worm in South Texas in 1883. By E. H. Anderson	31
Test of machinery for destroying the cotton-worm. By W. S. Barnard.	39
The tree-borers of the family Cossidæ. By J. S. Bailey	49
Tests of silk fiber from cocoons raised at the Department. By W. Mc-	
Murtrie-	56

- 2254. RILEY, C. V. Entomology in New York. <Rural New-Yorker, 15 December, 1883. S.-b. No. 42, pp. 85-86.
 - Review of J. A. Lintner's first report (for the year 1881) as State entomologist of New York.
- 2255. RILEY, C. V. Lucilia macellaria. <Sci. Amer., 15 December, 1883, v. 49, p. 373. S.-b. No. 42, p. 93.

 Critical review of paper of F. Humbert.
- 2256. RILEY, C. V. Dipterous larvæ in the human body. <Sci. Amer., 22 December, 1883, v. 49, p. 385. S.-b. No. 42, p. 90.
 - References to and notices of several articles on the occurrence of larvæ of Diptera in the human body; synonymy and distribution of Compsomyia [=Lucilia] macellaria; myasis caused solely by larvæ of Sarcophagidæ and Muscidæ; Æstridæ occur rarely and exceptionally in the human body.
- 2257. RILEY, C. V. Capitalizing specific names. < Papilio, September-December, 1883, v. 3, pp. 164-166. Separate: <[N. Y.], 3 pp. Comments on reasons given by W. H. Edwards and others for the uniform capitalizing of specific names. See No. 2172.
- 2258. R[ILEY], C. V. Townend Glover. <Papilio, September-December, 1883, v. 3, pp. 167-168.

 Obitnary notice.
- 2259. [RILEY, C. V.] The genus *Colias.* < Amer. Nat., [5] January, 1884, v. 18, pp. 74-76.
 - Review of paper of H. A. Hagen; discussion of the species of Colias in North America; plastic nature and classificatory characters of the genus.
- 2260. [RILEY, C. V.] Larval habits of the dipterous family *Dexidæ*. <Amer. Nat., [5] January, 1884, v. 18, pp. 76-77.
 - Notice of paper of F. Brauer; parasitism of Dexia rustica and of Phorostoma latum on the larva of Rhizotrogus and of Melanophora? diabrotica on Diabrotica vittata.
- 2261. [RILEY, C. V.] Entomological notes. < Amer. Nat., [5] January, 1884, v. 18, pp. 80-81.
 - Carpocapsa pomonella has been introduced into Australia, New Zealand, and Tasmania; occurrence of Myrmicoccla ochraceella in ants' nests.
- 2262. RILEY, C. V. Recent outbreaks of the army-worm. < Rural New-Yorker, 12 January, 1884, v. 43, p. 19. S.-b. No. 56, p. 145.

 Rare occurrence of *Leucania unipuncta* in 1882 and 1883, following its last abundance in 1881; occurrence of its larvæ at East Windsor, Vt., in June, 1883.
- 2263. [RILEY, C. V.] The harlequin cabbage-bug, etc. <Rural New-Yorker, 2 February, 1884, v. 43, p. 70. S.-b., No. 63, p. 41.

Habits of and means against Murgantia histrionica; effect of chemicals applied to the soil npon the colors of flowers.

- 2264. RILEY, C. V. Tribute to the memory of John Lawrence Leconte. <Psyche, November-December, 1883 [11 February, 1884], v. 4, pp. 107-110. Separate: <[Cambridge, Mass., 11 February, 1884], pp. 107-110. Notice: <Psyche, loc. cit., p. 119.
 - Biographical notice of J. L. Leconte; his work and writings; his personal character; disposition of his entomological collection.
- 2265. [RILEY, C. V.] Fruit culture in the South. < Washington Post, 26 February, 1884. S.-b. No. 42, pp. 119-120.

Interview with a reporter; means against insects infecting the orange.

- 2266. RILEY, C. V. Oviposition of the round-headed apple-tree borer. <Rural New-Yorker, 1 March, 1884, v. 43, p. 132, fig. 73. S.-b. No. 42, p. 85; No. 67, p. 3.
 - Notice of paper of C. G. Atkins; confirmation of the account of the method of oviposition of Saperda candida; description of this method; correction of several statements by W. Saunders in regard to the oviposition of different insects; description and figure of the egg of S. candida; figures of the burrows and hole of exit of this beetle; figures of the pupa and of the position of the egg when deposited.
- 2267. [RILEY, C. V., et al.] Third report of the United States Entomological Commission, relating to the Rocky Mountain locust, the Western cricket, the army-worm, canker-worms, and the Hessian fly; together with descriptions of larvæ of injurious forest insects, studies on the embryological development of the locust and of other insects, and on the systematic position of the Orthoptera in relation to other orders of insects; with maps and illustrations. <Washington: 1883 [6 March, 1884], pp. 14+347+12+92, 18 figs., 64 pls., 4 maps.

LETTER OF SUBMITTAL X
PREFACE. XII

PART I.—IN REFERENCE TO THE ROCKY MOUNTAIN LOCUST (Caloptenus spretus).

CHAPTER I:

Additions to the chronology of locust ravages in 1880 and 1881.....
The locust in 1880 in Texas, 3; in Colorado, 4; in Utah. 4; in 1881, in Texas, 6; in Utah, 7.

3

21

CHAPTER II:

The Rocky Mountain locust in Montana in 1880. By Lawrence Bruner

The country from St. Paul to Montana, 8—Between Bismarck and Fort Keogh, 9—Burning often impracticable, 9—Destroying locusts by ditches and kerosene eddies, 13—Ropes dragged to drive them, 15—"Drifts of hoppers," 16—Shooting against locust swarms, 16—Destroying by ditches and burning straw, 17—No more damage for three or four years, 17—Enemies of the locust in the Northwest, 17—Topography of western Dakota and Montana, 19.

CHAPTER III:

The Rocky Mountain locust in Wyoming, Montana, etc., in 1881.

By Lawrence Bruner.

Letter of submittal of report, by Lawrence Bruner, 21 — Gen-

Letter of submittal of report by Lawrence Bruner, 21 — General report, 22—Brief history of depredations, 22—The earlier

CHAPTER III—Continued.

ones, 23—Characteristics of the permanent region, 24—Settlement and other checks against locusts, 25—Distribution of the permanent regions, 25—Their physical peculiarities, 28—The arid region, 29—Permanent breeding grounds, 29—Their locations and relations, 30—The sources of locusts, 30—Interchange of swarms, 31—Regular migratory routes, 31—Agencies controlling migration, 32—Disadvantages of the temporary regions, 33—Period of hatching, 34—Period of growth, 35—Habits of the young, 36—Habits at maturity, 36—Causes of occasional over-increase, 37—Checks, enemics, 38—The efforts of experts baffled, 42—Tree culture, 42—Climatic checks, 43—Tree planting, 43—Flights affected by storms, 45—Number of annual broods, 46—Sub-permanent region, 46—Its relations, 47—Locust movements therein, 48—How to fight this insect, 49—Locust probabilities, 49—Recent swarms, 50.

CHAPTER IV:

Notes on other locusts and on the Western cricket, Anabrus. By Lawrence Bruner

Other locusts, 53—List of North American locusts north of Mexico, 55—The Western cricket, 61.

CHAPTER V:

Data obtained from solar physics and carthquake commotions applied to elucidate locust multiplication and migration, by A. H. Swinton

Importance of the central luminary, wide effects of variation in its potential energy, 65—On the tides, on chemical and organic activities, on the physical forces, on cpidemics, etc., 65, 66—Periods of sun-spots, electrical effects of, 66—Thermometric effects of, 67—Volcanic effects of, 68—Entomological effects of, 68—A new sun-spot table, 69—Statement and tabulation of sun-spot dates, 69—Comparison of sun-spot periods and insect periods, 73—American locusts diminished by the spots, 73—European confirmative examples of locusts and other insects, 74—Tabulation of rare insect captures, 79—Explained, 81—Migration and distribution affected, 81—Locusts again predicted in four or ten years, 83—Conclusions, 84.

PART II.—THE ARMY-WORM, CANKER-WORMS, AND THE HESSIAN FLY.

CHAPTER VI: By C. V. Riley.

53

65

CHAPTER VI—Continued.

tent of country injured, 133—Crops injured, 133—Amount of damage, 133—Previous season and crop, 134—Number of broods, 134—An accompanying cut-worm, 135—Natural enemies, 135—Army-worm correspondence in spring of 1882, 136—The invasion of 1880 in New Jersey, by Rev. Samuel Lockwood, 139—Performances of the worms, 139—The direction of travel, 141—Origin, eggs, etc., 143—Breeding spots, 143—Recapitulation, habits, and remedics, 145—Notes from Prof. C. V. Rilcy, the number of broods, hibernation, seasonal influences, 147—Bibliography, 148.

CHAPTER VII: By C. V. Riley.

Canker-worms, Paleacrita vernata, Anisopteryx pometaria. By C. V. Riley

Classification, 157—Two distinct insects concerned, 158—Differences between them, 159—Nomenclature, 162—Past history, 165—The spring canker-worm, 170—Range of the species, 170—Characters, 171—Habits and natural history, 172—Appearance of the worms, 173—Food-plants, 174—Modes of distribution, 175—Enemies, 175—Destructiveness of canker-worms, 178—The fall canker-worm, 179—Range of this species, 179—Its characters, 180—Habits and natural history, 181—Oviposition, 182—Season of appearance, 182—Pupation, 183—Food-plants, 183—Remedies and preventive measures applicable to both species, 183—Sticky substances, 183—Hanging tin band, 185—Troughs of oil, 186—Precautions and classification of contrivances, 189—Jarring and burning, 191—Washes and dustings, 191—Paris green, 192—Muriate of lime, 192—Sulphur plugged in trees, 193—Fall plowing, 193—Birds and parasites, 195—Different measures against the species, 196.

CHAPTER VIII:

PART III.—SCIENTIFIC RESULTS.

CHAPTER IX:

Descriptions of larvæ of injurious forest insects. By A. S. Packard, jr. Buprestidæ, 251—The flat-headed apple-borer, Chrysobothris femorata, 251—Chalcophora virginica, 252—Melanophila sp., 253—The flat-headed spruce-borer (Melanophila?), 254—The flat-headed peach- and cherry-borer, Dicerca divaricata Say, 255—Buprestid under hemlock bark, 255—Cerambycidæ, 256—Longicorn larva under hemlock bark, 256—Saperda on the willow, 256—The lesser pine-borer, Asemum mæstum Haldeman, 256—The oak-borer, Elaphidion parallelum Newman, 257—The common cak clytus,

157

198

263

271

273

277

2267. [RILEY, C. V., et al.]—Continued.

CHAPTER IX-Continued.

Xylotrechus colonus Fabr., 257—Clytus (?) larva on blaek-birch, 259—The ribbed rhagium, Rh. lineatum Oliver, 259—The lesser prionus, Orthosoma brunneum De Geer, 260—Unknown longicorn borer from an oak log, 261—Unknown longicorn borer in sycamore, 262.

CHAPTER X:

The embryological development of the locust. By A. S. Packard, ir. Formation of the blastoderm, 263—Origin of the primitive amedoid eells, 264—The blastodermic disc or primitive band, 264—The germinal groove, 264—Origin of the eellular or germinal layers, 264—Diagrammatic view of the origin of these layers, from Graber, 265—Embryonal membranes, 265—Serous membrane, 265— Amnion, 265—Tabular view of the eight embryonal layers, 265— Division of the embryo or primitive band into body segments, 266—Development of the appendages, 266; of the nervous system, 266; of the alimentary canal. 266—The stomodæum, 266— The proctodeum, 267—The mesenteron or stomach, 267—The pyloric appendages of the stomach, 267-The Malpighian tubes, 267—The genital glands, 267—The tracheæ and salivary glands, 267—Spinning glands, 267—Development of the wings, 268—Critieism of Gegenbaur's views, 269—Fritz Müller's views, 269— Speculation on the primary origin of wings, 270—Correlation of metamorphosis with the acquisition of wings, and of the latter with the ripening of the sexual organs, 271—Differentiation of mesonotum and metanotum dependent on the presence of wings, 271—Development of a pupal stage, 271—Genealogy of the orders of insects, 271.

The embryological development of orthopterous insects. By A. S. Packard, jr

Rapid development of the embryo in autumn, 272—Embryology of Gryllotalpa vulgaris, 272—Formation and contraction of the embryonal skins, 272—Formation of the digestive caual, 272—Functions of the mid-gut and hind-gut, 273—Origin of the Malpighian vessels, salivary glands, tracheæ, and ovaries, 273—The sections of eggs studied by the author, prepared by N. N. Mason, 273.

CHAPTER X-Continued.

The development of the bark-boring beetles *Hylurgops* and *Xyle-borus*. By A. S. Packard, jr.

280

282

286

Breeding habits of Xyleborus calatus, 280—Oyiposition of Hylurgus pinifex, 280—Egg of the Xyleborus, 24 hours after impregnation, 280—More developed egg, 280—Threads connecting the amnion with the serous membrane, 281—Dorsal view of the embryo, 281—Later stage in Hylurgops, 281—Later stage in Xyleborus, 281—Number of pairs of spiracles, 281—Structure and habits of freshly hatched Hylurgops, 281.

The number of segments in the head of winged insects. By A. S. Packard, jr

Views generally entertained on this subject, 282—Head composed of four segments, 283—The procephalic lobes form the antennal segment, 284—The clypeus and labrum are the tergal portion of this segment, 284—The epicranium is the pleural portion, 284—The occiput is the tergal portion of the labial segment, 284—The gular region probably the base of the labium, 285.

CHAPTER XI:

The systematic position of the Orthoptera in relation to other orders of insects. By A. S. Packard, jr.....

Review of the characteristics of the four lowest orders of winged insects, 286—Probable descent of Orthoptera, Pseudoneuroptera, and Dermatoptera from a Thysanuran form, 286-Method of the present study, 286—Characters of the Phyloptera, 287—Mouthparts, 287-Thorax, 287-Wings, 288-Abdomen, 288-Metamorphosis, 288-Nomenclature of external parts of Arthropoda, 288-Sequence of orders of Phyloptera, 288—Characters of Dermatoptera, 289; of Orthoptera, 289; of Pseudoneuroptera, 290-Want of uniformity in characters of Pseudoneuroptera, 290-The diagnostic characters superficial, 290—Structure of labium, 291— Relative proportions of head-parts, 291—Subdivisions of the order, 292—Characters of Platyptera, 292; of Odonata, 292—Of Ephemerina, 293—Characters of Neuroptera, 293—Ligula, 293—Thorax, 293—Wings, 293—Abdomen, 294—Subdivisions of the order, 294— Tabular view of the grand divisions of winged insects, 294-Diagrammatic view-of the genealogy of the insects, 295—Genealogy of the Hexapoda, 295; of Thysanura, 295; of Dermatoptera, 295; of Orthoptera, 296; of Pseudoneuroptera, 296; of Hemiptera, 297; of Neuroptera, 297—Derivation of cruciform larvæ, 297; of Sialidæ, 297; of thysanuriform larvæ, 297-Diversity of Neuropterous larvæ, 297-Larvæ of Sialidæ, 297; of Hemerobiidæ, 297; of Trichoptera, 298; of Panorpidæ, 298-Significance of the abdominal legs of Panorpid larvæ, 299-The hypermetamorphosis of Mantispa a key to the origin of a complete metamorphosis, 299-The aspect of the Neuroptera comparatively specialized and modern, 299—The Neuropterous labium a secondary product, 299—Origin of the Coleoptera, 299—The free, active larvæ of the carnivorous groups most nearly allied to the primitive form, 300—The scavenger and phytophagous larva show increasing degradation of development, 300-The relative form of the maxillæ a good index of the general development of the body in Coleoptera, 300-Structure of maxillæ in the several families, 300-Close resemblance of the mouth-parts of Elaterid

CHAPTER XI-Continueds

larvæ to those of Carabid larvæ, 301-The hypermetamorphosis of Meloidæ furnishes a clew to the probable origin of the different types of Colcopterous larvæ, 301-Brief description of the metamorphoses of Meloe, 301-Of Epicauta, 302-Comparison of the larval stages of Meloidæ with the different types of Coleopterous larvæ, 302-Origin of the Diptera, Lepidoptera, and Hymenoptera, 303—Presence of temporary abdominal appendages on each segment of Lepidopterous, Coleopterous, and Orthopterous embryos, 304—Description of the structure of Dermatoptera (Forficula), 304-Labia, 308-Larva of Forficula, 308-Orthoptera, 309—Blattariæ, 309—Close relation of Blatta and Termes, 310— Structure of Mantidæ, 310-Blatta the stem-form of the Orthoptera, 312-Mantis connects the Acrydii and the Blattariæ, 312-Structure of Phasmida, 312—Diapheromera, 312—Prisopus, 313— This genus connects the Phasmida with the Acrydii, 314-Structure of Acrydii, 314—Caloptenus, 314—Proscopia, 316—Conocephalus, 318-Locustaria (Anabrus), 318-Gryllida, 319-Gryllus, 319—Gryllotalpa, 320—Œeanthus, 322—Sequence of families of Orthoptera, 322—Structure of Pscudoneuroptera, 322—Corrodentia, 322-Perlidæ (Pteronarcys), 322-Psocidæ (Psocus), 325-Termitidæ, 326-Odonata, 329-Ephemerina, 333-Neuroptera, 335-Planipennia, 335-Sialidæ, 335-Hemerobiidæ, 338-Panorpidæ (Panorpa), 342-Trichoptera (Lininophilus), 344.

CHAPTER XII:

Note on the geographical distribution of the Rocky Mountain locust, illustrated with a colored zoö-geographical map of North America. By A. S. Packard, jr......

346

[3]

[6]

[8]

APPENDICES.

APPENDIX I:

Early references to the occurrence of the Hessian fly in North America.

Early newspaper references, [3]—In 1784-'85, at Long Island and
West Chester, [3]—In 1788, in Long Island, Delaware, New Jersey, and Pennsylvania, [3]—The fly resisted by certain varieties of wheat, [3-5].

APPENDIX II:

- B. Observations of the new crop gall-gnat. By Dr. Balthasar Wagner, Fulda, 1861. Translated by Carl F. Gissler.....

Literature, [8]—Preliminary consideration of the subimago and imago stages, [8]—Occurrence at Fulda, [8]—Breeding under nets, [9]—Seasonal and sexual differences, [9]—The fly short lived, [10]—Description of the adult, [10]—Of the male, [12]—Of the female, [13]—The winter generation, [14]—The egg, [15]—The maggot, [16]—The pupa or chrysalis, [19]—The perfect insect, [20]—Seasonal and sexual differences, [21]—Only the maggots hibernate, [22]—Single and double brooded, [23]—Summer generation, [23]—The new cereal gall-gnat as compared with Cecidomyia secalina Loew, [24]—Comparison of C. secalina Loew with C. destructor Say, [24]—Views of Loew, Fitch, and Herrick.

APPENDIX II—Continued.

discussed, [25]—The rye gall-gnat, C. secalina and C. destructor, all of one species, [28]—The name Hessian fly not justified, [28]—Historical evidence thereon, [28]—Attempt at a new theory of the origin of the insect, [31]—Of Asiatic origin?, [31]—Carried to North America from France or Spain, [32]—Parasites, [33]—Remedies, [34]—The removal of the barley after-growth, [34]—Picking by hand, [35]—Grazing, [35]—Mowing, etc., [36]—Sowing grain as bait, [36]—Plowing, burning, rolling, [37]—Richness of soil, [37]—Resistive varieties of wheat, [38]—Sowing late or early, [38].

APPENDIX III:

The Hessian fly in Silesia in 1869. By Professor Dr. Ferd. Cohn.... [39]
The Hessian fly, *Cecidomyia destructor*, and other Diptera observed destroying grain in Silesia, [39].

APPENDIX IV:

The Hessian fly recently detected extensively distributed in Russia, [41]—Two generations each season, [41]—Number of eggs at each deposition, [41]—Parasites, [41]—Doubtful species of Cecidomyia, [42]—Descriptions of species, [42].

APPENDIX V:

Conclusions of the author and others as to the nativity of Cecidomyia destructor, [43]—Not imported by the Hessian troops, [43]—
Probably here before the war, [45]—Not known in Germany before 1857, [46]—Mistakes corrected, [46]—A critical review of
its history, [47]—Marked difference between C. destructor and C.
secalina, [49].

APPENDIX VI:

Report on the Rocky Mountain locust in 1880. By John Marten... Report to Dr. Cyrus Thomas of observations, [50]—Damages from locusts and wheat flies in Minnesota, [50]—Previous damages from locusts in Dakota, [51]—In Montana, [52].

[50]

[55]

APPENDIX VII:

APPENDIX VIII:

Dates and localities of locust swarms that have attained the coast of Great Britain. Notes by A. H. Swinton, [57]—Occurrence of the migratory locust in Japan, [59]—Early notices of locusts in North America, [59]—Locusts at Fort Frontenac or Cataraqui, Canada, [59]; at Puerto Seguro Harbor, California, [60]—The locust in California in 1880, [60]; in Nevada, [60]—Locusts and coffee-trees in Guatemala, [60]—History of locusts in other countries, [61]—Locusts probably never will do so much damage again in North America, [61]—Utilization of locusts as food, [61]; of crickets as manure, [62]—Ravages of locusts in Russia in 1879 and 1880, [62]; in the Caucasus, [63]; in Elizavetopol, [63]; in the district of Rassachs, [64]—Plowing as a means of killing the

APPENDIX VIII—Continued.

locasts, [64]—Destruction of locasts in the district of Gori, [64]—Locusts in China in 1878, [65]—Oil as a means of destroying them, [65]-2,000,000 catties of eggs collected, [65]-Article on the extermination of the locust in China, [65-68]—Chinese bibliography of the subject, [65]—Chinese regulations regarding the extermination of locusts, [67]-Locusts in Cape Colony, South Africa, [68]—Ravages of nnfledged locusts, [68]—Locusts as food, [69]—Barriers as means of warding off and capturing locusts, [69]—Egg laying and hatching, [72]—Protection against flying locusts, [72]—Locusts in the Philippine Islands in 1878-'79, [72]—The locust plague in Bolivia, [72]—Locusts in India, [73]; in the Ottapidaram Taluq, [73]; in Sattur, [73]—Request of the Madras Government for information as to the movements of the locusts, [74]—Locusts in Kulapurum, Comaralingum, Kolumam, and in the Pulni Taluq, [74]—Difficulties in coping with locusts in southern India, [74]—Means against unfledged locusts, [75]— Locusts driven into the sea at Tuticorin and Trichendur, [76]— Recommendation of rewards for gathering locusts, [76]—Locusts at Peryapatty, [76]; in the Bellary district, [77]—Recommendation of means against locusts, [78]—Locusts at Madura, [78]— Destruction of young locusts by fire at Pothanore, [78]—Locusts at Tiroomoorthypovil, Dhully, and Jellipatty, [79]; in various parts of the Punjab, [79]—Means against the locusts, [79]—Remuneration for destruction of locusts, [80]—Locusts in Cnddapah district, [81].

APPENDIX IX:

Experience with the spring canker-worm. By Martin A. Howell, jr.. Means employed to save an orchard from the ravages of cankerworms, [82]—Pruning, manuring, and washing of the trees, [82]—Spraying of the leaves with soft soap and arsenite of soda in water, [83]—Belting the trees with sulphnr and adhesive mixtures, [83]—Endurance of the moths, [84]—Variation of color in eggs, [85]—Most practical means of relief, [85]—Adulterations of Paris green, [85]—Good force-pnmps, [85]—Distribution of canker-worms by the wind, [85].

- 2268. [RILEY, C. V.] The silk industry in the United States. <Sci. ence, 7 March, 1884, v. 3, pp. 290-292. S.-b. No. 42, pp. 82-84-Review of article of W. C. Wyckoff; early history and present status of silk culture in North America; criticism of the estimates of the amount of silk raised in early times; relations of silk-culture and silk-manufacture to free trade and protection in the United States.
- 2269. RILEY, C. V. The army-worm. < Washington, 1883 [20 March, 1884], pp. 89–152, pl. 1–2, with 1 p. expl. of pl. Advance reprint of principal portions: <Rept. [U. S.] Com. Agric. for 1881–1882, January, 1883, pp. 89–106.

Chapter 6 of the Third Report U. S. Entomological Commission. See No. 2267 for synopsis of contents.

2270. RILEY, C. V. Canker-worms. < Washington, 1883 [20 March, 1884], pp. 157–198, pl. 3, with 1 p. expl. of pl. Extract: < Amer.

- 2270. RILEY, C. V.—Continued.
 - Cult., 1884, v. 46: 29 March, p. 1; 5 April, p. 1. S.-b. No. 42, pp. 118–119; No. 57, pp. 86–87.
 - Chapter 7 of Third Report U. S. Entomological Commission. See No. 2267 for synopsis of contents.
- 2271. [RILEY, C. V.] The chinch-bug. Another entomologist expresses his views regarding the farmers' pest. <Watertown [N. Y.] Daily Times, 27 March, 1884, v. 24. S-b. No. 57, pp. 42-43. Reprint: <Ogdensburg Journal, 1884.
 - Notice of articles of J. A. Lintner; the occurrence of *Blissus leucopterus* in New York State in unusual abundance in 1882 and 1883 not a cause for great alarm; means against this insect. See No. 2252.
- 2272. RILEY, C. V. Notes on North American *Psyllidæ*. <Proc. Biol. Soc. Wash., 1884, v. 2, pp. 67–79. Separate: <[Washington], 10 April, 1884, pp. 67–79.
 - Notice of earlier studies of European and North American Psyllidw; list of 18 nominal species described from North America; systematic and synonymical list of the same; descriptions of Pachypsylla n. g. [p. 71], P. venusta, P. celtidis-mamma, and P. [Blastophysa n. s-g., p. 75] celtidis-gemma n. sp. [p. 74]; dichotomic table of the three species of Pachypsylla; descriptions of Ceropsylla n. g. [p. 76] sideroxyli n. sp. [p. 76], Rhinopsylla n. g. [p. 77] schwarzii n. sp. [p. 78].
- 2273. RILEY, C. V. Remarks on the bag-worm, *Thyridopteryx epheme-ræformis*. <Proc. Biol. Soc. Wash., 1882–1884, 1884, v. 2, pp. 80–83, figs. 1–3. Separate: <[Washington], 10 April, 1884, pp. 80–83, figs. 1–3.
 - Explanation of the more important and less known facts in relation to the life-history of *Thyridopteryx ephemeræformis*, especially in reference to the act of coition; figures of all stages and of the external male genitalia.
- 2274. RILEY, C. V. The use of naphthaline as an insecticide. <Science, 11 April, 1884, v. 3, pp. 455-456.
 - Review and summary of article of E. Fischer.
- 2275. RILEY, C. V. Entomography of Hirmoneura. <Science, 18 April, 1884, v. 3, p. 488.
 - Review of F. Brauer's observations on the life-history of Hirmoneura obscura.
- 2276. RILEY, C. V. Orthoptera. <Standard Natural History (The) ...
 Boston, Cassino, 1 May-27 July, 1884, v. 2, pp. 167-203, 1 pl.,
 figs. 243-285. Reprint: <Riverside Natural History (The) ...
 Boston and New York, Houghton, Mifflin & Co., 1888, v. 2, pp.
 167-203, 1 pl., figs. 243-285.
 - Systematic position, limitations, classificatory characters, transformations, molts, and geographical and geological distribution of the order; characters, habits, and distribution of the several families; mention of the principal North American and of some forcign species, with more particular accounts of many species, especially in regard to their stridulation and oviposition, with figures of a few species and of their eggs. Ectobia germanica, Phasmomantis carolina, Diapheromera femorata, Ecanthus niveus, E. latipennis, Orocharis saltator, Camptonotus scudderi, Cyrtophyllus concavus, Microcentrum retinerve, and Caloptenus spretus are particularly mentioned.
- 2277. RILEY, C. V. Reports of observations and experiments in the practical work of the division, made under the direction of the

entomologist, together with extracts from correspondence on miscellaneous insects. <Bull. No. 4, Div. Ent. U. S. Dept. Agric. [3 May], 1884, 102 pp., 4 figs.

CONTENTS.

Introduction	7
Report upon cranberry and hop insects. By John B. Smith	9
Observations on the Rocky Mountain locust and other insects in the	
Northwest during the summer of 1883. By Lawrence Bruner	51
Preliminary report of observations upon insects injurious to cotton,	
orange, and sugar-cane in Brazil. By John C. Branner	63
Report on the effects of cold on the scale insects of the orange in	
Florida. By Joseph Voyle	70
Extracts from correspondence	74
Additional notes on the cultivation of pyrethrum in the United States.	95

2278. RILEY, C. V. Silk-culture in the colonies. <Science, 9 May, 1884, v. 3, pp. 562-563.

Explanation of several technical terms used in silk-culture.

2279. RILEY, C. V. Maple-tree insects. <Sci. Amer., 24 May, 1884, v. 59, p. 325. S.-b. No. 63, p. 45.

Answer to inquiry of R. H. Cutler; food-plants of and means against *Pulvinaria innumerabilis*.

2280. RILEY, C. V. Acronycta betulæ n. sp. <Bull. Brooklyn Ent. Soc., May, 1884, v. 7, pp. 2-3, fig.

Description of Acronycta betulæ n. sp. [p. 2], reared from Betula nigra; seasons and habits of larva; comparison with related species.

- 2281. RILEY, C. V. Recent advances in economic entomology. <Proc. Philos. Soc. Washington, 1884, v. 7, pp. 10-12. Separate: <Washington, 1884, pp. 10-12. Reprint: <Kansas City Review of Science, May, 1884, pp. 13-15. S.-b. No. 63, pp. 43-44. Characteristics of economic entomology as a science and difficulties of its
 - Characteristics of economic entomology as a science and difficulties of its prosecution; the chief insecticide substances and their applicability; recent progress in mechanical appliances.
- 2282. RILEY, C. V. Department of insects. <Ann. Rept. of Regents Smith. Inst. for 1882, 1884, pp. 215-216.
 - List of accessions to the U.S. National Museum, with comments; note on the Glover plates and manuscripts.
- 2283. RILEY, C. V. Quelques mots sur les insecticides aux États-Unis et proposition d'un nouveau remède contre le Phylloxera. Communication faite à la Société d'Agriculture de l'Hérault à la seance du 30 Juin, 1884. <Montpelier, 1884, t.p. cover+8 pp. Reprint: <Messager Agricole, 10 July, 1884, ser. 3, v. 5, pp. 255-265. <La Vigne Amer., July, 1884, v. 8, No. 7, p. 207.
 - Resistant vines better than the best insecticides as a means against *Phylloxera* vastatrix; principal insecticides in use, especially arsenical substances, petroleum and pyrethrum; method of action and mode of application of these insecticides, especially application by the cyclone nozzle, and the subterranean application of kerosene emulsion as a means against *Phylloxera vastatrix*; tobacco vapor, under favorable conditions, would probably kill this insect.

340	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY	
2284.	[RILEY, C. V.] Remedies for various insects. < Prairie Farm 26 July, 1884, v. 56, p. 470. Sb. No. 59, p. 36. Comparative efficacy and other qualities of several insecticides, especial arsenical substances, tobacco vapor, petroleum emulsions, and pyrethruses.	ally
2285.	RILEY, C. V. Habits of Grapholitha olivaceana. <ent. 1884,="" 21,="" 67.="" august,="" bred="" curled="" from="" grapholitha="" ma.="" mo.="" of="" olivaceana="" p.="" solidago.<="" td="" tips="" v.=""><td>ıg.,</td></ent.>	ıg.,
2286.	RILEY, C. V. Rhyssa not lignivorous. Science, 28 November 1884, v. 4, p. 486. Parasitic habits of Rhyssa [= Thalessa] upon Tremex columba.	er,
2287.	RILEY, C. V. The apple-root borer. <rural 13="" 1884,="" 43,="" 50.="" 63,="" 831.="" <i="" against="" cember,="" larva="" learning="" means="" new-yorker,="" no.="" of="" p.="" sb.="" the="" v.="">Prionus laticollis.</rural>	De-
2288.	RILEY, C. V. A new insect injurious to wheat. <bull. 111-112.="" 111.="" 1884,="" 1884],="" 7,="" <[brooklyn,="" <i="" and="" brolyn="" december,="" description="" ent.="" female="" image="" larva,="" n.="" of="" p.="" pp.="" pupa,="" separa="" soc.,="" v.="" y.,="">Isosoma grande n. sp. [p. 1 reared from wheat stems in Indiana.</bull.>	te:
2289.	RILEY, C. V. The insects of the year. <science, 1884,="" 26="" 4,="" 565-568.="" calendar="" december="" each="" entomological="" for="" month="" of="" pp.="" td="" the="" v.="" year.<=""><td>er,</td></science,>	er,
2290.	RILEY, C. V. On the care of entomological specimens. < see ence, 9 January, 1885, v. 5, p. 25. Reply to editorial comments on the care of entomological specimens in National Museum.	
2291.	RILEY, C. V. Report of the entomologist. <ann. 10="" 1884,="" 1885,="" 285-41="" 285-418,="" 31="" 9+pp.="" 9,="" <washington:="" [u.="" agric.="" commissioner="" for="" january,="" pl.="" pl.<="" pp.="" rate:="" rept.="" se="" td=""><td>pa-</td></ann.>	pa-
	Introduction Scope of the report, 285—Climate and insect phenomena during the year, 285—Silk-culture, 286—Field force of the Bureau, 286—Suggested additions to field force, 287—Office force and its work, 287—Trip to Europe, 287—International Forestry Exhibition at Edinburgh, 287—Phylloxera in France, 288—Exhibit of the Bureau at the New Orleans Exposition, 288—Third report of the United States Entomological Commission, 288—Special reports, 288—Suggestions, 288—Illustrations, 288.	285
	Cabbage cut-worms General characters, habits, and natural history, 289—Natural en-	289 289
	emies, 290—Damage to cabbage, 290. The dark-sided cut-worm, Agrotis messoria. Synonymy, 290—Spread, 290—Climbing habits, 290—Natural his-	290
	tory, 290—Larva and moth popularly described, 290. The granulated cut-worm, Agrotis annexa	291
	The shagreened cut-worm, Agrotis malefida	292

Distribution, 292—Points of difference from Agrotis annexa, 293—

Early states described, 293.

2291 Rile	EY, C. V.—Continued.	
	BAGE INSECTS—Continued.	
	The W-marked eut-worm, Agrotis clandestina	293
	Food-plants, 293—Habits, 294—Larva and moth popularly described, 294.	2017
r	The greasy eut-worm, Agrotis ypsilon	294
	Synonymy, 294—Distribution, 294—Habits, 294—Food - plants, 294—Hibernation, 295—Larva and moth popularly described, 295—Description of egg, 295.	
Г	The speekled eut-worm, Mamestra subjuncta	296
	Distribution, 296—Habits, 296—Appearance, 296.	
Γ	The glassy eut-worm, Hadena devastatrix	296
Т	The variegated eut-worm, Agrotis saucia. Distribution, 297—Food-plants, 297—Egg and early states described, 298—Rapidity of development, 298.	297
Rei	medies for eabbage eut-worms	298
	Unsatisfactory remedies, 298—Wrapping the stem, 298—Trapping the worms, 299—Ditching, 299—Fall plowing, 299—The great value of poisoned bait, 299.	
	her eabbage insects	300
7	The imbricated snout-beetle, <i>Epicarus imbricatus</i>	300
Т	The wavy-striped flea-beetle, Phyllotreta vittata	301
	Food-plants, 301—Habits, 301—Injury, 302—Early states, 302—	001
	Remedies, 302.	
Z	Zimmermauu's flea-beetle, Phyllotreta zimmermanni	304
	Food-plants, 304—Habits of larva, 307—Characters as compared with the wavy-striped flea-beetle, 305—Early states described, 306—Other species of the group, 307—Their habits, 307—Nat-	
rr.	ural enemies, 307—Pleurotropis phyllotretæ n. sp. deseribed, 308. The Colorado eabbage flea-beetle, Phyllotreta albionica	308
1	How it differs from the preceding species, 308—Mention of a Pacific coast species, 308.	900
Т	The Colorado potato-beetle, Doryphora 10-lineata	308
r	The harlequin cabbage-bug, Murgantia histrionica	309
r	The tarnished plant-bug, Lygus pratensis	312
	Distribution, 312—Food-plants, 312—Synonymy, 312—Supposed	
	earnivorous habit, 312—Effects of its puncture, 313—Variation in color, 313—Habits, 313—As an enemy to strawberries, 314—Percenting recourses, 215—Applications, 215	٠
п	Remedies, 314—Preventive measures, 315—Applications, 315. The false elinch-bug, Nysius angustatus	315
1	Synonymy, 315—Food-plants, 316—Habits, 316—Remedies, 316.	010
Т	The eabbage plant-louse, Aphis brassica	317
	An imported insect, 317—Damage, 317—Habits, 317—Life-history, 318—Natural enemies, 318—Remedies, 318.	
r	The eabbage Authomyia, Anthomyia brassica	319
	Introduced from Europe, 319—Life-history, 320—Natural enemies, 320—Remedies, 321.	

2001	D C V C	
2291.	RILEY, C. V.—Continued.	
	CABBAGE INSECTS—Continued.	
	The cabbage Oscinis, Oscinis brassica	322
	First found at St. Louis, 322—Habits, 322—Descriptive, Oscinis	
	brassicæ n. sp., 322.	000
	The Rocky Mountain locust, Caloptenus spretus	323
	Successful introduction of a parasite (Apanteles glomeratus) of the	
	imported cabbage-worm	323
	GENERAL TRUTHS IN APPLIED ENTOMOLOGY	323
	Introductory, 323—Benefits derived from insects, 324—Influence	
٠	of civilization upon insect increase, 324—Lossos from insects,	
	324—Knowledge which the economic entomologist should pos-	
	sess, 325—Habits of the grape phylloxera, 325—Number of	
	species of insects, 326—The different insecticides, 326—Arsen-	
	ical compounds, 327—Petroleum, 327—Kerosene emulsions,	
	328—Pyrethrum, 328—Its cultivation in California, 328—Its	
	influence on mammals, 329—Insecticides to be used against	
	root-feeding insects, 329—Mechanical contrivances, 329—The	
	eddy-chamber or cyclone nozzle, 330. Kerosene emulsions	330
	Their importance as insecticides, 330—Use in orange groves,	330
	330 —Warning against imperfect emulsions, 331—Formula for	
	kerosene and soap emulsions, 331—Importance of a stable	
	emulsion, 331—Use of a force-pump in making emulsions, 331—	
	Trees injured by unemulsified oil, 332—The use of ley washes	
	in California, 333.	
	MISCELLANEOUS INSECTS	334
	The American Cimbex, Cimbex americana	334
	Injury to willows, 334—A new habit, 334—Eggs and mode of	
	oviposition, 335—Habits of young larva, 335—Remedies, 336—	
	Other willow enemics, 336.	
	The streaked cottonwood leaf-beetle, Lina scripta	336
	Value of the cottonwood in the West, 336—Damage by the beetle	
	in 1884, 336—Other food-plants, and former injuries, 337—	
	Natural history, 338—Remedies, 338—Apparatus for poisoning,	
	339—Varieties of the beetle, 340.	
	The Southern buffalo-gnat, Simulium. pecnarum	340
	Losses in former years, 340—Damage done by European species,	
	341—Early states of European species, 342—Early states of	
	American species, 342—Recent ravages in the Southwest, 343—	
	Remedies, 344—Smudges, 344—Body applications, 344—Sug-	
	gestions, 345.	
	The angoumois grain-moth, Gelechia cerealella	345
	Abundance, 345—Origin, 345—Past history, 346—Natural history	
	and method of work, 346—Number of annual generations, 347—	
	Results of its work, 347—Parasites, 348—Remedies, 348—	
	Baking and concussions, 348—Quarantine, 348—Bisulphide of	
	carbon, 349—Preventive measures, 349—Description of eggs,	
	350,	250
	The cottony maple scale, Pulvinaria innumerabilis	350
	Distribution, 350—History and synonymy, 350—Life history, 351—	
	Food-plants, 352—Mode of spreading, 353—Natural enemies,	
	354—Remedies, 355. The graphermy fruit worm. Acrehasis againsi	255
	The cranberry fruit-worm, Acrobasis vaccinii Natural history, 355—Remedies, 356—Descriptive, Acrobasis vac-	355
	cinii n. sp., 356—Systematic position, 357.	
	other in spa, ood by significant posterior, not.	

91. I	RILEY, C. V.—Continued.	
	MISCELLANEOUS INSECTS—Continued.	
	The larger wheat-straw Isosoma, Isosoma grande	;
	The phytophagic habits of the genus, 357—Mr. Webster's obser-	
	vations, 357—Descriptive, Isosoma grande n.sp., 358—Oviposition,	
	358—Only females known, 358.	
	SILK CULTURE	;
	Sending out eggs, 359—Reports, 359—The manual of instructions,	
	359—Appointment of Philip Walker as special agent, 359—	
	Diseases, 359—Pasteur's process, 359—A Government reeling	
	establishment, 360—Correspondence; 360—The Serrell reel,	
	360—Home reeling, 361—Herbelin's filature, 361.	
	REPORTS OF AGENTS	;
	Rust of the orange. By H. G. Hubbard	
	Discoloration of the fruit, 361—Time of appearance, 361—It is	
	not a fuugus, 362—The mite on the leaves, 362—First appear-	
	ance of mites on the fruit, 363—Attacks of mites always fol-	
	lowed by rust, 363—Interval between disappearance of mites	
	and appearance of rust, 363—Description of the mite, 363—Life	
	history, 364—Food, 364—Wandering habits, 365—Numerical	
	abundance, 365—Rings of rust on fruit, 366—Influence of	
	weather, 366—Agencies which assist in the distribution of the	
	mites, 366—Food-plants, 367—Effects of attacks on foliage, 367—	
	Rusted fruit, 367—Introduction and spread of the mite, 367—	
	Periods of increase, 368—Geographical distribution, 368—Reme-	-
	dies, 368—Influence of soil and methods of cultivation, 368—	
	Preventive measures, 368—Application of insecticides, 369—	
	Whale-oil soap, 369—Sulphur, 370—Kerosene, 372—Creosote,	
	372—Potash, 372—Pyrethrum, 373—Lime, 373—Ashes, 373—	
	Cantion, 373.	
	Second report on the causes of the destruction of the evergreen and	
	other forest trees in northern New England and New York. By A.	
	S. Packard, jr	
	Localities visited, 374—Destruction of spruces in northern New	
	York, 374—Probably caused by bark-boring beetles, 375—Destruction of spruces in northern Maine, 376—Discussion of	
	causes, 376—Condition of the hackmatack in 1884, 377—The	
	reddish-yellow spruce-bud worm, 378—Its injuries, 378—Its	
	classification, 378—Its habits, 378—Remedies, 379—The com-	
	mon longicorn pine-borer, 379—Abundance, 379—Mode of ovi-	
	position, 380—Description of egg, 380—Habits and damage	
	done, 381—The sugar-maple borer, 382—Its burrows, 382—Its	
	eggs and larvæ, 382—Kerosene as a remedy, 383.	
	Insects affecting fall wheat. By F. M. Webster	
	The wheat-straw Isosoma, 383—Oviposition, 384—Description of	
	its egg, 385—Number of generations, 385—Effect of its work on	
	wheat-straw, 386—Table o position of larvæ in straw, 386—	
	Natural enemies, 387—Wheat and grass saw-fly, 387—Appear-	
	ance, 388—Parasites, 388—Wheat midge; 389—Natural enemies,	
	389—Dates, 389—American Meromyza, 389—Dates, 390—The	
	flea-like negro-bug, 390—Dates, 390—Description of egg, 390—	
	The tarnished plant-bug, 391—The soldier-bug, 391—The field-	
	cricket, 391—Cutting off wheat, 391—Insects affecting rye, 392—	
	Insects affecting oats, 392—The stalk-borer, 392—Dates, 393.	

2291. I	RILEY, C. V.—Continued.	
	REPORTS OF AGENTS—Continued.	
	Report upon insects affecting the hop and the cranberry. By J. B.	
	Smith The hop Aphis, 393—Its first appearance, 393—The cranberry fruit-worm, 394—The fire-worm, 395—Cranberry leaf-folder, 395—Its dimorphism, 396—A cecidomyid larva infesting pears at Meriden, Conn., 396—Description of larva and infested fruit, 396—Pupation, 397—Distribution, 397—Varieties of pear infested, 397—Probably an imported pest, 398—Remedy, 398.	393
	Notes from Nebraska. By Lawrence Bruner The Rocky Mountain locust in 1884, 398—Other species of locusts, 399—The snowy tree-cricket, 399—The chinch-bug, 399—Cutworms, 400—The imported cabbage-worm, 400—Its natural enemies, 400—Other cabbage-worms, 401—False caterpillars on grass and scdge, 401—Cimbex americana on willow hedges, 401—injury to cottonwoods by the striped beetle, 402—The Colorado potato-beetle, 403.	398
	± /	403
	Chinch-bug notes The so-called "invasion" of St. Lawrence County, New York, in 1883, 403—Mr. Lintner's prediction and his reasons therefor, 403—Reasons for dissenting from this view, 404—The result in	403
	Notes on the grape Phylloxera	405
	Regarding the grape Phylloxera in Russia, 405—In reference to the treaty of Berne and the prohibition of the introduction of bulbs and cuttings from the United States into Germany, 407—The grape Phylloxera in graperies; legal questions arising, 408.	
		41 0
	The box Psylla found in the United States, 410—The dwarfing of oaks by Mallodon melanopus, 410—The clover-seed midge, 411—The potato-stalk weevil, 411—The red-humped prominent, 411—The varying anomala, 412—White-lined morning-sphinx, 412—The apple-tree tent caterpillar, 412—Corn bill-bugs, 413—A swarming mite, 413—A new enemy to white roses, 413—A beetle eating peach-leaves, 413—Effect of cold on the eggs of barklice, 413—Ravages of grain-weevils in Florida, 414—Fuller's rose-beetle, 414—The blood-sucking Conorhinus, 414—The catalpa sphinx, 415—Notes on cotton worms, 415—Pyrethrum, 416.	
		416
2292. 1	RILEY, C. V. General truths in applied entomology. Ess: <trans. 153–159.="" 1884,="" agric.="" ga.="" pp.="" sep.<="" soc.,="" state="" td="" v.="" —,=""><td>•</td></trans.>	•
	rate: <macon, 153-159.<="" 1884,="" cover,="" ga.,="" pp.="" td="" tp.=""><td>Re-</td></macon,>	Re-
	print: <rept. 1884,="" 32<="" [u.="" agric.="" commissioner="" for="" pp.="" s.]="" td=""><td>23-</td></rept.>	23-

330. <Gardener's Chronicle, 1885, v. 23; 20 June, pp. 785-786; 27 June, p. 818. S.-b. No. 61, p. 6; 10-11. See: <Ga. Crop

- 2292. RILEY, C. V.—Continued.

 Rept., June, 1884, p. 33. <Sci. Amer. Suppl., 7 February, 1885, v. 19, p. 7588. S.-b. No. 61, p. 12.

 See No. 2291 for synopsis of contents.
- 2293. R[ILEY], C. V. Codlin moth. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2, p. 281, fig.
 Natural history, remedies, and figures of Carpocapsa pomonella.
- 2294. R[ILEY], C. V. Colorado potato-beetle. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2. p. 311, fig.

 Summary of the eastward spread of Doryphora 10-lineata; natural history,

enemies, remedies, and figures of the same.

- 2295. R[ILEY], C. V. Cotton-worm. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2, pp. 432–433, figs.

 History, ravages, life-history, remedies, and figures of Aletia xylina.
- 2296. R[ILEY], C. V. Curculio, plum. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2, p. 494.

 Natural history and means against Conotrachelus nenuphar.
- 2297. RILEY, C. V. Insect named. < Daily Gate City [Keokuk, Iowa], 18 February, 1885. S.-b. No. 63, p. 31.

 Brief notes of Olfersia americana.
- 2298. RILEY, C. V. A new remedy for the imported cabbage-worm. <Rural New-Yorker, 28 February, 1885, v. 44, p. 132. S.-b.</p>
 No. 63, p. 50. See: <Colman's Rural World, 5 March, 1885, v. 38, p. 76. S.-b. No. 63, p. 8. <Prairie Farmer, 7 March, 1885, v. 57, p. 153. S.-b. No. 61, p. 26. <Amer. Farmer, 1 April, 1885, v. 4, p. 105. S.-b. No. 67, p. 60. <Sci. Amer., 23 May, 1885, v. 52, p. 322. S.-b. No. 61, p. 19. <Indiana Farmer, 30 May, 1885. S.-b. No. 61, p. 22.</p>

Ice-cold water applied during the heat of the day causes the death of the larva of *Pieris rapæ*.

- 2299. RILEY, C. V. The collection of insects in the National Museum.

 <Science, 6 March, 1885, v. 5, pp. 188-189. S.-b. No. 61, pp. 7-8.

 Answer to criticisms of C. H. Fernald; the insects in the National Museum cared for by the entomologist of the Department of Agriculture.
- 2300. RILEY, C. V. Ants and aphides. <Rural New-Yorker, 14 March, 1885, v. 44, p. 171. S.-b. No. 61, p. 19.
 - Answer to inquiry of J. McFarland; habits of ants in relation to plant-lice; subterranean plant-lice attended by ants; ants in northern States beneficial rather than injurious; ground can be cleared of them by injection of bisulphide of carbon or a strong kerosene emulsion.
- 2302. RILEY, C. V. Jumping spiders. <Rural New-Yorker, 11 April, 1885, v. 44, p. 250. S.-b. No. 63, p. 33.

 Answer to inquiry of H. B. S.; habits of Attus sp.

- 2303. RILEY, C. V. Circular No. 9 [U. S. Department of Agriculture, Division of Entomology]. <[Washington:] 1 May, 1885, 1 p., fo:, 1 fig. Reprint: <Pacific Rural Press, 16 May, 1885. S.-b. No. 63, p. 27. <Weekly Times-Democrat [New Orleans], 23 May, 1885. S.-b. No. 61, p. 23. <Rural Californian, June, 1885, v. 80, p. 122. <S.-b. No. 61, pp. 20–21. <Sci. Amer. Suppl., 6 June, 1885, v. 19, p. 7859. S.-b. No. 63, pp. 25–26. <Pacific Rural Press, 16 May, 1885, v. 29, p. 469. S.-b. No. 63, p. 27. See: <Farmer's Review, 21 May, 1885, v. 14, p. 362. S.-b. No. 63, p. 29. N. Y. Weekly Sun, 27 May, 1885. S.-b. No. 63, p. 29. N. Engl. Farmer, 16 May, 1885. S.-b. No. 63, p. 29.
 - Conditions under which the Government will buy silk-worm eggs; how to select moths to produce eggs.
- 2304. RILEY, C. V. The imported elm-leaf beetle. Its habits and natural history and means of counteracting its injuries. <Bull. No. 6, Div. Ent. U. S. Dept. Agric. [14 May], 1885, 20 pp., 1 pl., 1 fig.
 - Treats of Galeruca xanthomelana; its habits and natural history; description, figures, remedies, and methods of application of insecticides.
- 2305. RILEY, C. V. Destroying Cicadas: Scurfy apple bark-louse. <Rural New-Yorker, 23 May, 1885, v. 44, p. 353. S.-b. No. 61, p. 24.
 - Answer to inquiries of J. A. K.; means against Cicada [= Tibicen] septendecim and Chionaspis furfurus.
- 2306. RILEY, C. V. [Poisonous spider.] <Rural New-Yorker, 23 May, 1885, v. 44, p. 354. S. b. No. 61, p. 23.
 - Reply to statements of Dr. W. H. G.; no spider known whose foot-tracks kill leaves or trees.
- 2307. [RILEY, C. V.] Expected advent of the locust. <Sci. Amer., 23 May, 1885, v. 52, p. 320. S.-b. No. 61, p. 9. <Farmer's Home Journ., 13 June, 1885. S.-b. No. 63, p. 18. <Orange Co. [N. Y.] Farmer, 28 May, 1885. S.-b. No. 63, p. 23. <See: <Sci. Amer., 20 June, 1885, v. 52, p. 389. S.-b. No. 63, p. 28. Simultaneous appearance of a 17-year (Cicada [= Tibicen] septendecim) and a
 - Simultaneous appearance of a 17-year (Cicada [= Tibicen] septendecim) and a 13-year (C. [= T.] tredecim) brood; localities of the two broods; life-history and habits.
- 2308. RILEY, C. V. Rust of orange. <Rural New-Yorker, 16 May, 1885, v. 44, p. 355, figs. 196, 197. S.-b. No. 61, pp. 18–19, figs. 196–197. Reprint: <Colman's Rural World, 18 June, 1885, v. 38, p. 195, 2 figs. S.-b. No. 61, pp. 31–32, 2 figs.
 - Description, habits, and means against *Phytoptus ęleivorus*; figures the same and its eggs; figure of a rusty orange.
- 2309. [RILEY, C. V.] Red-ants. <Sci. Amer., 21 May, 1885, v. 52, p. 183. S.-b. No. 66, p. 35.
 - Recommends pyrethrum, kerosene, hot water, and naphthaline as means against red-ants in houses.

- 2310. [RILEY, C. V.] Ridding the ground of cut-worms. <Rural New-Yorker, 30 May, 1885, v. 44, p. 368. S.-b. No. 63, p. 22.

 Answer to inquiry of V. R. R.; means against the larvæ of Agrotidæ.
- 2311. [RILEY, C. V.] An entomological breakfast. <N. Y. Times, 2 June, 1885. S.-b. No. 63, p. 1. See: <Belvidere [Ill.] Standard, 23 June, 1885. S.-b. No. 63, p. 7. Cicadas and Aerididæ as food.
- 2312. RILEY, C. V. The periodical or seventeen-year Cicada. < Harper's Weekly, 6 June, 1885, v. 29, p. 363, 4 figs. S.-b. No. 61, pp. 24-26, 4 figs.
 - Records and map showing distribution of brood XXII (septendecim) and of brood VII (tredecim); habits, enemies, and figures, with details of structure of Cicada [= Tibicen] septendecim; figure of twig punctured by the same and of twig healed after the puncture.
- 2313. [RILEY, C. V.] The winged pests of the West. <St. Louis
 Globe-Democrat, 9 June, 1885. S.-b. No. 61, p. 11.
 - Prediction that the northern States of the Mississippi Valley will escape serious damage from locusts this year; reasons for this opinion; ravages of Camnula pellucida in California.
- 2314. RILEY, C. V. The periodical or seventeen year Cicada. < Amer. Grange Bulletin, 11 June, 1885. S.-b. No. 63, pp. 16-17. Chronological record, natural history, and popular names of Cicada [= Tibicen] septendecim.
- 2315. RILEY, C. V. The periodical Cicada. An account of *Cicada* septendecim and its tredecim race, with a chronelogy of all broods known. <Bull. No. 8, Div. Ent. U. S. Dept. Agric., [17 June], 1885, 46 pp., 8 figs. Second edition, 13 July, 1885.
- 2316. RILEY, C. V. Notes on joint worms. <Rural New-Yorker, 20 June, 1885, v. 44, p. 418, figs. 215–218.

 Habits, means against, and figure of Isosoma hordei; criticism of A. J. Cook's
- views on Isosoma; figures I. tritici and I. grande.

 2317. RILEY, C. V. Beetles in the corn-fields. < Daily Gate City
 - [Keokuk, Iowa], 23 June, 1885. S.-b. No. 61, p. 17.

 Answer to letters of J. M. Evans and J. M. Schaffer; Agonoderus pallipes injurious to seed corn; recommends soaking the seed corn in Paris green or London purple.
- 2318. RILEY, C. V. Notes on the periodical Cicada. <Science, 26 June, 1885, v. 5, pp. 518-521. Reprint: <Sci. Amer. Suppl., 27 June, 1885, v.19, pp. 7905-7906. S.-b. No. 61, pp. 47, 48; No. 63, p. 74.
 - Record showing distribution of broad XXII (septendecim) and of broad VII (tredecim) in 1885; the specific value of the different forms; long period of underground development; its life-history; food of the larva, methods of burrowing, and transformations; the Cicada versus civilization.
- 2319. RILEY, C. V. The Chester onion pest. <Orange County [N. Y.] Farmer, 2 July, 1885. S.-b. No. 61, pp. 1–2. See: <Rural New Yorker, 5 December, 1885, v. 44, p. 829. S.-b. No. 61, p. 151.
 - Extract from Rept. U. S. Ent. for 1884; remedies against cabbage cut-worms; the poisoned ball system and kerosene emulsion effective against the onion cut-worm; formula for a soap-kerosene emulsion.

- 2320. RILEY, C. V. Premature appearance of the periodical Cicada. <Science, 3 July, 1885, v. 6, pp. 3-4.
 - Criticism of L. F. Ward's record of the occurrence and song of Cicada [= Tibicen] septendecim. See No. 2326.
- 2321. RILEY, C. V. Periodical Cicada in Massachusetts. <Science, 3 July, 1885, v. 6, p. 4.
 - The occurrence of Cicada [= Tibicen] septendecim in southeastern Massachusetts needs confirmation.
- 2322. RILEY, C. V. Destructive insects of the year. <Rural New Yorker, 11 July, 1885, v. 44, p. 464. S.-b. No. 61, p. 8.
 - Injuries and means against Agrotis malefida, Anthonomus musculus and Eury-creon rantalis; localities and food-plants of the last.
- 2323. [RILEY, C. V.] [Grasshopper ravages in California.] <Rural New-Yorker, 11 July, 1885, v. 44, p. 470. S.-b. No. 61, p. 2.

 Ravages committed by Melanoplus [= Caloptenus] devastator and Caloptenus differentialis; recommends the use of coal-oil pans against them.
- 2324. RILEY, C. V. Pests of the strawberry. <Rural New-Yorker, 18
 July, 1885, v. 44, p. 484. S.-b. No. 61, p. 17.
 - Answer to inquiry of J. H. J.; means against Agrotis tricosa, Phoxopteris fragariæ, Eccopsis permundana and Emphytus [=Harphiphorus] maculatus.
- 2325. RILEY, C. V. The imported elm-leaf beetle. <Harper's Weekly, 18 July, 1885, v. 29, p. 463, 1 fig. S.-b. No. 61, pp. 21-22, fig. Natural-history descriptions and figures of all stages of Galeruca xanthomelæna; means against the same.
- 2326. RILEY, C. V. Premature appearance of the periodical Cicada. <Sci. Amer. Suppl., 15 August, 1885, v. 20, p. 8021.
 - Criticism of L. F. Ward's record of the occurrence and song of *Cicada* [= *Tibicen*] *scptendecim*; an introductory note states that this article was prepared for Science, but that it appeared in that journal in a mutilated and weakened form. See No. 2320.
- 2327. RILEY, C. V. The cyclone nozzle. <Rural New-Yorker, 22 August, 1885, v. 44, p. 567. S.-b. No. 61, p. 34. Facts relative to the invention of the cyclone nozzle.
- 2328. RILEY, C. V. A new remedy against destructive locusts. < Rural New-Yorker, 29 August, 1885, v. 44, p. 577. S.-b. No. 61, p. 33.
 - Recommends the use of poisoned bait; formula of the same.
- 2329. RILEY, C. V. The influence of climate on *Cicada septendecim*. <Enton. Amer., August, 1885, v. 1, p. 91.
 - Records the transfers of eggs of brood XXII (septendecim) of Tibicen septendecim to the extreme southern States where no septendecim brood is known to occur, and of brood VII (tredecim) to northern States where no tredecim brood is known to occur.
- 2330. RILEY, C. V. Department of insects. <Ann. Rept. of Regents Smith. Inst. for 1883, 1885, pp. 239-244.
 - Brief mention of work accomplished in the U.S. National Museum; list of accessions to the collection.

- 2331. RILEY, C. V. Insects of the year. < Prairie Farmer, 3 September, 1885, v. 57, p. 567. S.-b. No. 63, p. 37; 134. See: < Entom. Amer., December, 1885, v. 1, pp. 176-177.
 - Ravages of Agrotis messoria, Anthonomus musculus, Eurycreon rantalis, Cecidomyia destructor, and Acrididæ on the Pacific coast.
- 2332. RILEY, C. V. On the parasites of the Hessian fly. <Proc. U. S. Nat. Mus., 14–17 September, 1885, v. 8, pp. 413–422, pl. 23. Abstract: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, pp. 332–334. <Amer. Nat., November, 1885, v. 19, pp. 1104–1105.
 - Description, synonymy, habits, and figures of Merisus destructor (Say); M. subapterus n. sp. [p. 416, pl. 23, f. 2: p. 333]; Eupelmus allynii French; Tetrastichus productus n. sp. [p. 419, pl. 23, f. 5: p. 333]; and Platygaster herrickii Packard.
- 2333. RILEY, C. V. Enemies of the black-walnut and willow. <Rural New-Yorker, 19 September, 1885, v. 44, p. 632. S.-b. No. 61, p. 35.
 - Habits, description of eggs, larva, and adult of Datana ministra; description of eggs, larva, and adult of Cimbex americana; means against both species.
- 2334. RILEY, C. V. The song-notes of the periodical Cicada. <Science, 25 September, 1885, v. 6, pp. 264–265. Reprint, with additions: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, pp. 330–332. Translation: <Stett. Ent. Zeit., 1886, jahrg. 47, pp. 158–160. See: <Science, 11 September, 1885, v. 6, p. 225. <Kansas City Review, October, 1885, p. 171.
 - Description of the three prevalent notes of Cicada [= Tibicen] septendecim.
- 2335. RILEY, C. V. The probabilities of locust or "grasshopper" injury in the near future, and a new method of counteracting their injury. <Colman's Rural World, 29 October, 1885, v. 38, p. 348. S.-b. No. 63, pp. 32-33. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, pp. 519-520. <Proc. 6th Meeting Soc. Prom. of Agric. Sci., Manhattan, Kans., 1886, pp. 38-39. See: <Mich. Christian Herald, 3 September, 1885. S.-b. No. 63, p. 39. <Amer. Grange Bull., 5 November, 1885. S.-b. No. 63, p. 34. <Prairie Farmer, 10 October, 1885, v. 57, p. 669. S.-b. No. 61, pp. 34-35.
 - The advance of settlement and cultivation will prevent such widespread injuries of *Caloptenus spretus* as occurred between 1874 and 1877; use of poisoned baits.
- 2336. RILEY, C. V. The present status and future prospects of silk-culture in the United States. <Entom. Amer., October, 1885, v. 1, pp. 139-140. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, p. 516. See: <Science, 11 September, 1885, v. 6, p. 236.
 - Abstract; headings under which the subject is discussed; summary of conclusions.

- 2337. RILEY, C. V. The grain moth. <Rural New-Yorker, 7 November, 1885, v. 44, p. 744. S.-b. No. 63, p. 56.
 Habits, descriptions, ravages, and means against Asopia farinalis.
- 2338. [RILEY, C. V.] Profits of silk-culture. <Rural New-Yorker, 26
 December, 1885, v. 44, p. 885. S.-b. No. 63, pp. 50-51.

 Answer to inquiry of L. M. S.: notuel cost of producing silk in different parts.

Answer to inquiry of J. M. S.; actual cost of producing silk in different parts of the United States.

- 2339. RILEY, C. V. Aletia xylina vs. A. argillacea. <Entom. Amer., December, 1885, v. 1, pp. 161–163.
 - Reasons for considering Aletia xylina the correct name for the cotton-worm; advance print from No. 2343.
- 2340. [RILEY, C. V.] [Euphanessa mendica, etc.] < Entom. Amer., December, 1885, v. 1, pp. 170–171, 173, 176.
 - Remarks on papers read at the A.A.A.S.; Euphanessa mendica considered a geometrid; some characters of the Lepidoptera; life-habits of Egerida.
- 2341. RILEY, C. V. On the hitherto unknown mode of oviposition in the *Carabidæ*. Abstract: <Proc. Amer. Assoc. Adv. Sci. for 1884, 1885, v. 33, pp. 538-539.
 - History of Chalanius impunctifrons traced from egg to imago; eggs laid singly on leaves of trees inclosed in mud or clay.
- 2342. RILEY, C. V. [Report of the Edinburgh International Forestry Exhibition.] < Rept. U. S. Dept. Agric. for 1884, 1885, pp. 167–179.

Brief mentiou of some of the insect collections exhibited.

2343. RILEY, C. V. Fourth report of the United States Entomological, Commission, being a revised edition of Bulletin No. 3, and the final report on the cotton-worm, together with a chapter on the boll-worm, with maps and illustrations. <Washington: 1885, [3 February, 1886], pp. 38+399+147, 45 figs., 64 pl., 2 maps.

1

5

CHAPTER II:

Characters, habits, and natural history

Distinct states of growth, 5—The worm must hatch from an egg,
5—Description of egg, 5—Where the egg is laid, 5—Number of
eggs to a leaf, 6—Time elapsing before hatching, 6—Unhatched
eggs perish when frozen, 6—Insects' and other eggs mistaken for
eggs of Aletia, 6—Characters of worm, 6—Description of newlyhatched worm, 6—Number of molts, 7—Different colors of worm,

CHAPTER II-Continued.

7-Habits of worm at different stages, 7-Jumping habit, 7-Only known to feed on cotton and one another, 8—Odor of the worm, 8-Migrations of worm, 8-Method of pupation, 8-Formation of cocoon, 8-The worm never burrows in the ground, 8-Distinctive characters of the chrysalis, 9-Duration of chrysalis state, 9-Distinctive characters of the moth, 9-Sexual differences in the moths, 9-Different habits of the moths at night and by day, 10-Their strong flight, 10-Their position at rest, 10-How soon the female begins to lay, 10-Her prolificacy, 10-Food of the moth, 10-It injures fruits, 11-Structure of the tongue, 11-Mention of other Noctuida which injure fruits, 11-Time elapsing from one generation to another, 11-This will average about one mouth, 12-Time of year when the first worms appear, 12-Former opinions erroneous, 12-Dates of earliest appearance variable, 12-Worms of all sizes found in Florida and southern Georgia in the latter part of March, 12-First worms always few and scattered, 12-They multiply and spread irregularly, 12-Their progress governed by the season and latitude, 13-The third generation often called the first, 13-Number of annual generations, 13-There are at least seven in southern Texas, 13-Gradual progress and succession of broods, 14—The second generation exceptionally very injurious, 14— Why not usually so, 14-Extent of migratory flights of the moths, 15-Probable existence of northern food-plants, 15-Causes and seasons of migration, 15-Behavior of migrating moths, 15-The question of hibernation, 15-Different former opinions and beliefs concerning hibernation, 16-Discussion of these opinions, 16—The chrysalides are killed by a temperature below 22° F., 16—Parasited chrysalides can bear greater cold, 16-Fallacy of the belief that the chrysalis winters underground, 16-Ease with which erroneous conclusions can be drawn from mistaken identity, 17-The chrysalis of Aspila virescens mistaken for that of the Aletia, 17-Chrysalides of other cotton larvæ found underground in abundance, 17-The chrysalis of Aletia killed by burial, 17-Ability of the moth to survive the winter, 18-Unreliability of most testimony as to the hibernation of the moth, 18-Other moths easily mistaken for it, 18-Hypena scabralis, 18-Phoberia atomaris, 19-Leucania unipuncta, 19—Absence of testimony to the survival of the moths beyond March, 19-Theory of annual introduction of the species from some southern foreign country, 19-Statement and discussion of Grote's arguments and of others in support of this theory, 20-Arguments in favor of the hibernation of the moth, 21-Both immigration and hibernation may occur, 21-Summary of the evidence, 22-Aletia hibernates only as a moth and only in the extreme south, especially in Texas, 22.

CHAPTER III:

338	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	
2343.	RILEY, C. V.—Continued. CHAPTER III—Continued. remedies, 35, 36—Arsenic, 36—Paris green, 36-38—Planting jute, 38—Machinery, 38—London burple, 38—Pyrethrum, 38. CHAPTER IV: The cotton-worm in other countries. Geographical distribution of Aletia xylina, 39—Other insects injurious to cotton in the Eastern Hemisphere, 39—Occurrence of Aletia xylina on the Pacific coast of Mexico, 39-40—Ou the Gulf coast of Mexico, 40-41—In Yucatan, 41—In the West Indies, 42—In the northern countries of South America, 42-43—In Brazil, 43-44.	3
	Chapter V: On the anatomy of Aletio. By Edward Burgess and C. S. Minot Circumstauces under which this chapter was prepared, 45—External anatomy of larva, 45—True lcgs, 45—Prolegs, 46—Colored markings, 46—Stigmata, 46—Internal anatomy of larva, 47—Head, 47—Ganglia, 47—Digestive canal, 47—Malpighian vessels, 47—Salivary glands, 47—Dorsal vessel, 47—Stomach, 48—External anatomy of imago, 48—Method of preparing the exoskeleton for examination, 48—Head and appendages, 48—Prothorax, 48, 49—Mesothorax, 49—Metathorax, 49, 50—Supposed organ of hearing, 50—Abdomen, 50–51—Spiracles, 51—Lcgs, 51—Scales, 51–52—Scale-pores, 52—Proboscis, 52—Spines, 53—Internal anatomy of imago, 53—Digestive canal, 53—Pharynx, 53–54—Salivary glands, 54—Œsophagus, 54—Food reservoir, 54–55—Stomach, 55—Malpighian vessels, 55—Intestines, 55—Aorta, 55—Nervous system, 55—Ganglia, 55—Terminal body segments and organs of reproduction, 56—Brush-sac, 56—Male organs of reproduction, 56—Testes, 57—Vasa deferentia, 57—Penis, 57—Female organs of reproduction, 57—Ovaries, 57—Sebaceous glands, 57—Vagina, 58—Copulatory pouch, 58—Terminal abdominal segments of the female, 58.	4
	CHAPTER VI: The cotton belt. By Prof. E. A. Smith A. General features of the cotton States Region included in the cotton belt, 59—Climate, 59—Winds, 59— Rainfall, 59-61—Temperature, 61-62—Geological sketch, 62-63— Topography, 63-64—Soils, 64-65—Agricultural subdivisions, 65— Regions of forest growth, 66-67.	5
	B. Description of the agricultural-subdivisions. The alluvial region, 67–68—The lower prairie region, 68–69—The long-leaf pine region, 69–71—The oak uplands region, 72–73—The upper prairie region, 73–75—The red and brown loam region, 75–	ϵ

The gneissic region, 79-80. Chapter VII:

Terrestrial and meteorological influences affecting the worm.

Condition of soil and plant connected with the appearance of the first worms, 81—The earliest worms are confined to the "low lands," 81; and to luxuriant plants, 82; and to the vicinity of winter shelters, 82; and to the same localities, 82—Influence of wet weather on the development of the worms, 83—Severe rains with gales destroy both worms and crop, 83—Late cold rains do the same, 84—Frequent summer rains favor the development of the worms, 84—Hot, dry weather destructive to them, 84—Artifis

77—The sandy and siliceous lands of the older formations, 77–79-

CHAPTER VII-Continued.

cial drought produces the same effect, 84—Indirect influences of wet weather in favoring the development of the worms, 85—İmmunity of the worms from their enemies in wet weather, 85—Drowning of ants by heavy showers, 85—Wet weather prevents poisoning and working the cotton, 85—Letter from J. W. Du Bose on the influence of winds on Aletia, 85–86.

CHAPTER VIII:

Natural cnemies General remarks, 87—Importance lessened by the use of arsenical poisons, 87—Vertebrate enemies. 87—Quadrupeds, 87—Birds, 87— English sparrow, 88—Toads and lizards, 89—Invertebrates, 89— Spiders, 89-Ants, 90-Species of ants destroying Alctia, 90-Hubbard's observations on ants, 92—The leaf-cutting ant, 94— Wasps, 94—Coleoptera, 95—Tiger-beetles, 95—Ground-beetles, 95—Lady-birds, 96—Soldier-beetles, 96—Heteroptera, 97—List of species, 97—Diptera, 99—Orthoptera, 99—Neuroptera, 100—Lepidoptera, 100-Parasites, 101-List of species, 101-The Trichogramma egg-parasite, 102—The cotton-worm Microgaster, 104— Comstock's Euplectrus, 105—Elachistus euplectri, a secondary parasite, 106—The common flesh-fly, 107—Cyrtoneura stabulans, 108—Is it parasitic?, 108—Tachina-flies, 109—Their habits, 109— The watchful Pimpla, 111—The ring-legged Pimpla, 113—Cryptus nuncius, 113-The ovate Chalcis, 114-The devouring Tetrastichus, 115-Species that are easily mistaken for parasites of Aletia, 115—Hexaplasta zigzag, 115—Phora aletia, 116.

CHAPTER IX:

Preventive measures

Mode of cultivation, 120—Improving cotton seed, 120—Forcing the young plants, 120—Transplanting from hot-beds, 120—Objections, 121—Frequent cultivation, 121—Topping the cotton, 121—Fertilizers, 121—Sulphuric acid on seeds, 121—Late planting, 121—"Worm-proof cotton," 121—Diversified agriculture, 122—Rotation of crops, 122—Jute as a protection, 122—Other supposed protective plants, 123—Protection of natural enemies, 123—Immunity of cotton under trees, 124—Preventing oviposition of the moth, 124—Futility of decoctions for this purpose, 125—Road dust, 125—Early application of direct remedies, 126—Concerted action, 126—Early poisoning and hand-picking, 126—Cotton-worm warnings, 127—Destroying chrysalides accidentally carried into gin-houses, 127—False theories, 127—Burning the stalks, 127—Winter plowing, 127—Salt, 127.

CHAPTER X:

Remedies; means of coping with the insect; substances used for its destruction

Destruction of the eggs, chrysalides, and moths, 128—Destruction of the eggs impracticable, 128—Little chance for successful destruction of the chrysalides, 128—Destruction of the moth, 129—Lights and fires for attracting the moths, 129—Indifferent success with lamps used at Columbus, Tex., 130—Importance of using lamps carly in the season, 130—Apparent success with lamps near Hearne, Tex., 131—Great attractiveness of the electric light, 131—Movable lights, 131—Poisoned sweets and fluids, 131—Fondness of the moths for ripe fruit, 132—Killing moths by poisoned fruit,

87

120

CHAPTER X-Continued.

132—Method of using poisoned liquids, 132—No results from using poisoned baits late in summer, 133-Poisoning the glands of the plant, 133—Cotton-leaf essence and its attraction to the moth, 134—Hand-picking, 135—Mechanical means of killing the worms, 136—Shaking off the worms, 136—Poisoning the worms, 136— Progress in the use of insecticides, 137—Classification of insecticides, 137—Importance of preparing materials in advance, 138— Arsenical compounds, 138—Safety in their use, 138—Difficulty in determining minimum quantities, 139—Overdoses of poisons, 139 -General rules in dry applications, 140-Mixing devices, 141-Ingredients, 141—Wet application, 142—Principles to be followed and ingredients to be used, 142-Comparison of dry and wet applications, 142—Paris green, 143—Historical data, 143—Liquid application, 143-Dry application, 144-Minimum quantities, 144 -Patents on Paris-green combinations, 146-Arsenic, 147-Commercial arsenic, 147—Arseniate of soda, 147—Fowler's solution, 148-Johnson's dead-shot, 148-Texas cotton-worm destroyer, 149 -London purple, 149-Manufacture and analysis, 149-History of its use, 150—Experience in Alabama in 1880, 150—Advantages and disadvantages, 151-Dry application, 151-Wet application, 152—Other mineral substances, 153—Salt and saltpeter, 153—Snlphur, 154—Red lead, 154—Road dust, 154—Oils and allied substances, 155—Kerosene, 155—Former methods of application, 155 -Invention and perfection of emulsions, 156-Method of preparing emulsions, 157—Formula for perfected emulsion, 158—Experiments with imperfect emplsion in 1880, 158-Experiments with perfected emulsions, 160—Oil of creosote, 162—Oil of tar, 162— Gas-tar water, 162—Carbolic acid, 163—Cotton-seed oil, 163— Vegetable insecticides, 164—Pyrethrum, 164—History, 164—Mode of cultivation, 165—Success in cultivating the plant in America, 166—Preparation of the powder, 167—Its use as an insecticide, 168—Advantages and disadvantages, 168—Active principle in Pyrethrum, 169—Its effects on cotton-worms, 169—Imported vs. Californian powder, 170—Dry application, 170—Minimum quantities of dry powder, 171-Dry powder mixed with flour and other ingredients, 172-Application in fumes, 174-Alcoholic extract, 174—Experiments with extract obtained by distillation, 174— Experience with extract obtained by repercolation, 176—Use of the powder in simple water solution, 177-Tea or decoction of Pyrethrum, 178-Effect of Pyrethrum upon other insects, 179-Prospects for the use of Pyrethrum for the cotton-worm, 180-Ox-eye daisy powder and its nselessness as an insecticide, 180— Extracts and decoctions from various plants, 181-Difficulties in the way of discovering new vegetable insecticides, 182-Mode of preparing the extracts and diffusions, 183-List of the plants experimented with, 184-Effect of alcohol upon the worms, 187-Yeast ferment and fungns infection, 188-Dr. Hagen's recommendations of the use of yeast as an insecticide, 188-Objections to Dr. Hagen's plan, 189-Negative results obtained by the Commission, 190.

CHAPTER XI:

Machinery and devices for the destruction of the worm. By W. S. Barnard, Ph. D

CHAPTER XI-Continued.

- I. Spray nozzles, classification, preferred kinds, 191-Many-punctured nozzles, 191-196—Preferred construction, 191-193—Straining and cleaning vs. clogging, 192-Eddy-roses, their operation, construction, and leading importance, 192-193-Plug-roses, action and construction unsatisfactory, 193-194-Johnson's, Melcher's, Dawson's, Foster's, 193-Lynch's, 194-Colliding jets, gas-jets, superiority, Daughtrey's, Weber's, Prouty's, 194-T-roses, 194-Yeager's, Warner's, improvements, 195-Divided rose-heads, clutch-head of Mast, Foos & Co., Foss', Fox's, Barrows', Vose's, 195-Peripheral roscs [divided], Mclcher's, Yeager's, Ruhmann's, 196-Rose-combinations of Barry, Prentice, etc., 196-Slot nozzles, 196-205—Operation, 196—Preferred construction, 196-201—Disadvantages, improvements made, 197-201-Eddy-chambered, 197-199—Lip construction, 198-199—Inside cleaner, 199—Simple slotnozzles, Fowler's, Mallory's, Iske's, 201-202-Plug slot-nozzles, Allen's, Ruhmann's, Johnson's, "The Niagara," Pinter's, 202-Removable slots of Long, Vestal, and Merigot, 203-Jawed slots, "The Boss Nozzle," Raymond's and Perkin's, Smith's, Moffet's, 203-Williams', Ruhmann's, Pinter's, 204-Stanton's, 205-Side slots, Schier's, Melcher's, 205—Deflector nozzles, 206-211—Definition, use, applicability, relative merits, 206-Drip-waste, clogging, 206-Simple constructions made, 206-208-Conformations for narrow, wide, and even sprays, 206-207-Removable deflectors, 207-Hollings', Douglas's, Nickerson's, 208-Hayden's, Killam's, Lewis', Schier's, Barrett's, 209-Ruhmann's, Binkley's, Schier's compound, 210—Schier's and Polansky's, 211—Centrifugal nozzles, 211-221-Operation, kinds, choice, 211-The new and most perfect sprayers, 212-Eddy-chambered, 212-219-Conformation and operation, 212-Clogging, cleaning, construction principles, 213—Forms described, 214—Whistle-jets, 215-216—For blast-atomizers, 216—Eddy-jets proper, 216-219—Involute form, cone form, 216-Convex and concave forms, 217-Direct discharge, proximal, diagonal discharge, 217—Distal diagonal discharge, 218-Centrifugal uose-pieces, double-cone or doublechambered form, and chambered plug form, 218-Direct spray and solid jet, 219-Fistular spray nozzles, hose-pipes, 219-With rotary segment, Hotz's, Clifford's, Gielow's, Johnson's, Hoyer's, Clarke's, 220—With cross-plug, McGaffey's, Johnson's, Gray's, Gielow's, Hosford's, 221—Spray-wheels, 221.
- II. Centrifugal throwers, 221–226—Their character and operation, 221—Rotated orifices, pronged reels, brush poison-throwers, 222—Brush fibers, 222–223—Rotary polishing brushes, improved construction, fiber strength, spring and density, 223—Feeding the brushes with liquid, 223–225—With powder, 224—A simple brush thrower of powder, 224–225—Advantages over-sifters, velocity of rotation, 225—Wisewell's, 225–226—Whisps and brooms, rotated recesses, 226.
- III. Blowers of poison, 226-252—Rotary blowers, 226-235—Kinds and success of, 226-227—For blowing powder, 227-232—Hoppers and adjustable feeders, 227-228—Blower encasements and blast-pipes, 228-229—Blast forks and deflectors, 229—Improved light rotary blowers, 229-230—Hauled blowers, compound rotary blower, Aframe and legged swivel wheels, 230—Rotary velocity, 231—

CHAPTER XI—Continued.

Hurd's rotary powder blower, 231-232—For blowing liquids, 232-235—Improved feeders of liquid to rotary blower, 232-233—Dripcatcher, forked blast spray, 233—Darnell's rotary liquid-blower, 233-234—Hurd's ditto, and Perl's rotary fume-blower, 234— Force-blast rotary blowers—Oscillating blowers, 235-251—Bellows power, construction, durability, improvements, 235-236-For blowing powders, 236-243-Hermetic powder-box, 236-Feeders of powder to blasts, 236-237-Pipes, forks, and nozzles for these powder blasts, 237-238—Hauled compound bellows powderer, to wagon, with motor, etc., 238-Cultivator bellows-blower, 238-239-Knapsack and horse-back bellows powderers, 239-Improved small hand-bellows powderers, 239-241-With forks, 241—Allen's powderer, 241—Common powder bellows, Woodason's, Hendley's improved, 242—Stelle's, etc., 243—For blowing fluids, 243-249—Importance, resuction, feeding by blast-suction, blast-pressure, gravitation, gauges, 243-244—Blast spray conductors and nozzles, reverberatory, 244—Whistle-jcts and agitation chambers, 245—Common blast atomizers, 245—Improved automatic-feeding blast sprayers, 246-249-Reatomizing, reverberatory nozzles, 247-Nether blast sprays, directing pipes and nozzles, 248—Compound combinations, 249-Peck's blast sprayer, and Wallace's, 249—Reciprocating or pistoned blowers, 249-251— Improved air-pump apparatus, 249-250—Air-pumps of Humphryville, Rumsey, ctc., 251—Generator blowers, 251-252—Steinmann's vaporizer, 251.

CHAPTER XII:

Machinery and devices for the destruction of the worm—continued.

IV. Pneumatic compression squirters, 253-261—Kinds, advantages, available fire-extinguishers, 253-254—Carbonic anhydride, apparatus, practicability, 255—Danger, antidotes, safety constructions, 255-256—Simple generators, 257—Rotary force-blast, compression ejectors, 257—Oscillating bellows, pneumatic compression ejectors, 257—Reciprocating or pistoned ditto, 258-261—Simple kind to construct, beer forcers applicable, Worswick Co.'s, Weindel's, Rumsey's, Douglas's, 258—Daughtrey's underspray theory and machine, 258-261—The author's devices, 259—Weber's nozzles, 260.

V. Solid compression squirters of poison, 261-283-Kinds, 261-Rotary force-pumps, 261-262—Hydraulic bellows, 262—Oscillating force-pumps, 262—Reciprocating force-pumps, 262-283—Kinds, 262-Hydronettes and fountain pumps, 262-269—May's patent, Tyler's, Servants', Staples', Whitman's fountain pump, 263—Hydronettes, double-acting, Deakin's, Rumsey's, cost, 264—Author's improvements in knapsack apparatus, 264-265-In knapsack fireextinguishers, Condict's and Doty's, 265-In horseback apparatus, Warner's, author's, 265-In cart or wagon, 265-268-Calahan's carts, legged wheels, tongue or shafts, 266-Wagon use, Trelease's report, 266-268—More economic devices, 268—Other pumps not in trade, 268-269—Aquapults, aquajects, aquarius, excelsior, hydropult, 269-271-Operation, cost, 269-Aquapult, Johnson's patents, Douglas', Prouty's, 269-270-Johnson's syringe, Douglas' aquarius, Rumsey's aquaject, 270 — Deakin's excelsior, Vose's hydropult, 271-Bucket pumps and knapsack

CHAPTER XII-Continued.

pumps, 271-274—Lewis's, 271—Korth's and less desirable kinds, Stoner's, Mallory's Kaiser's, Dix's, 272-Crandal's, Hollaud's knapsack extinguishers, Douglas's, Stanton's, 273—Bucket poisoners, McDonald's, Allen's watering-pot improvements, 273-Amor's and Lane's can-syringe, Hull's, Wisner's, 274—Barrel and tank pumps and appurtenances, 274-283-Single acting, discharging below the piston, Melcher's, 274-Voglesang's, Ruhmann's, 275-Polansky's, Schier's, Butman's, Ball's agitator, 276-Evenden's, Helmecke's, 277-Yeager's pump and nozzle, 277-278-Pinter's, 278—Single-acting, discharging from above the piston, Chipley's, Weith's, 278—Available cistern or well pumps, 278-279—Barrow pumps, truck pumps, garden engine pumps, windmill pumps, Blunt's Lotus pump, the Pendulum and Index pumps, counter pumps, 279—Double-acting, force-pumps proper, 279-283—The best, 279—Kinds characterized, Vose's hydropult, the Champion, 280-Ramsdeu's, reduction-cylinder pumps, patentees and manufacturers, 281-282—Author's agitator barrel pump, found most satisfactory, 282-283—Conduits, frames, portage, and combinations of appurtenances, 283—Johnson's cotton-spraying machine, 284— Jones', Binkley's, '285—Goodiu's, Wolfram's, 286—Author's uuderspraying accessories, 288-Plan, skid, mixing, and straining funnel, 288—Adjustments of pipes, forks and nozzles, 289-297; fork modifications, 290-292—Pendent pipes, 292, conformability, lightness, cheapness, 292-293—Tube substances, 293—Author's Aframe machines, 293-297—Maximum width undersprayed, 295— Pipe adjustments to row widths, flexible systems superior, 296-297.

VI. Gravitational distributors, 297-309—For liquid, 297-302—Kinds, author's tripod automatic sprinkler, 297-298—Schank's sprinkler, Taylor's, 298—Robinson's, 299—Suction force-pumps and windlass elevators, 299—Bilge-pumps, 300—Horseback automatic sprinklers, watering-pot method, Willie's sprinkler, 300—Ramsey's, 301—Knapsack automatic sprinklers, 301—Gray's, Ruggle's, Townsend's, 302—Automatic hand-sprinklers, watering-pots afoot and on horseback, 302—For dry poisons, sifters, kinds, 302–309—Disadvantages, reciprocating sieve machines, hand-sieves, 303–304—Sifting bag, Hurd's sifter and blower, Goodheart's duster and sprinkler, 304—Rotary-sieve machines, 304–307—Robinson's duster and sprinkler, Davis' duster, Levy's, 305—Taylor's duster and sprinkler, 306—Reciprocating-stirrer sifters, Willie's, 307—Rotary-stirrer sifters, Young's, 307—Smith's, 308—Eldridge's, 309.

CHAPTER XIII:

Machinery and devices for the destruction of the worm—concluded.

VII. Insect manipulators, mechanical treatment, dislodging, crushing or stifling the worms or chrysalids, 310—Sweeping, knocking, or jarring off the worms; friction drags, fringes; beaters, 310—Collecting and despatching means, 311—Crushing, Helm's sweeper and crusher, 311—Ewing's sweeper and stifler, 312—moths, Wood-Smith's, 313—Iske's catching trays, 314—Traps for the 314-321—Kinds of lures, lighttraps, 314—Lewis's, McQueen's, Rigel's, 315—Walker's and others, 315-316—Lamps in motion, Leblanc's, Fordtran's, 316—Bait traps, Heard's, Garrett's, 317—Traps combining light and bait, author's net trap, 317-319, thought

CHAPTER XIII—Continued.

the best, 318—Indiscriminate killing wrong, the best bait, 319—Stith's trap, 319—Pugh's, Garrett's, 320—Binkley's, 321.

322

345

355

CHAPTER XIV:

History of the literature and bibliography...
History of the literature, 322—From 1802 to 1828, 322—From 1829 to 1847, 323—From 1848 to 1854, 324—From 1855 to 1871, 325—From 1869 to 1874, 326—From 1874 to 1878, 327—From 1878 to 1880, 328—From 1880 to 1881, 329—Bibliography up to and including the year 1881, 329–344.

CHAPTER XV:

Insects liable to be mistaken for Aletia..... Confusion of Aletia with other moths, 345—With Aspila virescens, 345—Characters of the most important of these moths to be illustrated in this chapter, 345-List of these moths, 345-Account of Anomis erosa Hübner, 345—Its geographical distribution, 346— Structure of its eggs, 346—Distinguished from egg of Aletia xylina, 346—Its seasons, 346—Habits of larva, 346—Characters of moth, 347—Detailed description of egg, 348—Larva, 348—Pupa, 349— Scasons and food-plants, 349—Account of Anomis texana n. sp., 350—Habitat of A. exacta, 350—Of A. texana, 350—Larva distinguished from that of Alctia xylina, 350—Pupa distinguished, 350—Account of Leucania unipuncta Haworth, 350—Reference to discussions about this species, 350—Its geographical distribution,. 351—Oviposition, 351—Food-habits of worms, 351—Pupation, 351 -Number of broods, 351-Hibernation, 351-Account of Aspila virescens, 351—Synonymy, 351—Confounded with Aletia xylina only in pupa state, 351—Pupæ of the two species distinguished, 352-Food-plants of A. virescens, 352-Moth distinguished from that of Aletia xylina, 352—Account of Drasteria erechtea (Cramer), 352—Its geographical distribution, 352—Variations in size of moth, 352-Food-plants of larva, 352-Habits of moth, 352-Number of broods, 352—Oviposition, 353—Colors of larva, 353—Account of Laphygma frugiperda, 353—Its food-habits, 353—Synonyms, 353—Account of Platyhypena scabra (Fabr.), 354—Geographical distribution, 354—Food-plants, 354—Number of broods, 354 -Reference to descriptions of larva, 354-Euplectrus platyhypenæ Howard, bred from larva, 354—Account of Phoberia atomaris (Hübner), 354—Reference to descriptions and figures of moth, 354-Food-plant of moth, 354.

CHAPTER XVI:

The boll-worm (Heliothis armigera Hübner)

Introductory, 355—One of the foremost of our injurious insects, 355—Extent of its ravages, 355—Nomenclature, 357—Synonyms, 358—Popular names, 358—Geographical distribution, 358—Foodplants, other than cotton, 359—Corn, 359—Tomato, 361—Tobacco and other Solanacew, 362—Leguminosw, 362—Cucurbitacew, 363—Malvacew, 363—Other food-plants, 363—Characters and transformations, 364—The egg, 364—The larva, 365—The pupa, 370—The imago, 371—Number of broods, 372—Hibernation, 373—Summary of the distinguishing points compared with Aletia, 374—Egg, 374—Larva, 374—Pupa, 374—Adult, 374—Natural enemies, 375—Remedies, 377—Early planting, 378—Low corn vs. high corn, 378—Fall plowing, 378—Destruction of the moths, 379—Lights

0242	Print C V Continued	
4040.	RILEY, C. V.—Continued. CHAPTER XVI—Continued.	
	and poisoned sweets, 379—Hand-picking, 380—Poisoning, 381—	
	Pyrethrum, 381—Bibliography, 382.	
	EXPLANATION TO PLATES	385
	Plate I-III, 385—IV-VII, 386—VIII-XI, 387—XII-XVI, 388—XVII-	
	XX, 389—XXI-XXV, 390—XXVI-XXVIII, 391—XXIX-XXXII,	
	392—XXXIII-XXXVIII,393—XXXIX-XLII,394—XLIII-XLVII,	
	395—XLVIII-LIII, 396—LIV-LVII, 397—LVIII-LXI, 398—	
	LXII-LXIV, 399.	
	APPENDICES	[1]
	Preface to appendices	[3]
	Report of H. G. Hubbard	[5]
	Report of observations made in 1881 upon Aletia and other insect	[o]
	enemies of cotton, in the State of Florida, [5]—Cotton injured	
	more by Dysdercus suturellus than by Aletia xylina, [6]—And	
	more by rust-mite than by either, [7]—Great destruction of	
	Aletia eggs by Trichogramma pretiosa, [7]—Successions of broods	
	of Aletia, [7]—Distribution of worms on plant, [8]—Tabular	
	statement of periods of time occupied in each stage of growth,	
	[10]—Locality of deposition of eggs on plant [12]—Proportions.	
	of light and dark colored worms, [12]—Effect of shade in protect-	
	ing cotton, [12]—Periods of different stages, [13]—Copulation, [13]—Position of moths at rest, [13]—Experiments with poisons,	
	[14]—Pyrethrum, [14]—London purple, [15]—Poisoning the	
	moths, [15]—Yeast, [16]—The boll-worm, Heliothis armigera	
	Hübn., [16].	
	APPENDIX II:	
	Report of Prof. R. W. Jones	[17]
	The cotton boll-worm, Heliothis armigera, [17]—Its importance, [17]	
	-Food-plants, [17]-Natural history, [18]-The egg, [18]-The	
	larva, [18]—Chrysalis, [19]—The moth, [19]—Weather, [19]—	
	Natural enemies, [19]—Cotton army-worm, Aletia xylina, [19]—First appearance observed in 1880, [19]—Experiments with	
	poisoned sweets, [20]—With adhesive sweets, [20]—Gradual dis-	
	appearance of moths in October, [20]—Preparation of vegetable	
	substances as insecticides, [20]—Pyrethrum, [21]—Experiments	
	with Pyrethrum, [22].	
	APPENDIX III:	
	Report of J. P. Stelle	[25]
	First appearance of Aletia xylina and Heliothis armigera in Texas	
	in 1880, [25]—Cotton blight, [25]—Its symptoms, [25]—Its	
	causes, [26]—Boll rot, [26]—Its symptoms, [27]—Its cause [27] —The flare, [27]—Its symptoms, [27]—Its probable cause, [27]	
	-Other food-plants than cotton for Aletia, [27]—None found for	
	the larva, [27]—Many for the moth, [27]—Food of larva of	
	Heliothis armigera, [28]—Annoyances to the cotton-worm, [28]—	
	Common salt, [28]—Saltpeter, [28]—Road dust, [28]—Open	
	spaces, [28]—Trees, [29]—Shade, [29]—Natural enemics, [29]—	
	Birds, [29]—Ants, [29]—Other insects, [30]—The yeast ferment	
	remedy, [31]—Pyrethrum powder, [31]—Other vegetable poisons,	
	[32]—Arsenical poisons, [32]—London purple, [32]—Paris green,	
	[33]—Arsenic, [33].	

[37]

[49]

[55]

[59]

_	_			_	7	_	Ť		•	_	_	 Ī	_
	A	PI	E	N.	DI	X		IV	:				

Reports of Dr. E. H. Anderson For 1880, [37]—Seasons and habits of A. xylina, [37]—Of Heliothis armigera, [37]—Localities of first appearance of A. xylina in the season, [37]—Reasons for this first appearance, [37]-[38]—Hibernation of this insect in some form, [38]-Causes of undue multiplication of the worms, [38]—Inflnence of ants on the number of worms, [38]—Habits of ants, [38]—Other insects gathering sweets from the cotton-plant, [39]—Jnte growing amidst cotton has no effect on the worms, [39]—Saturnia io found eating cotton, [39]—An enemy of the cotton-worm found, [39]—Experiments with London purple, [39]—With Pyrethrum extract, [39] —With yeast, [40]—Eggs of insects found on cotton-leaf, [40]— Characteristics of season of 1880, [40]—Report for 1881, [40]— Diary of observations on issuance, egg-laying, and hibernation of moths, and on weather, from October 3 to December 31, [40]-[44]—Observations of Heliothis armigera, [41]-[43]—Report for 1882, [44]—Observations on ants and aphides, [45]—Abundance and sudden disappearance of Laphygma frugiperda, [45]—Scarcity of cotton-worms, [45]—Experiments with Pyrethrum, [45]— [48]—Weather in September, [46]-[47]—Scarcity of Heliothis armigera in cotton, [46]-[47]—Its preferred food-plants, [47].

APPENDIX V:

APPENDIX VI:

Report of Judge William J. Jones.

Obstacles to investigation in 1880, [55]—Exemption of scattered patches of cotton-plants from depredation, [55]—Answers to questions in Circular No. 7, [55]—[56]—Cotton killed by a fungus, [56]—Application of poisons to cotton, [56]—Effect of poisons upon eggs and larvæ, [56]—No other insects observed to be injurious, [56]—Use of lights recommended to destroy moths, [56]—[57].

APPENDIX VII:

Reports of consuls and consular agents on the cotton crop and its enemies in Mexico, Central and Sonth America, and the West Indies....

Introduction, [59]—Circular letter of inquiry, [59]—Replies: from Merida, Mexico, [60]—From Tampico, Mexico, [60]—From Martinique, W. I., [60]—From Trinidad, W. I., [61]—From Manzanillo, Mexico, [61]—From Mazatlan, Mexico, [62]—From Bahia, Brazil, [63]—From Maricaibo, U. S. of Colombia, [64]—From Vera Cruz, Mexico, [65]—From Pernambuco, Brazil, [67]—From Bogotá, U. S. of Colombia, [68]—Notes on insects

2343.	RILEY, C. V.—Continued.	
	APPENDIX VII—Continued.	
	injurious to the cotton-plant in the Republic of Mexico, by D.	
,	H. Strother, U. S. consul-general, [70].	
	APPENDIX VIII:	
	Answers to Circular No. 7	[71]
ĕ	From D. M. Hamilton, St. Francisville, West Feliciana Parish, La., [71]—From R. A. Lee, Evergreen, Ala., [73]—From J. M. Wolkom, Henderson, Tex., [74]—From P. S. Clarke, Hempstead, Waller Co., Tex., [75]—From L. D. Hoyt, Livingston, Sumter Co., Ala., [77]—From F. S. Shields, Lake Concordia, Concordia Parish, La., [79]—From G. E. Gillespie, M. D., Natchitoches, La., [81]—From H. O. Dixon, Jackson, Miss., [84]—From F. L. Yoakum, Larissa, Cherokee Co., Tex., [85]—From O. H. Perry,	
	Perry Co., Ala., [86]—From J. W. Grace, Walterborough, Colleton Co., S. C., [86]—From F. M. McMeekin, Jamcstown, Alachua Co., Fla., [87]—From H. P. Bee, San Antonio, Tex., [88]–[89]—From E. H. Anderson, Kirkwood, Miss., [90]—Condensed summary of the habits of the worm, from Dr. D. L. Pharcs, Woodville, Miss., [92].	
	Notes	[93]
	Note 1	[95]
	Hübner's description of Aletia argillacea, with translatiou, [95].	
	Note 2	[95]
	Reasons for rejecting Hübner's description of Aletia argillacea, [95] —Wherein this description and the figures differ from A. xylina, [95]—Doubts of authors concerning A. argillacea, [95]—Search for the type of A. argillacea, [96]—Condition of the Sommer collec- tion, [96]—How specimens of A. xylina are labeled therein, [96].	F063
	Note 3. No published, full, and complete description of the earlier states of A. xylina extant, [96]—Reprint of descriptive portion of Dr. C. W. Capers' article "On the cotton caterpillar," [97]—History of ravages of A. xylina prior to 1828, [97]—Sudden abandonment of cotton-plants, [97]—His description of moth, [98]—Egg, [98]—Larva, [98]—Smell of larva, [98]—Habits, [98]—Food-plant, [98]—Pupation, [98]—Description of pupa, [98]—Remedies, [99]—Detailed description of egg, [99]—Six stages of larva, [99]—Pupa, [100].	[96]
	Note 4 Difference in structure of prolegs, distinguishing Aletia xylina from	[100]
	Anomis texana, [100].	
	Note 5	[100]
	Resemblance of larva of <i>Plusia dyaus</i> to that of <i>Aletia xylina</i> , [100]—Proportion of light and dark specimens early and late in the season, [100].	
	Note 6	[100]
	Cotton leaves blotched by young larva of Spilosoma acrea, [100].	
	Note 7	[100]
	Larva of Aletia xylina fed on Ipomæa eommutata, [100]—Abutilon and Phytolacca defoliated by different larvæ, [100]—Said to feed on "salve bush," [100].	
	Note 8	[100]

348 BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY. 2343. RILEY, C. V.—Continued. Notes—Continued. Note 9 [101] Notice of Wm. Trelease's "Nectar: what it is and some of its uses," [101]—Teleology and dysteleology of nectar glands of cottonplant, [101]. Rapidity with which the broods of A. xylina follow one another in midsnmmer, [101]—Time of first appearance of worms, [101]— Number of broods, [101]—Prolificacy of moth, [101]—Importance of natural checks upon its increase, [101]. Influence of winter temperature on time of first appearance of worms, [101]. Theories of hibernation of A. xylina, [101]—Proof of hibernation of moth, [102]—importance of this proof, [102]. Influence of latitude upon time of hatching of insects, [102]. Number of broods of A. xylina previously recognized, [102]. Possible food-plants of larva of A. xylina, [102]—There must be some besides cotton, [102]—Failure to find any other, [102]— Feeding of larva of Anomis erosa on Urena lobata, [102]-Value of fiber of U. lobata, [102]—Geographical distribution of U. lobata, [103]—Eggs and larva of Anomis erosa distinguished from those of Aletia xylina, [103]—Examination of malvaceous plants in herbarinm of U. S. Department of Agriculture, [103]—Disadvantages of such an examination, [103]—Plants on which eggs were found in the herbarinm, [103]—Petition for aid in obtaining evidence of the food-plant of A. xylina in the more northern States, [103]-List of malvaccons plants growing in these States, with localities, [103]. Note 16 [104] Description of larva of Aspila vireseens, [104] Note 17...... [104] Travels of E. A. Schwarz in 1878 and 1879, [104]—Reference to published accounts of the results of his investigations and those of others on hibernation of A. xylina, [104]. Platyhypena seabra, [104]—Characters of larva, [104]—Food-plants

of Jarva, [104]—Popation and hibernation, [104]—Characters of

Note 19 [104]

Seasons of larva of Phoberia atomaris, [104].

pupa, [104].

2343. RILEY, C. V.—Continued. Notes-Continued. References to discussion of J. P. Stelle's claim to have first recommended publicly the use of Paris green for A. xylina, [105]. References to descriptions and figures of brush-sacs of moths, [105]. Worms worse in wet weather than in dry, because more protected from enemies, [105]—Localities of first appearance of worms are those of least molestation, [105]. Note 25 [106] Appetite of swine for cotton-worms, [106]—Worms eaten by dogs and cats, [106]. Reference to list of birds of southern States, [106]. Range of English sparrow in United States, [106]—In hotter portions of the country it is confined to towns and villages, [106]. Report by Dr. Gco. Marx on spiders found on cotton, [106]—Preliminary list of spiders which destroy insects noxious to agriculture, [106]—List of spiders observed to devour larvæ of Aletia xylina, [106]—How they capture their victims, [106]—Food-habits of Theridula spharula, [107]—Observations on habits of Oxyopes viridans, by H. G. Hubbard, [107]—Ants captured by larvæ of a Cicindela, [107]. Description of image of Trichogramma pretiosa, [107]. Metamymar n. g. and M. aleurodis n. sp., provisionally named, [107].Description of image and larva of Apanteles aletice, [108]. Apanteles aletia parasited by Eupelmus sp., [108]. Description of image of Euplectrus comstockii, [108]. Description of image of Elachistus euplectri n. sp., [108]. Note 34 [109] Sarcophaga sarraceniæ distinct from S. carnaria, [109]-Points of difference between the Sarcophage of America and Europe generally, [109]—Additional specific characters of S. sarracenie, [109]. Description of image of Tachina aletia, [109]. Description of image of Tachina fraterna, [109]. Differences between tachinid larva and that of Sarcophaga, [109]— Differences of puparia, [109]—Reference to description of larva of Senometopia atropivora, [110]; of larva and pupa of Tachina villica, [110]—Description of larva and puparium of Sarcophaga sarracenia, [110]; of larva of Belvoisia bifasciata, [110]-Difference of larva of B. bifasciata from that of Tachina concinnata,

2343.	RILEY, C. V.—Continued.	
	Notes—Continued.	
	[110]—Structure of spiracles in normal form of tachinid puparium, [110]—Description of the puparium, [110]; of puparium of Belboisia bifasciata, [111].	
	Note 38	Г1111
	Synonymy of Cryptus conquisitor, [111]—C. pleurivinctus erroneously given as a synonym of C. annulicornis, [111].	[]
	Note 39	[111]
	Synonymy of Cryptus samia, [111].	
	Note 40	[111]
	Description of larva and pupa of <i>Chalcis ovata</i> , [111]—List of Lepidoptera parasited by <i>Ch. ovata</i> , [111]—Variations in size or imago, [111].	
	Note 41	[1117
	Synonymy of Tetrastichus esurus, [111]—Description of imago,	
	Γ1117.	
	Note 42	[111]
	Description of image of Hexaplasta zigzag, [111]—Difficulty of defining the families Chalcidide, Proctotrupide, and Cynipide, [112]—Food-habits of these families, [112].	
	Note 43	[112]
	Phora aletia not a true parasite, [112]—Habits of this species, [112].	
	Note 44	[112]
	The vast majority of the moths attracted to light said to be males, [112]—If this is so, the usefulness of fires and lights as a remedy is almost nothing, [112].	•
	Note 45	[112]
	Antidotes for arsenical poisoning, [112].	
	Note 46	[112]
	Non-fertilized blossoms destroyed by morning showers or spraying, [112]—When to make wet applications of poison, [112].	F1107
	Note 47	[II3]
	Note 48	Г1137
	Proportions of Paris green mixture, [113].	Litol
	Note 49	Γ 11 37
	Effect of kerosene and kerosene emulsion on cotton-plants, [113].	
	Note 50	
	Pyrethrum willemoti probably a synonym of P. roseum, [113]—Ref-	
	erence to Willemot's paper on P. willemoti, [113].	
	Note 51	[113]
	Growth of productive pyrethrum industry in California, [113]—Effect of pyrethrum on warm-blooded animals, [113]—Pyrethrum	
	recommended as a disinfectant and germicide, [113].	
	Note 52	Г1131
	Cost of production compared with price of pyrethrum, [113].	Litol
	Note 53	[113]
	Enemies of Aletia xylina killed by poisoning the worms, [113].	_ •
	Note 54	[113]
	Positiveness with which various plants have been recommended as insecticides, [113].	
	Note 55	[114]
	Insects injuring dog-fennel, [114],	

2343. RILEY, C. V.—Continued. Notes-Continued. Note 56 [114] Tests of spraying machinery, by Dr. W. S. Barnard, [114]—Diffieulties in the use of machinery in irregularly-planted fields, [114]—Means of overeoming these difficulties, [114]—Conveyances for underspraying apparatus, [115]—Management of apparatus, [115]—Rapidity of service, [115]—Quantity of poison required, [115]—Suecess of stirrer-pump device, [115]—Construction of stirrer-pump, [115]—Construction of nozzle-pipes, [116]—Adjustment of these pipes, [116]—Success of Eddy-chamber nozzles, [116]—A elosed system of pipes advisable, [116]— High pressure should always be used, [117]—Adjustment of deseending pipes, [117]—Flexile joints recommended, [117]—Advantages of flexile pipes, [117]—Lifting of desecnding pipes in turning, [118]—Devices for lateral shifting of pipes, [118]— Summary of eonelusions from experiments, [118]. Note 57 [119] Recommendation to use Paris green in 1872, [119]—Extract from report made on this subject by J. P. Stelle, in 1880, [119]—Comment on the same, [120]. Description of image of Anomis texana n. sp., [120]. Description of egg and pupa of Drasteria ercchtea, [120]. Reprint of J. W. Boddie's description, in 1850, of image of Phalana zea (Heliothis armigera), [121]. Reprint of A. R. Grote's description of Heliothis umbrosus, [121]. Remarks on specimens sent from Bahia, by R. A. Edes, in 1880, [121]—Characters of egg, larva, and pupa of Anomis sp., [121]— Insects found in cotton bolls, [121]. Probably Macrosila rustica stated to feed on cotton, [121]. Larvæ of Aletia xylina received from Vera Cruz, [121]—Recurrence of eotton-worms at irregular periods, [122]. 2344. RILEY, C. V. Reports of experiments with various insecticide substances, chiefly upon insects affecting garden crops, made under the direction of the entomologist. <Bull. No. 11, Div. Ent. U. S. Dept. Agric., [February 26], 1886, 34 pp. CONTENTS. Experiments with insecticides..... 5 Report of experiment sat La Fayette, Ind. By F. M. Webster 9 Report of experiments at Ames, Iowa. By H. Osborn 23 Report of experiments at Trenton, N. J. By T. Bennett 2345. [RILEY, C. V.] [The study of entomology.] < Entom. Amer., March, 1886, v. 1, pp. 225-227. Remarks made at a meeting of the A. A. S.; books useful for beginners,

- 2346. [RILEY, C. V.] [Habitat of Mezium americanum.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 14. Occurrence of Mezium americanum in old hay.
- 2347. [RILEY, C. V.] [Arctic insects.] < Proc. Ent. Soc. Wash., [30] March], 1886, v. 1, pp. 14-15. Remarks on collection of insects made at Point Barrow, Alaska; prevalence

in Arctic regions of species common to America and Europe; occurrence of Urocerus flavicornis in Alaska.

- 2348. [RILEY, C. V.] [Habits of Isosoma.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 15. Phytophagic habits of the genus Isosoma; I. grande observed ovipositing.
- 2349. [RILEY, C. V.] [Cranberry fruit-worm.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 15. Names Acrobasis vaccinii as the well-known cranberry fruit-worm.
- 2350. [RILEY, C. V.] [Rhyssa lunator.] < Proc. Ent. Soc. Wash., [30] - March], 1886, v. 1, p. 15. States that Rhyssa [= Thalessa] lunator is parasitic and not lignivorous.
- 2351. [RILEY, C. V.] [Tiphia and Rhipiphorus.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 15. Refers to O. Lugger's statements in regard to the habits of Tiphia and Rhipi-
- phorus as manifestly incorrect. 2352. [Riley, C. V.] [Insects attracted to light.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, pp. 15-16.

Disfigurement of buildings by the insects and spiders attracted to the electric light on the dome of the Capitol.

- 2353. [RILEY, C. V.] [Parasitic Coleoptera.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 16. Parasitic habits of Aleochara unthomyie [=nitida].
- 2354. [RILEY, C. V.] [Scenopinus.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 16. Larva of Scenopinus sp. infesting the blanket of a Navajo Indian.
- 2355. RILEY, C. V. Annual address of the president. < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, pp. 17-27.
 - Brief comments upon the histories or injuries of Agrotis fennica, Hadena devastatrix, Nematus erichsoni, Phytonomus punctatus, P. nigrirostris, Pulvinaria innumerabilis, Systachus leucophaus, and Phylloxera vastatrix; notes on the egg parasites of the Acridida; mode of oviposition of some Carabida; advantages of Washington from an entomological standpoint.
- 2356. [RILEY, C. V.] [Gall-making moths.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 30.

Walshia amorphella, previously known as a gall-maker on Amorpha fruticosa, bred from roots of loco weed; Euryptychia saligneana distinct from Pædisca scudderiana.

2357. [RILEY, C. V.] Sphida, Grote.] < Proc. Ent. Soc. Wash., [30] March], 1886, v. 1, p. 30.

Considers S. obliquata synonymous with Arzama densa.

- 2358. [RILEY, C. V.] [Food of Calopteron and Photinus.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 31.
 - Larva of Calopteron terminale gregarious and lignivorous; Photinus pyralis feeds on snails.
- 2359. RILEY, C. V. A carnivorous butterfly larva. <Science, 30 April, 1886, v. 7, p. 394.
 - Distribution and recorded food-habits of Feniseca tarquinius; Aphididæ the normal food of the same.
- 2360. RILEY, C. V. Entomology. Professor Riley to Dr. Shaffer. <Daily Globe [Keokuk, Iowa], 2 May, 1886. S.-b. No. 63, p. 30. Injuries of and means against Abia caprifolium [= Zarwa inflata].
- 2361. RILEY, C. V. A carnivorous butterfly larva.—Plant-feeding habit of Feniseca tarquinius. <Amer. Nat., June, 1886, v. 20, pp. 556-557.
 - Supposed food-plants of the larva of Feniseca tarquinius; first publication of proof that it feeds on plant-lice; Schizoneura tessellata, Pemphigus fraxinifolii, and P. imbricator the species preyed upon.
- 2362. RILEY, C. V. Thrips—Leaf hoppers. <Gardener's Mo. and Hortic., June, 1886, v. 28, p. 174. S.-b. No. 61, p. 56.
 - Letter to Duncan Rhind; remedies against Erythroneura vitifex [= Typhlocyba vitis].
- 2363. RILEY, C. V. Report of the entomologist. <Ann. Rept. [U. S.] Commissioner Agric. for 1885, 1886, pp. 207-343, 1 map, 9 pl. Separate: <Washington: June, 1886, pp. 10+137+10.

TABLE OF CONTENTS.

Introduction	207
Silk-culture, 207—Review of the divisional work in silk-culture,	
207—Necessity for a home market for cocoons, 205—Proposed es-	
tablishment of a filature at Washington, 208—Silk culture in	
France, 209—Destructive locusts, 209—Injurious insects of the	
year, 209—Correspondence of the Division, 210—Work of the Di-	
vision in economic ornithology, 210—Establishment of an api-	
cultural station, and the work of the Division in apiculture, 211—	
Work of other divisional agents, 212—Relations of the Division to	
the National Museum, 213—Publications of the Division, 213—	
Office force, 213.	
SILK-CULTURE	214
General work of the division, 214—Distribution of eggs, 214—States	214
in which greatest interest is felt, 214.	
**	214
Distribution of mulberry trees.	214
Number distributed, 214—Danger of indiscriminate distribution,	
214—Osage orange, 214—Russian mulberry, 215.	015
Distribution of eggs	215
Variety distributed, 215—Designation of races, 215.	
Home-raised vs. imported eggs.	215
Feeling against importation, 215—Reasons for importation last year,	
215—Provisions for the next distribution, 216—Eggs rejected,	
216—Labor involved in silk-culture, 216.	
Establishing of filatures—Cost of producing reeled silk	216

2363. RILEY, C. V.—Continued,

SILK-CULTURE—Continued.	
Importance of filatures, 216—Establishing of stations, 217—Work	v.
at Philadelphia station, 217—Work and results at New Orleans	
filature, 217—The Serrell automatic silk reel, 218—Application of	
the saving of this recl to New Orleans figures, 219.	
Work on the Pacific coast	219
Last season's experimental crop, 219—Incorporation of the Ladies'	
Silk-Culture Society of California, 219—The station at Piedmont,	
219-Proposed work at the station, 220-Work of the State board,	
220—Attempt to form a stock company, 220.	
On the precautions necessary in the production of pure silk-worm	
eggs and in properly wintering them	221
Pasteur's classification of diseases, 221.	
Flaceidity (Flacherie)	221
Symptoms and consequences, 221—Action of worms, 221—Color,	
221—Odor, 221—Nature and treatment, 222—Microscopic diagno-	
sis, 222—The bacillus of flaceidity, 222—Examination of the	
chrysalis, 222.	
Pébrine	223
Symptoms, 223—The black spots, 223—Microscopic diagnosis, 223—	
Appearance of chrysalis, 224—Tests for the determination of	
pébrine, 224—Difficulty of using the worm for test, 224—Advantage	
of using the moth, 224—Isolation and examination of the moths,	
224—Maillot's method, 224—Examination of chrysalides, 224—	
Selection of cocoons, 225—Moth cells, preparation, 225—Fasten-	
ings for cells, 225—Preparation of moth for examination, 226—	
The corpuscle of pébrinc, 226.	
Wintering the eggs	227
Colors of eggs, 227—The micropyle, 227—Proper temperature and	
humidity, 227—Hibernating boxes, 228—Acknowledgments, 228.	
MISCELLANEOUS INSECTS	228
Destructive locusts or "grasshoppers," Acridida	228
Unusual damage, 228—The Rocky Mountain locust, 228—Proba-	,
bilities for 1886, 229—The California migratory locust, 229—	
Abundance the present year, 229—The destructive species in	
previous years, 230—Geographical range, 230—Characters of the	
species, 231—Remedies, 232—New remedy, 232—Non-migratory	
species, 232—The red-legged locust, 232—The differential locust,	
233—The two-striped locust, 233—The lesser locust, 233.	
The periodical Cicada, Tibicen septendecim	233
Preparation of a Bulletin, 233—A seventeen-year and a thirteen-	
year race, 234—Our announcement in 1868, 234—Dr. Smith's man-	
uscript, 234—Dr. Phares's earlier discovery, 234—Two distinct	
forms or varieties; specific value of the different forms, 234—	
First discovery, 234—Points of difference, 235—How the species	
should be catalogued, 235—The long period of underground de-	
velopment, 235—The food of the larva, 236—Method of burrow-	
ing of the larva, 237—The transformations, 237—Rising of the	
pupæ, 237—Five phases of ecdysis, 238—Time required for trans-	
formation, 238—The Cicada vs. civilization, 238—Influence of	
domestic animals, 238; of building of towns, 238; of the En-	
glish sparrow, 238—Song notes of the Cicada, 239—Notes of other	
insects, 239—First or phar-r-r-aoh note, 239—Second note or	
"screech," 240-Third or "chirping" note, 240-Other notes,	

258

265

270

276

2363. RILEY, C. V.—Continued.

MISCELLANEOUS INSECTS-Continued.

240-Variation in time of appearance, 240-Enemies of the Cicada, 241—Supposed sting of the Cicada, 241—Oviposition, 242-Injury caused to fruit trees; remedies and preventive measures, 242—Small injury in larva state, 243—Destruction of the insects as they issue from the ground, 243; by Pyrethrum powder, 243; by Pyrethrum water, 243; by kerosene emulsion, 244— Experiments with substances that kill by thwarting exnviation, 244—Carbolic acid, 244—Acetic acid, 245—Alcohol, 245—To prevent ovipositing, 245—Kerosene emulsion, 245—Consideration of the two 1885 broods, 246—Circular letter, 246—Brood VII, 247— Brood XXII, 248—Points of contact of the two broods, 249— Geographical distribution of Brood VII, 250-Distribution of Brood XXII, 251-Summary of distribution and future appearance of different broods, 252-Influence of climate upon the races, 254—Experiment in the transfer of eggs, 255—Persons assisting, 255—Description of the placing of the different lots of eggs, 256—The Cicada in 1886, 257.

The garden web-worm, Eurycreon rantalis.

Great damage the present year, 265—A wide-spread species, 265—Popular descriptions of different states, 265—Former injuries, 266—Localities of damage in 1885, 266—Food-plants, 267—Habits and natural history, 267—Number of annual generations, 267—Habits of larvæ, 268—Cocoon, 269—Natural enemies, 269—Remedies, 269—Arsenical poisons, 269—Machine for jarring the worms from the plants, 270.

The dark-sided cut-worm, Agrotis messoria

Injury to onions; new habit, 270—Habits and natural history, 270—Number of broods, 270—Hibernation, 271—Remedies, 272—Treatment of land in spring, 272—Poisoned balls, 272—Kerosene emulsions, 272—Report of John B. Smith, 273—Report of Thomas Bennett, 274.

The strawberry weevil, Anthonomus musculus.

Past history, 276—In Maryland, 276—In Missouri, 276—In Michigan, 276—Injury in 1885, 276—Damage on Staten Island, 276—Habits and natural history, 277—Mr. Smith's observations, 277—Diversity of habit in the genus, 278—Natural history of other species of Anthonomus, 278—Inquilinous species, 279—Speices having different habits, 279—Remedies, 279—Wolf's soap, 279—Kerosenc emulsion, 280—Pyrethrum, 280—Repellants, 280—Characters and synonomy, 280—Descriptive, 281—Variations of A. musculus, 282—Comparative differences between A. musculus and A. suturalis, 282.

2363. RILEY, C. V.—Continued.

MISCELLANEOUS INSECTS—Continued.

Decisive steps for eradication recommended, 283—Life-history and habits, 283—Laying of eggs and growth of larvæ, 283—Habits of larvæ, 284—The cocoon, 284—One annual generation; dates of issuance of adults, 284—Parasites, 285—Remedies, 285—Is it an introduced species?, 285—Reasons for so considering it, 285—Schmidberger's account of the European pear midge, 286—Dr. Joseph Mik's opinion, 287—Descriptive, 287—Imago, 287—Larva, 288—Pupa, 288—The classificatory value of the genitalia, 288.

REPORTS OF AGENTS.

289

Report on the locusts of the San Joaquin Valley, Cal. By D. W. Coquillett

289

Letter of transmittal, 289—Letter of instructions, 289—Location, 290—Work of the locusts in general, 290—Species most destructive, 291—The devastating locust, 291—The ash-colored locust, 296— 292—Where did these locusts hatch?, 292—Consideration of surrounding topography, 293—How the young locusts may be distinguished, 293-Influence of submersion on hatching, 294-Cause of the abundance in this valley in 1885, 294—Early hatching, 294— Lack of rain, 294—The differential locust, 295—Abundance, 295— Egg-pod, 295—Oviposition, 296—Flight, 296—The yellow locust, Abundance, 296—Flight, 296—Cannibalism, 296—Oviposition, 297-Other species of locusts, 297-Table showing relative abundance, 297—Injury committed by the locusts, 297—To grape-vines, 298—To grain-fields and vegetable-gardens, 298—Natural enemies, 298—Birds and poultry, 298—Wasps, 298—Mites, 299—Flies, 299— Remedies, 299—Collecting in windrows of dry stems and then burning, 299—Rolling, 300—Kerosene emulsion, 300—Burning sulphur, 300—Horse troughs, 300—Bran, arsenic, and sugar trappoisoning, 300-Buhach, 302.

Report on the abundance of the Rocky Mountain locust in 1885. By
L. Bruner.

303

Letter of submittal, 303—The great abundance of native species, 303—Direction of flight of the Rocky Mountain species, 304—Probabilities for 1886, 305—Effect of weather upon probabilities, 305—Scarcity of parasites, 305—Abundance of Melanoplus atlanis and Camnula pellucida, 306—List of locusts noticed in larger numbers than usual, 307.

308

311

333

2363. RILEY, C. V.—Continued.

REPORTS OF AGENTS-Continued.

332—Descriptive, 333.

products, 343.

Third report on the eauses of destruction of the evergreen and other forest trees in northern New England. By A. S. Packard 319 General review, 320—Condition of the spruce on the east of Maine, 321-Exemption from insect attack, 321-Condition of the hackmatack in 1885, 321—Verification of last year's predictions, 322— The white-pine weevil, 322—Eggs, 322—Larva, 323—Pupa and its cell, 323—Distinguished from other species, 323—Life-history in brief, 324—Effects of its work, 324—Prevention, 325—The spruce Epizeuxis, 325—Discovery of food-plant, 325—Compared with other species, 325—Descriptive, 326—The spruce plume-moth, 326— Habits, 326—Descriptive, 326—The pine Pheocyma, 327—Tho evergreen Cleora, 327-Habits, 327-Descriptive, 327-The fir Paraphia, 328—Descriptive, 328—The spruce Therina, 328—Deseriptive, 329—The pine Therina, 329—The pine Amorbia, 329— The V-marked Cacecia, 329—Habits, 329—Descriptive, 330—Tho

hiekory Eecopsis, 330—Habits, 330—Descriptive, 331—The variegated Eccopsis, 331—Déscriptive, 331—The white-heart hickory Geleehia, 331—Habits, 332—Descriptive, 332—The willow Teras, 332—Descriptive, 332—The purple willow Gracilaria, 332—Habits,

Report on experiments in apiculture. By N. W. McLain......

Letter of submittal, 333—Economy in the production of wax, 334—
Experiments in wax-feeding, 334—Device for feeding sugar syrup and for other purposes, 334—Description, 334—Wintering bees, 335—Loss from thirst, 335—"Roaring in the hive," 336—Device for furnishing water, 336—Bees vs. fruit, 336—House built for experiment, 336—Details of experiments, 337—Conclusions, 338—Fruits used, 339—Experience of Richard Rees, 339—Artificial fertilization, 339—Desirability of accomplishing it, 339—Recent experiments, 340—Experiments with larval and pupal queens unsuccessful, 340—Experiments with imago queens, 341—Conclusions and ground for hope, 342—Bee forage, 342—Improved races, 342—Importance of apiculture, 342—Statistics, 343—Necessity and scope, 343—Desirability of accurate erop reports of apiarian

2364. RILEY, C. V. Miscellaneous notes on the work of the Division of Entomology for the season of 1885, prepared by the Entomologist. <Bull. No. 12, Div. Ent. U. S. Dept. Agric. [13 July], 1886, 46 pp. 1 plate.

CONTENTS.

Report on the production and manufacture of Buhaeh. By D. W. Co-	
quillett	7
Additions to the third report on the eauses of the destruction of the	
evergreen and other forest trees in northern New England. By A. S.	
Packard, jr	17
The periodical Cicada in southeastern Indiana. By A. W. Butler	24
Notes of the year	32

2365. R[ILEY], C. V. Hessian-fly. <Suppl. Encyclopædia Britannica, 9th Ed., 1886, v. 3, p. 324, fig.

Natural history, remedies, and figures of Cecidomyia destructor.

- 2366. R[ILEY], C. V. Locust. <Suppl. Encyclopædia Britannica, 9th Ed., 1886, v. 3, pp. 626-628, fig., map.
 - Ravages, life-history, enemies of, and means against Caloptenus spretus; migrations and their cause; map of the region overrun; figure of adult.
- 2367. RILEY, C. V. Some popular fallacies and some new facts regarding Cicada septendecim L. <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, p. 334.
 - Variety cassinii is not the race tredecim of Tibicen septendecim; twigs with eggs do not necessarily break off or die to insure the hatching of the larva.
- 2368. RILEY, C. V. Report of the curator of the department of insects in the U. S. National Museum for 1884. <Ann. Rept. of Regents of Smith. Inst. for 1884, 1885 [1886], pp. 185-188.
 - List of accessions, with notes; recommendations for utilizing the Glover plates; needs of the Department.
- 2369. RILEY, C. V. Notes on Feniseca tarquinius, Fabr. <Ca. Ent., October, 1886, v. 18, pp. 191-193.
 - Comments on article by W. H. Edwards; records made at the Department of Agriculture proving the carnivorous habits of the larva of Feniseca tarquinius.
- 2370. RILEY, C. V. Two useful lives. <Sci. Amer., 29 January, 1887, v. 56, p. 64. S.-b. No. 63, p. 122. See: <Wine and Fruit Grower, January, 1887, v. 9, p. 10. S.-b. No. 61, pp. 55-56. Brief obituary notices of Louis Bazille and Jules Lichtenstein.
- 2371. RILEY, C. V. Mr. Hulst's observations on *Pronuba yuccasella*. <Entom. Amer., March, 1887, v. 2, pp. 233–236.
 - Criticism of the observations of G. D. Hulst; summary of the method of oviposition of *Pronuba yuccasella*; pollination of *Yucca* by the same.
- 2372. [RILEY, C. V.] Fruit pest extermination. <San Diego Mirror, 5 April, 1887. S.-b. No. 63, pp. 88-89.

 Means against Coccide.
- 2373. [RILEY, C. V.] [Californian orange insects.] < Daily Herald. [Los Angeles], 9 April, 1887. S.-b. No. 63, p. 88.

 Means against Coccidæ.
- 2374. [RILEY, C. V.] Our bugs. <San Francisco Daily Examiner, 16
 April, 1887. S.-b. No. 63, p. 80; 86–87.
 Food-plants, ravages, and means against the white-scale.
- 2375. [RILEY, C. V.] [Remedies and appliances.] < Press and Horticulturist, 16 April, 1887. S.-b. No. 61, pp. 82-84.
 - Replies to inquiries made at the annual convention of the State Board of Horticulture of California; means against the codlin-moth; methods of spraying; differences between allied scales; their methods of dispersal and means against them.
- 2376. RILEY, C. V. Bumble-bees vs. red-clover. <Rural New-Yorker, 23 April, 1887, v. 46, p. 270. S.-b. No. 61, pp. 36-37.
 - Revival of unsettled questions; red-clover sterile in the absence of bumble-bees in New Zealand; some fertilization by other insects probable; introduction of bumble-bees into New Zealand; their rapid propagation there; beneficial effects on red-clover.

2377. RILEY, C. V. Young grasshoppers. <San Francisco Examiner, 25 April, 1887. S.-b. No. 63, p. 85.

Interview with reporter; ravages of and means against Melanoplus [= Caloptenus] devastator.

2378. RILEY, C. V. Our shade trees and their insect defoliators. Being a consideration of the four most injurious species which affect the trees of the capital; with means of destroying them. <Bull. No. 10, Div. Ent. U. S. Dept. Agric. [7 May], 1887, 69, pp. 27 figs.

CONTENTS.

LETTER OF SUBMITTAL	5
Introduction	7
FOUR PRINCIPAL LEAF-EATERS	8
The imported elm leaf-beetle, Galeruca xanthomelana	8
An importation from Europe	8
Habits and natural history	8
Remedies	11
More recent experience at the Dcpartment	14
Past history of the clms in question, 14—Condition and charac-	
teristics of the grove in 1882 and 1883, 14—Extent of injury in	
1882 and 1883, 14—Preferences of the elm beetles for certain	
varieties, 15—Effects of arsenical poisons on insect and plant,	
15—Preventive effects of the poison, 16—Treatment with Lon-	
don purple, 17—Preparation of the poison, 17—Effects of the	
mixture, 17—Treatment with Paris green, 18—Mechanical	
means of applying the poison, 19.	
The bag-worm, Thyridopteryx ephemeraformis	22
Habits and natural history	22
The eggs, 22—The larva and its bag, 23—Pupation, 25—The	
imago or perfect insect, 25.	
Geographical distribution	26
Food-plants	27
Enemies	27
The white-marked tussock-moth, Orgyia leucostigma	29
Habits and natural history	29
The eggs, 29—Development and characters of the larva, 29—	
Habits of the larva, 30—Pupation, 30—The imago, 30—Hiberna-	
tion, 31—Number of annual generations, 31.	
Food-plants	31
Natural enemies and parasites	31
Geographical distribution	33
The fall web-worm, Hyphantria cunea	33
Natural history	33
Limitation of broods, 33—The eggs, 34—The larva, 35—Pupa and	
cocoon, 36—The moth, 36.	
Injury done in 1886	37
Proportionate injury to different plants and shade trees	40
Peculiar effect of defoliation upon some plants	42
Enemies of the web-worm other than insects	43
Predaceous insect enemies.	44
Fungus diseases of the web-worm	46
Experiments to obtain percentage of diseased caterpillars, 47.	

2378. RILEY, C. V.—Continued.
Contents—Continued.
The parasites of the web-worm
Telenomus bifidus Riley, 48—Meteorus hyphantriæ Riley, 49—Apan-
teles hyphantriæ Riley, 50—Limneria pallipes Prov., 51—Tachina
sp., 52.
SUMMARY OF THE HABITS OF THE FOUR SPECIES. 53
Remedies and preventive measures
Winter work55
One simple preventive remedy for all
Pruning and burning
Mulehing
Influence of tree-boxes
Whitewashing of trunks
BIRDS: THE ENGLISH SPARROW
THE FUTURE OF OUR TREES. PRUNING
TREES WHICH ARE UNINJURED 64 GOOD AND BAD EFFECTS OF OUR TREES 64
Index
2379. RILEY, C. V. Remarks on the insect defoliators of our shade-
trees. <new 1887,="" pp.<="" t-p.+12="" td="" york,=""></new>
Report of address made before the New York farmers, 10 March, 1887; ravages
and means against Galeruca xanthomelana, Thyridopteryx ephemeraformis,
Orgyia leueostigma and Hyphantria eunea.
2380. RILEY, C. V Variable moulting in Orgyia. <ent. mag.,<="" mo.="" td=""></ent.>
May, 1887, v. 23, p. 274.
The female of Orgyia levelstigma undergoes four molts, the male undergoes
three; tendency of individual larvæ of all orders to vary from the normal number in the species; whenever there is a discrepancy in size of the sexes
the smaller undergoes a less number of molts; number of molts when
not sexual dependent on food supply; molting correlated with rate of
growth and nutrition.
2381. RILEY, C. V. Pedigree moth-breeding. <ent. mag.,="" may,<="" mo.="" td=""></ent.>
1887, v. 23, pp. 277–278.
Recommends Seriearia mori as the most favorable insect to experiment with;
its tendency to vary under new conditions.
2382. RILEY, C. V. Reports of observations and experiments in the
practical work of the division, made under the direction of the
· ·
entomologist. <bull. 13,="" agric.<="" dept.="" div.="" ent.="" no.="" s.="" td="" u.=""></bull.>
[3 June], 1887, 78 pp., 4 figs.
CONTENTS.
Introduction
Report on locusts in Texas in the spring of 1886. By L. Bruner 9
Fourth report on insects injuring forest and shade trees. By A. S. Pack-
ard, jr
Report on Nebraska insects. By L. Bruner
Tests with insecticides on garden insects. By W. B. Alwood
Report on Ohio insects. By W. B. Alwood

	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY. 3	61
	RILEY, C. V.—Continued. CONTENTS—Continued. A record of some experiments relating to the effect of the puncture of some hemipterous insects upon shrnbs, fruits, and grains, 1886. By F. M. Webster. Notes from Missouri for the season of 1886. By M. E. Murtfeldt	54 59 66
2383.	RILEY, C. V. A new apple pest. <sci. 11="" 112.="" 138;="" 148.="" 18="" 185.="" 1887,="" 216.="" 29,="" 384.="" 40,="" 50.="" 56,="" 61,="" <colman's="" <gardene="" adult="" amer.,="" and="" arsenical="" by="" description="" food-plants,="" habits,="" haltica="" hortic.,="" july,="" june,="" larva="" mo.="" no.="" of="" p.="" poisons.<="" punctipent="" rural="" sb.="" species="" subdued="" td="" the="" v.="" world,=""><td>23 er's 11 ;</td></sci.>	23 er's 11 ;
2384.	RILEY, C. V. A destructive cricket in Louisiana. <floridal 1,="" 1887,="" 1888,="" 2="" 20="" 576.="" 61,="" 64-reprint:="" 7,="" 87-88.="" <insect="" against="" and="" cotton,="" destructive="" dispatch,="" gryllus="" irish="" june,="" life,="" means="" no.="" october,="" p.="" peas,="" potatoes.<="" pp.="" ravages="" sb.="" sp.,="" sweet="" td="" to="" tobac="" v.=""><td>65.</td></floridal>	65.
2385.	RILEY, C. V. Strawberry borers. < Pacific Rural Press, 25 June 1887, v. 33, p. 559. Sb. No. 61, p. 90; No. 63, p. 110. Letter to I. A. Wilcox; recommends bisulphide of carbon as a means again Egeria impropria.	
2386.	RILEY, C. V. Life-history of the Icerya. <pacific 18="" 1887,="" 2="" 2394="" 25="" 33-34,="" 565;="" 60-64,="" 61,="" 9="" 9.="" advance="" agriculture="" commissioner="" contents.<="" figs.="" for="" from="" july,="" june,="" no.="" of="" p.="" pre="" proof="" report="" rural="" s.="" sb.="" see="" synopsis="" td="" u.="" v.=""><td>pp.</td></pacific>	pp.
2387.	[RILEY, C. V.] Cut-worms. < Pacific Rural Press, 25 June, 18 v. 33, p. 578. Sb. No. 63, p. 108. Means against cut-worms.	87,
2388.	RILEY, C. V. Reports of observations and experiments in the practical work of the division made under the direction of the entomologist. <bull. 1="" 14,="" 1887,="" 4="" 62="" [3="" agr.="" august],="" dept.="" div.="" ent.="" figs.<="" no.="" pl.,="" pp.,="" s.="" td="" u.=""><td>the</td></bull.>	the
	CONTENTS.	
	Introduction	7

Introduction	7
Report on insects injurious to garden crops in Florida. By W. H. Ash-	·
mead	9
Report on buffalo gnats. By F. M. Webster	29
Native plums. How to fruit them. They are claimed to be practically	
Curculio-proof. By D. B. Wier	39
The Serrell automatic silk-reel. By P. Walker	52
2389. RILEY, C. V. The Icerya or fluted scale, otherwise known as the	he
cottony cushion-scale. <bull. 15,="" del<="" div.="" ent.="" no.="" s.="" th="" u.=""><th>t.</th></bull.>	t.
Agric. [18 August], 1887, 40 pp. <san 16="" apr<="" diego="" th="" union,=""><th>il,</th></san>	il,
1887. Sb. No. 61, pp. 90-97. <los 13="" 188<="" angeles,="" april,="" th=""><th>37.</th></los>	37.

2389. RILEY, C. V.—Continued.

S.-b. No. 61, pp. 97-104. <Los Angeles Tribune, 14 April, 1887. S.-b. No. 61, pp. 104-111. See: <Pacific Rural Press, 23 April, 1887, v. 33, pp. 361-362; 364. S.-b. No. 61, p. 48; No. 63, p. 76. <Riverside Daily Press, 12, 13 April, 1887. S.-b. No. 61, pp. 65-68; No. 63, pp. 98-105. <Press and Horticulturist, 16 April, 1887. S.-b. No. 61, pp. 74-82. <Florida Dispatch, 2 May, 1887, v. 7, pp. 385-388. S.-b. No. 63, pp. 75; 77. <Pacific Rural Press, 2 July, 1887, v. 34, p. 9.

CONTENTS.

Letter of submittal	5
Introduction	7
The scale-insects of the orange in California, and particularly the	
Icerya or fluted scale, etc	9
Notes on Icerya-Its probable origin the Islands of Bourbon and	
Mauritius	27
The use of gases against scale-insects.	35

2390. [RILEY, C.V.] [Classification of insects.] < Entom. Amer., September 1887, v. 3, p. 102.

Comments on the address of J. H. Comstock before the A. A. A. S.

- 2391. [RILEY, C. V.] [Pronuba and its connection with the pollination of Yucca.] < Entom. Amer., September, 1887, v. 3, pp. 107-108. Record of the results of recent experiments on the pollination of Yucca and the agency of Pronuba in this work.
- 2392. RILEY, C. V. Beschreibung einer den Birnen schädlichen Gallmücke (? Diplosis nigra Meig.). <Wiener Entomol. Zeit., September, 1887, v. 6, pp. 201–206, 3 figs.
 - Discovery of the species in America; its habits and life-history; detailed description of all stages; possible difference between European and American specimens; suggests *D. pyrivora* for the latter, if distinct.
- 2393. RILEY, C. V. Some important discoveries in the life-history of the hop-plant louse (*Phorodon humuli* Schrank). <Soc. for Prom. Agric. Sci., September, 1887, 1, No. 9, p. 205. Reprint: <Sci. Amer. Suppl., 24 September, 1887, v. 24, p. 9781. S.-b. No. 61, pp. 117–119. <Gardener's Mo. and Hortic., October, 1887, v. pp. 369–311. S.-b. No. 61, p. 124.
 - Résumé of recent discoveries in the life-history of *Phorodon humuli*; proof of its migration from plum to hop; life of the egg on plum in winter; spring migration to hop; number of broods thus far observed; probable course of later broods.
- 2394. RILEY, C. V. Report of the entomologist. <Ann. Rept. [U. S.] Commissioner Agric. for 1886, 1887, pp. 459-592, 11 pl. Separate: <Washington, September, 1887, pp. 459-592+6+9, 11 pl.

TABLE OF CONTENTS.

2394. RILEY, C. V.—Continued-

TABLE OF CONTENTS—Continued.

and canstie potash, 459—Remedy for the California red-seale (Aspidiotus aurantii), 460—Experiments made by Mr. Coquillett and Mr. Koebele, 460—Emulsifying oil, 460—Southern bnffalo gnats, 461—Results of investigations made on the Southern bnffalo gnat and the turkey gnat, 461—Common fall web-worm (Hyphantria cunea), 461—Imported elm-leaf beetle, 461—Spraying appliances, 461—Alternation of generation with the jointworms, 462—Injurions insects of the year, 462—Hop Aphis (Phorodon humuli), 462—Serrell antomatic reel for silk, 462—Apicultnral station at Anrora, Ill., 462—Work of the Division of economic ornithology, 462—International exhibition of machinery and contrivances for applied remedies against fungiand insects at Florence, 464—Publications of the Division during the year, 464—Proposed publications, 464—Work of field agents, 465—Office force, 465.

Introductory, 466—Geographical distribution, 466—In Australia, 466-In Cape Colony, 467-In New Zealand, 467-Importation of the species into California, 468-Its spread and present limitation in California, 469—Food-plants, 471—Original food-plant of Icerya purchasi, 471-Its food-plants in Sonth Africa, 471-Its food-plants in New Zealand, 472—Its food-plants in California, 472—Characters and life-history, 474—The egg, 475— The female larva, first stage, 475—Female larva, second stage, 476-Female larva, third stage, 476-The adult female, fourth stage, 477—The egg-sae, 478—The male larva, probable seeond stage, 478-Male larva, third stage, 479-The male pupa and eoeoon, 479—The adult male, 480—Rate of growth of the different stages, 481—Habits, 481—Exndation of the honeydew, 482-Mode of spread and distribution, 483-Natural enemies, 484—Birds, 484—Predaeeous insects, 484—Parasites, 487— Remedies and preventive measures, 488-Importation of parasites, 488—Preventive action, 489—Spraying with insecticides, 489—Fumigating, 491—Bandages around the trunk, 491—Conclusion, 491.

Geographical distribution, 493—Early history, 493—Time of appearance, 494—Duration of an invasion, 494—Character of a swarm, 495—Mode of attack, 496—Animals injured, 497—Effect of the bites, 497—How animals protect themselves, 498—Preventives, 499—Remedies for the bites, 501—Attacking man, 501—Damage done in various years, 501—Popular opinions about the early states of the buffalo gnats, 502—Habits and natural history, 503—The egg, 503—The larva, 505—Habits of the larvæ, 505—Food of the larvæ, 507—Pupa and eoeoon, 508—The imago, 509—Number of broods, 509—Enemies of the buffalo gnat, 510—Descriptive, 511—Simulium pecuarum n. sp., 512—Simulium meridionale n. sp., 513—Remedies tried and proposed against the larvæ, 514—Overflows and buffalo gnats, 515.

2394. RILEY, C. V.—Continued.

TABLE-OF CONTENTS—Continued.	
The fall web-worm, Hyphantria cunea. Natural history, 518—Limitation of broods, 518—The eggs, 519—	518
The larvæ, 519—Pupa and cocoon, 520—The eggs, 513—	
done in 1886, 521—Proportionate injury to different plants and	
shade trees, 522—Peculiar effect of defoliation upon some	
plants, 525—Enemies of the web-worm other than insects,	
525—Predaceons insect enemies, 526—Fungus disease of the	
web-worm, 527—Experiments to obtain percentage of discased	
caterpillars, 529—True parasites of the web-worm, 530—Sec-	
ondary parasites, 530—The Telenomns egg-parasite, 531—The	
Meteorns parasite of the web-worm, 531—The Microgaster par-	
asite of the web-worm, 533—The Limneria parasite of the web-	
worm, 534—The Tachina parasite of the web-worm, 534—Rem-	
edies, 535—Pruning and burning, 535—Mulching, 536—Arsen-	
ical poisons, 536—Emulsions of kerosene, 538—Naphtha, 538.	539
Joint-worms, Isosoma sp	555
539—Other recent appearances, 540—Its identity with <i>I. nigrum</i>	
Cook, 541—Parasites, 542—The wheat-straw Isosoma, 542—	
Dimorphism of I. tritici and I. grande, 543—Occurs in California,	
544—In Kansas, 545—Parasites, 546.	
SILK CULTURE	546
Appropriations for the current year, 546—Establishment of a fil-	
ature at Washington, 546—Osage orange vs. mulberry, 546—	
Rendition of osage cocoons, 547—Mr. Serrell's opinion, 547—	•
Satinage, 547—The Serrell reel; cost of work up to the present	
time, 548—Expenses tabulated, 548—Chances for improvement,	
549—Distribution of eggs, 549—Reasons for purchasing foreign	
eggs, 549—Improper choking of cocoons, 550—A simple apparatus for choking, 551—Cocoons produced in the United States	
in 1886, 551—Tabulated by States, 552.	
REPORT OF AGENTS.	552
Report on remedies for the cottony cushion-scale. By D. W. Co-	
quillett	552
Letter of transmittal, 552—General considerations, 553—Canstic	
potash, 554—Caustic soda, 555—Hard soap, 555—Soft soap, 555—	
Kerosene emulsions, 556—Tobacco, 556—Sheep dip, 557—To-	
bacco soap, 557—Vinegar, 557—Paris green, 557.	
Report upon supplementary experiments on the cottony cushion-	
scale, followed by a report on experiments on the red-scale.	
By A. Koebele Letter of transmittal, .558—Introductory, .558—Soap solutions,	558
558—Preparation of soap, 559—Resin compounds, 559—Lye so-	
lution, 560—Bisnlphide of carbon, 560—Kerosene emulsion,	
560—Experiments, 560—Experiments on fumigation with bi-	
snlphide of carbon, 569—Experiments on red-scale (Aspidiotus	
aurantii), 569.	
Insects affecting small grains and grasses. By F. M. Webster	573
Letter of transmittal, 573.	
Insects affecting fall wheat	573
The wheat-straw Isosoma (Isosoma tritici, Riley), 573—The Amer-	
ican Mcromyza (Meromyza americana, Fitch), 574—The com-	
panion wheat fly (Oscinis is sp.), 574.	

2394. RILEY, C. V.—Continued.	
TABLE OF CONTENTS—Continued.	
Insects affecting barley	575
The white-grub (Lachnosterna fusca, Frohl.), 575—The barley	
root-louse (Schizoneura sp.), 575—The grain Aphis (Siphonophora	
avenæ, Fab.), 576.	
Insects frequenting or depredating upon buckwheat	576
Supposed exemption from insects, 576—Mode of observation,	
576—Signification of symbols, 576—List of insects, 576.	
Insects affecting timothy, 578.	
The glassy cut-worm (Hadena devastatrix, Brace), 578—Outbreak	
in Indiana in 1886, 578—Doubt as to species concerned, 578—	
Method of work, 579—Traveling, 579—Parasites, 579—Damage,	
580-Condition of fields in October, 580-The grain Spheno-	•
phorus (Sphenophorus parvulus, Gyll.), 580.	
Insects affecting white-clover	580
The flavescent clover weevil (Sitones flavescens, Allard), 580-	
Infests alsike, 580—Mode of attack, 580—Distribution, 580—	
Eggs and larvæ, 580—Hibernation, 580—Summary of life-his-	
tory, 581—The clover-stem maggot (Oscinis sp.), 582—Descrip-	
tive, 581—Time and place of oviposition, 581—Number of	
broods, 581.	
Report on experiments in apiculture. By N. W. McLain	58
Letter of transmittal, 583—The "quaking disease," 583—Nature	
of the disease, 583—Odor of infested hives, 583—Treatment	
with brine and soda, 583—Another form of the disease, 584—	
The foul-brood disease, 584—It is contagious, 584—Live pollen	
the medium, 584—Treatment, 584—Results of treatment, 585—	
How the disease spreads, 587—The control of reproduction,	
587—Difficulties, 587—Experiments in 1885, 588—Device for	
holding the queen, 588—Differences in drones, 589—Proportion	
of impotent individuals, 589—Mr. Baldwin's experiments at	
Clarksville, Mo., 589—Comments, 590—Fertilization in con-	
finement, 590—Apparatus, 590—Results of experiments, 590—	
Proposed experiments, 591.	
2395. RILEY, C. V. The Hessian-fly in England; its origin; its I	past
its future. <london 17="" 1887.="" no<="" october,="" sb.="" td="" times,=""><td>61</td></london>	61
pp. 144–147.	
Date of the introduction of Cecidomyia destructor into England; proba	hilita
that it has been introduced from continental Europe since the tin	•
Curtis; conditions in Great Britain unfavorable for its increase.	по о
•	, ,
2396. RILEY, C. V. The problem of the hop-plant louse fully sol	
. (Gardener's Chronicle, 22 October, 1887. Sb. No. 61	, pp
133-135. Reprint: < Mark Lane Express, 31 October, 188	37, v
57, pp. 135–137. S. b. No. 63, pp. 136–140.	
Life-history, migrations, and mode of hibernation of Phorodon humuli.	
2397. RILEY, C. V. On the luminous larviform females of the Phe	on ac
2001. INILEY, O. V. OH THE TUMINOUS TAILVIIOTHI TEMPATES OF THE FIRE	Jugo.

dini. <Ent. Mo. Mag., [December], 1887, v. 24, pp. 148-149.

Structural characters of the larval Phengodini; food of Zarhipis; characters of eggs, young larvæ, and female larvæ of Zarhipis and Phengodes; female Phengodini considered an archetypal hexapodal form; relations between phosphorescence and differentiation of the sexes.

- 2398. RILEY, C. V. [Introduction of the Hessian-fly into England.] <Trans. Ent. Soc. London, [December], 1887, pp. 45-48, Proc. Discussion of the date of introduction of Cecidomyia destructor into America and England; concludes from a study of the historical evidence, the distribution and parasites of the insect, that it was probably introduced into England about three or four years ago.
- 2399. RILEY, C. V. Poisonous insects. <Reference Handbook of the Medical Sciences, 1887, v. 5, pp. 741-760, figs. 2971-3020.

 An exhaustive illustrative review of the Arachnida, Myriapoda, and Hexapoda which secrete a poison injurious to man; descriptions of their life-histories; the manner in which the injury is inflicted and the remedies for the same.
- 2400. RILEY, C. V. The problem of the hop-plant louse [Phorodon humuli, Schrank] in Europe and America. <Rept. Brit. Assoc. Adv. Sci., 1887, pp. 750-753. Separate: <pp. 1-3. See: <Nature, 13 October, 1887, v. 36, pp. 566-567. <Gardener's Chronicle, 17 September, 1887. S.-b. No. 61, pp. 333-334.
 - Life-habits of *Phorodon humuli*; effects of extreme heat and of meterological conditions; natural enemies and means against the *Phorodon*.
- 2401. RILEY, C. V. On *Icerya purchasi*, an insect injurious to fruit trees. <Rept. Brit. Assoc. Adv. Sci., 1887, p. 767. Separate: <p. 1. See: <Nature, 20 October, 1887, v. 36, p. 592.
 - Summary statement of food-plants, original home, synonomy, and means against *Icerya purchasi*.
- 2402. RILEY, C. V. On the luminous larviform females in the *Phengodini*. <Rept. Brit. Assoc. Adv. Sci., 1887, pp. 760-761. Separate: <pp. 1-2. See: <Entom. Amer., September, 1887, v. 3, p. 107. <Proc. Amer. Assoc. Adv. Sci., 1887 [May, 1888], v. 36, p. 262.
 - Résumé of facts relating to the history, characters, and life-habits of the luminous larviform females in the *Phengodini*; bearing of these facts on the theory of evolution.
- 2403. RILEY, C. V. [Scale on Euonymus latifolia?] <Sci. Amer., 14

 January, 1888, v. 58, p. 27. S.-b. No. 61, p. 148.

 Means against Chionaspis euonymi and other Coccida; formulæ of kerosene emulsions.
- 2404. [RILEY, C. V.] [Larval habits of Lixus.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 33.

 Lixus macer bred from stems of Chenopodium hybridum and from Helianthus; larva of L. parcus a gall producer on stems of Amelanchier.
- 2405. [RILEY, C. V.] [Girdling habits of *Pædisca obfuscata*.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 33.

 Larva of *P. obfuscata* spins a web over the orifice at the amputated end.
- 2406. [RILEY, C. V.] [Early stages of Aphorista vittata and Epipocus punctatus.] <Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 37.
 - Comparative characters between corresponding stages of the two species.

- 2407. [RILEY, C. V.] [Food-habits of Feniseca tarquinius.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 37.

 List of Aphidida preyed upon by larva of F. tarquinius.
- 2408. [RILEY, C. V.] Notes on *Phengodes* and *Zarhipis*. < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, pp. 62-63.

Structural characters of the luminous larvæ of *Phengodes* and *Zarhipis* and of an unnamed form from Nevada; difference between the perfect female and the larva.

- 2409. [RILEY, C. V.] [Trees injured by spiders.] <Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 84.

 Growth of trees retarded by spider-webs.
- 2410. [RILEY, C. V.] [Remarks on exhibited specimens.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, pp. 84-86.
 - 1. Notes on the life-habits of Ægeriidæ. Habits and early stages of Melittia gloriosa, Sciapteron robiniæ, Phemonoe 5-caudata, Ægeria impropria, A. albicornis, and A. pyri.
 - 2. Color variation in the larva of Agraulis vanillae. Colorational variation between eastern and western larvae of A. vanillae.
 - 3. Miscellaneous insects. Food-plants of Eumenia atala and Cloantha derupta; habitat of Dendrotettix quercus n. g. et sp.
- 2411. RILEY, C. V. Further notes on *Phengodes* and *Zarhipis*. < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, pp. 86–87.
 - Comparative characters of the larvæ of *Phengodes* and *Zarhipis*; life-habits and metamorphoses of the same; colorational and structural characters of the larva and larviform female of *Zarhipis*; description of the eggs of *Zarhipis*.
- 2412. [RILEY, C. V.] [Remarks on exhibited specimens.] < Proc. Ent. Soc. Wash., [13-March], 1888, v. 1, pp. 87-89.
 - 1. Notes on the eversible glands in larvæ of *Orgyia* and *Parorgyia*, with notes on the synonomy of species. Presence of glands, probably scent organs, in the larvæ of *Orgyia* and *Parorgyia*; synonomy and food-plants of some species of *Parorgyia*.
 - 2. Further remarks on *Phengodes*. Comparative characters of larva and larviform females of *Phengodes laticollis*.
 - 3. Interesting Lepidoptera. Characters of Syntomeida sp., and of an undetermined moth.
- 2413. RILEY, C. V. The British pest. Worthlessness of the sparrow as an insect-killer. <National Tribune, 26 April, 1888.
 - Result of the examination of the stomach contents of 522 sparrows, of which 92 only, or 17 fo per cent., contained insects, a large proportion of these being innoxious or actually beneficial species; review of recorded observations in North America.
- 2414. RILEY, C. V. Elm-tree depredators. <Newark [N. J.] Press and Register, 10 May, 1888.
 - Report of an address before the Newark Board of Trade; life-history and means against Galcruca xanthomelæna.
- 2415. RILEY, C. V. On the original habitat of *Icerya purchasi*. < Pacific Rural Press, 12 May, 1888, v. 35, p. 425.
 - Australia probably the true home of Icerya purchasi; its distinctness from I. sacchari.

2416. RILEY, C. V. The buffalo-gnat problem in the lower Mississippi Valley. Abstract: <Proc. Amer. Assoc. Adv. Sci. for 1887, [May], 1888, v. 36, p. 362.

Result of late investigations on species of Simulium.

2417. RILEY, C. V. Systematic relations of *Platypsyllus*, as determined by the larva. <Sci. Amer. Suppl., 2 June, 1888, v. 25, pp. 10356-10358, 4 figs.

Review of the literature concerning the systematic position of *Platypsyllus* castoris; additional facts confirming G. H. Horn's view of the coleopterous nature of the insect.

48

51

TABLE OF CONTENTS.

Damage by the chinch-bug, 48—Codling-moth, 48—Hop Aphis, 48—Work of Division in regard to silk-culture in the past year, 49—Recent experiments on the fluted or cottony cushion-scale of California, 49—Hydrocyanic acid gas and arseniuretted hydrogen, 49—Possible introduction into California of the parasites of Icerya from Australia, 50—Recent experiments on other scale-insects, 50—Professor Osborn's report on the more important insects of the season in Iowa, on the chinch-bug, on hop insects in Wisconsin, 50—Mr. Bruner's report, 50—Mr. Webster's report on the southern buffalo-gnat, 50—The year as a whole one of comparative immunity, 50—Apiculture, 50.

THE CHINCH-BUG, Blissus leucopterus. By L. O. Howard..... Introductory, 51—Past history, 51—Earliest appearance, 52—First described by Say, 52-Called "Mormon louse" in Illinois in 1840, 52-First recorded appearance in Missouri, Iowa, Indiana, and Wisconsin, 52—Its damage in the years 1864 and 1868, 52—Estimates of loss in 1871, 52-Great injury in 1874 in Missouri and adjoining States, 52-First injurious appearance in New York in 1882, 53—Damage in succeeding years, 53—Geographical distribution, 53-Indigenous east of the Rocky Mountains, north and south, 53—States in which it does the greatest damage, 53—In Cuba, 54-West of the Rocky Mountains, 54-Specimens from California and Mexico, 55—Injury during 1887, 55—Review of localities and damage this season by the statistician of the Department, 55—Table showing losses in States most damaged, 56— Summary of counties in these States, 57-Food-plants, 57-Cultivated grains and cultivated and wild grasses, 57—Rice, 58—Polygonum, 58—Stages of growth—Descriptive, 59—Egg, 59—Larval stages, 59—Pupa, 59—Imago, 59—Original description, 59—Le Baron's description, 59-Fitch's varieties, 60-Riley's variety melanosus, 60-A new variety, 60-Number of broods and hiber-

nation, 60—First accurate statement, 60—Hibernation, 60—Influence of severe cold, 61—Odor reveals hiding places, 62—Habits, 62—Flight, 62—Oviposition, 62—Young larvæ, 63—Growth, 63—Migration, 63—Habits on corn, 64—Preparations for hibernation, 64—Erroneous statement as to oviposition, 65—Exceptional habits, 65—Natural enemies and diseases, 65—Insect enemies, 65—No true internal insect parasite yet known, 65—A possible hair-

2418. RILEY, C. V.—Continued.

THE CHINCH-BUG-Continued.

worm parasite, 65-Lady-bird enemies, 65-Weeping lacewinged fly, 66-True bugs which prey upon it, 66-Inefficacy of lady-bugs when plant-lice are present, 66-Vertebrate enemies, 67-Common quail, 67-Other birds, 67-Quail laws, 67-Diseases, 68-Dr. Shimer's account of his observations on the epidemic in 1865, 68—Professor Forbes' investigations, 69—His studies of bacterial diseases of other insects, 70-Professor Riley's comments, 70-Wet weather and chinch-bug, 71—Professor Forbes' experiments, 71—Comments, 71—Wet weather and the disease, 72—Dr. Thomas' theory, 72-Professor Riley's comments, 73-An anonymous prediction, 73-Table of temperature and rain-fall in North Carolina, 74-Official records of precipitation in chinch-bug States for 1885, 1886, and 1887, 74—Remedies and preventives, 75—Earlier recommendations, 75—Preventions, 75—Clean cultivation, 75—Diversified farming, 75-Rotation of crops, 76-Early sowing and manuring, 76-Rolling, 76-Sowing an unattractive crop with wheat, 76-Direct winter remedies, 77-Burning, 77-Fall plowing and harrowing, 77.—Gas lime, 77.—Trapping, 77.—Trampling, 77-Direct summer remedies before migration, 77-Irrigation, 78—Burning, 78—Prevention of migration—Direct remedies during and after migration, 79-Ditching, 79-Tarred boards or tar alone, 80-Sowing strips of plants distasteful to the bugs around the fields to be protected, 80-Sowing strips of favored food around the fields to be protected, 80—Hot water and soap-suds, 80-Kerosene emulsion, 80-Professor Forbes' experiments, 81-Mr. Hubbard's formula, 81-Professor Atkinson's test, 81-Professor Osborn's experiments, 82-Mr. Warren's letter, 83-Bogus chinch-bugs, 83—False chinch-bug, 83—Insidious flower-bug, 84— Ash-gray leaf-bug, 84—Flea-like negro-bug, 84—Bibliographical list, 84.

THE CODLING-MOTH, Carpocapsa pomonella. By L. O. Howard..... Introductory, 88-Remarks on the bibliography, 88-Geographical distribution, 89—Date of introduction into America, 89—The insect popularly described, 89—Larva, 89—Cocoon, 89—Moth, 90— Habits and natural history, 90—Broods, 90—Its round of life, 90— Irregularity of development, 91-Number of larvæ in a single apple, 91-Larvæ of the second generation, 91-Hibernation in the larval state, 92—Apparent exception to this rule, 92—Cocoons found in apple barrels, 92-Habits of the insect in the North of Germany, 92—Food-plants, 92—Stone fruits of the Rosaceæ less infested, 92-European records of its occurrence in walnuts and oak-galls, 92-A closely allied species known to feed upon walnuts, 94—Natural enemies, 94—Birds, 94—A hair-worm parasite, 94—True hymcnopterous parasites, 94—Predaccous insects which feed upon the larvæ and pupæ, 95-Remedies, 95-The destruction of windfalls-feeding and trampling-the use of sheep and hogs, 96—Jarring or picking infested fruit from the trees, 96— Killing the moth, 97—Miss Walton's observations, 97—Mr. De Long's observations, 97—Capture of the moth with baits, 98— Insect-catching flowers, 98—Summary of the question of attracting the moth, 98-Quotations from Professor Riley's Fourth Missouri Report, 99-Attraction of moth impractical, 99-Trapping the worm-bandages, shingle traps, etc., 100-"Trimble hay-

88

2418. RILEY, C. V.—Continued.

THE CODLING-MOTH-Continued.

band system," 100—Professor Riley's rules concerning this system, 100—The Wier shingle trap, 101—Cloth bands, 101—Dr. Chapin's work in California in 18-2, 102—Spraying the trees with arsenical mixtures, 103—Professor Cook's statements, 103—Professor Forbes' conclusions from his experiments in 1885, 104—Mr. Goff's experiments at Geneva, N. Y., during 1885, 106—Supplementary conclusions by Professor Forbes, 106—Mr. Alwood's experiments in Ohio the past season, 109—Results, 110—Mr. Cushman's letter, 111—The consideration of these poisons in California, 111—Time of day for spraying, 112—Notice of Mr. Dixon's paper, 113—Apparatus for applying the arsenical mixtures, 114.

115

123

143

147

154

SILK-CULTURE—REPORT OF THE YEAR'S OPERATIONS. By Philip Walker Distribution of silk-worm eggs, 115—European egg-producers, 115—The establishment of Signor Susani, of Milan, 115—Deydier establishment at Aubenas, France, 117—Mulberry trees, 117—The Cattaneo nurseries, 117—Experimental stations, 118—The Royal Sericultural Experimental Station at Padua, 118—Experimental silk filature at Washington, 119—Production and purchase of cocoons, 120—Co-operating organizations, 122.

REPORT ON THE GAS TREATMENT FOR SCALE-INSECTS. By D. W. Coquillett

Letter of transmittal, 123—The gas treatment for scale-insects, 123—Extract from Mr. Craw's paper, 124—Results of Dr. Dimmock's experiments on various insects with pure gases, 125—The tent, 126—Apparatus for operating the tent, 126—The McMullen tent, 126—The Wolfskill fumigator, 127—The Titus fumigator, 128—The Culver fumigator, 129—The gas, 129—The dry cyanide process, 129—The dry gas process, 130—The cyanide and soda process, 131—Remarks, 132—Agitating the air in the tent, 133—Experiments, 134—Index to experiments, 142.

Report on experiments against scale-insects. By Albert Koebele Letter of submittal, 143—The value of arsenic as an addition to the kerosene emulsion, 143—Results of various experiments with resin compound, 143—Experiments with resin compound, 146—Experiments on Aphididæ with resin compound, 146.

berry saw-fly, 152—Wheat wire-worm, 153.

REPORT UPON THE INSECTS OF THE SEASON IN IOWA. By Herbert
Osborn

Letter of transmittal, 154—The turf web-worm or sod-worm, 154—Extracts from crop report, 155—Habits and life-history, 156—Description of different stages, 158—Remedies, 159—Natural enemies, 160—Other mention of the species and related forms, 160—The wheat-head army-worm, 160—Extracts from Iowa crop report, 160—Blister beetles, 161—The false chinch-bug, 162—Notes on miscellaneous insects, 162.

2418.	RILEY,	C. V	.—C	ontin	ned.
ZIIO.	TATTIELY	U. 1	•—	OHULL	$u \circ u$

Report on the season's observations in Nebraska. By Law-	
rence Bruner	164
Letter of submittal, 164—Introduction, 164—Colorado potato-	
beetle, 164—Chinch-bug, 165—Causes of increase, 165—Locusts,	
167—Cabbage insects, 168—The codling-moth, 168—Forest-tree	
insects, 168—The American Cimbex, 169—Other insects, 170.	
REPORT ON EXPERIMENTS IN APICULTURE. By N. W. McLain	170
Letter of submittal, 170 — Diseases of bees, 171 — Bacillus alvei	
(Cheshire), 174—Quotations from Professor Cheshire, 172—Treat-	
ment, 174—Starved brood, 174—Symptoms, 175—Remedy, 175—	
The control of reproduction, 175.	

U. S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

of

THE MORE IMPORTANT CONTRIBUTIONS

 $\mathbf{T}0$

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

 $B\Gamma$

SAMUEL HENSHAW.

INDEX

TO

Parts I, II, and III.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1890.



INDICES TO PARTS I, II, AND III.

SYSTEMATIC INDEX OF THE NEW NAMES PROPOSED BY B. D. WALSH AND BY WALSH AND RILEY.

[The Walsh and Riley names are followed by W. & R. The first number following the name refers to the number of the paper in the list, the second to the page where the species is first described. As is well known, the Walsh collection was destroyed in the Chicago fire in October, 1871. A few of the types sent to Drs. Hagen and Riley are in the collections of the Museum of Comparative Zoology at Cambridge and the National Museum at Washington. Those at Cambridge are indicated by a *, those at Washington by **.]

Perlidæ-Continued.

ARACHNIDA.

ACARINA.
Acaridæ.

Acarus pruni-crumena, 373; 43. s.-ænigma (Cecidomyia), 46; 608. s.-scmen (Cecidomyia), 46; 606.

MYRIAPODA.

CHILOGNATHA.

Iulidæ.

Cambala multistriatus (Iulus), 193; 34 = annulata, Say.

HEXAPODA.

PSEUDONEUROPTERA.

Psocidæ.

Psocus amabilis, 24; 362.*
bifasciatus, 39; 183.*
confluons, 39; 185.*
conterminus, 39; 185.*
geologus, 24; 362.*
lichenatus, 39; 183.*
madescens, 39; 186.*
permadidus, 39; 185.*
perplexus, 24; 361.*
purus, 24; 361.*
rufus, 39; 185.
semistriatus, 24; 361.*

Perlidæ.

Acroneura rupinsulensis, 24; 363.*
Chloroperla brunnipennis, 24; 367.*
fumipennis (Perla), 24; 366.*
nana, 24; 367.*

Perla decipiens, 24; 364.* elongata, 24; 366.*1 flavescens, 24; 363.*1 producta, 24; 365.* varians, 24; 364.* Ephemeridæ. Bætis sicca, 24; 371.*1 Bætisca, 24; 378. Cloe dubia, 24; 380.*1 ferruginea, 24; 379.* fluctuans, 21; 379.* mendax, 24; 381.* Ephemera flavcola, 24; 377.* myops, 39; 207.* Ephemerella, 24; 377. consimilis, 24; 378. excrucians, 24; 377.*1 Heptagenia, 39; 197. cruentata, 39; 205.* maculipennis, 39; 206. simplex, 39; 204.* Hexagenia, 39; 197. Palingonia flavoscens, 24; 373.*1 pulchella, 24; 375.*1 terminata, 24; 376.*1 vittigera, 24; 373.* Pentagenia, 39; 196. quadripunctata, 39; 198. Potamanthus? odonatus, 24; 372.

Siphlurus interlineata (Bætis), 39; 190=

femorata, Say.

Agrion binotatum, 24; 387.*

dentiforum, 39; 236.*

hageni, 24; 386.

Odonata.

39; 234.*1

```
Jassidæ-Continued.
Odonata-Continued.
                                                        Typhlocyba anrea, 22; 3.
   Cordulia? molesta, 39; 254.
                                                                     binotata; 22; 3.
   Gomphus amnicola, 24; 396.*2
                                                                     pallidula, 22; 3.
              consobrinus, 39; 242.*
              fluvialis, 24; 394.*1
                                                    Tingitidæ.
                                                        Gargaphia amorphæ (Tingis), 45; 409.**
              grasilinellus, 24; 394.*1
                                                                   tiliæ (Tingis), 45; 408.**
              quadricolor, 39; 246.
                                                    COLEOPTERA.
              vastus, 24; 391.*1
              ventricosus, 39; 249.
                                                    Scolytida.
   Herpetogomphus? rupinsulensis, 24; 388.
                                                        Scolytus fagi, 220; 58.
   Hetærina pseudamericana, 39; 223.*
                                                     Calandridæ.
             rupamnensis, 39; 230.*1
                                                        Sphenophorus zeæ, 337; 117** = sculptilis
             rupinsnlensis, 24; 383.
                                                           Uhler.
             scelerata, 39; 267.
                                                     Curculionidæ.
             texana, 39; 227.
                                                         Anthonomus cratægi, 197; 266.
   Lestes inæqualis, 24; 385.
                                                                      syncophanta, 197; 265.
   Macrogomphus? spiniceps, 24; 389.
                                                                      tessellata, 197; 267.
                                                         Apion languinosum, 197; 269 = walshii, Smith.
   Macromia flavipennis, 24; 398.
              illinoiensis, 24; 397.
                                                        Coccotorus prnnicida (Anthonomus), 33; 372
                                                          =scutellaris, Lec.
   Ophiogomphus mainensis, 39; 255.
                                                        Conotrachelus cratægi, 35; 37.
ORTHOPTERA.
Phasmidæ.
                                                                       puncticollis, 34; 21.
    Diapheromera velii, 45; 410.*
                                                     Chrysomelidæ.
HEMIPTERA.
                                                         Fidia viticida, 272; 87.**
Coccidæ.
                                                        Physonota 5-punctata (Cassida) W. & R.
    Chionaspis harrisii (Aspidiotus), 2; 308**=
                                                          767; 4**= unipunctata, Say.
                  furfurus, Fitch.
                                                     DIPTERA.
                salicis-nigræ (Aspidiotus), 373;
                                                     Trypetidæ.
                  40 = salicis, Linn.
                                                         Trypeta pomonella, 373; 33.**
   Pulvinaria acericola (Lecanium) W. & R., 389;
                                                     Anthomyidae.
                                                         Homalomyia leidyi, 382; 138.
                 14** = innumerabilis, Rath.
               macluræ (Lecanium) W. & R., 389;
                                                                      prunivora, 382; 138.
                 14 = innumerabilis, Rath.
                                                                      wilsoni, 382; 138.
Aphidida.
                                                     Tachinidæ.
   Aphis bella, 27; 299.*
                                                        Nemoræa militaris (Senometopia) 6; 367**=
          carduella, 27; 300.
                                                          leucaniæ Kirkp.
          quercifoliæ, 27; 298.
   Byrsocrypta pseudobyrsa, 27; 306.
                                                        Pipiza radicum W. & R., 495; 83**= 1 femo-
                 vagabunda, 27; 306.*1
                                                          ralis Loew.
    Calaphis, 27; 301.
                                                     Midaidæ.
             betulella, 27; 301.*
                                                         Midas fulvipes, 46; 306.
   Pemphigus formicarius, 27; 308.
                                                     Cecidomyidæ.
               formicetorum, 27; 308.*
                                                         Cecidomyia albovittata, 40; 621.*
               ulmi-fusus W. & R., 518; 109.*
                                                                     cornuta, 46; 625.
    Phylloxera caryæ-globuli, 27; 309.**
                                                                     cratægi-bedeguar, 376; 79.
               caryæ-semen (Dactylosphæra)
                                                                     orbitalis, 46; 623.
                 373; 23.*
                                                                     q-pilulæ (Cynips), 41; 481.
                                                                     s.-batatas, 46; 601.**1
    Schizoneura cornicola (Eriosoma?), 27; 304.
                                                                     s.-brassicoides, 46; 577.**1
                fungicola (Eriosoma?), 27; 304.
Jassidæ.
                                                                     s.-cornu, 46; 570. 197; 224.
   Chloroneura, 22; 42 = Empoasca, Walsh.
                                                                     s.-coryloides, 46; 588.*
   Empoa albicans, 22; 4.
                                                                     s.-gnaphaliodes, 46; 583.*
    Empoasca, 22; 3.
                                                                     s.-hordeoides, 46; 599.
               abnormis (Chloroneura), 22; 4.
                                                                     s.-nodulus, 46; 599.*
               consobrina, 22; 4.
                                                                     s.-rhodoides, 46; 586.*
               malefica (Chloroneura), 22; 4=
                                                                     s.-siliqua, 46; 591.*1
                                                                     s.-strobilisens, 46; 582. 197; 223.
                 viridescens, Walsh.
                                                                     s.-strobiloides, 46; 580.
               maligna (Chloronenra), 22; 4=ob-
                 tusa, Walsh.
                                                                     s.-triticoides, 46; 598.*1
               obtusa, 22; 4.
                                                                     s.-verruca, 46; 606.
               viridescens, 22; 3.
                                                                     v.-coryloides W. & R., 518; 107.**
    Erythroneura australis, 22; 4.
                                                                     v.-pomum W. & R., 518; 106.**
                  octonotata, 22: 4.
                                                        Diplosis annulipes, 46; 629.
```

ziczac, 22; 4.

atricornis, 46; 628.

¹ Also in the National Museum.

The pages refer to the separate.

Midaidæ—Continued.	Ichneumonidæ.
atrocularis, 46; 626.	Acænitus rupinsulensis, 385; 144.
decem-maculata, 46; 631.	Bassus bicapillaris, 385; 88.
helianthi-bulla, 197; 228.	ruficrus, 385; 86.
septem-maculata, 46; 631.	semifasciatus, 385; 86.
LEPIDOPTERA.	tripicticrus, 385; 85.
Tortricidæ.	Catocentrus, 385; 98.
Grapholitha prunivora (Semasia), 373; 80.**	Chorinæus cariniger (Polyrhabdus), 385; 98.
Phoxopteris fragariæ (Anchylopera), W. & R.,	Cryptocentrus, 385; 156 = Mesoleius, Holmg
499; 90.**	Cryptus albicaligatus; 385; 82.
·	albisolcatus, 385; 80.
Pyralidæ.	atricollaris, 385; 72.**
. Acrobasis nebulo (Phycita), 2; 308 = indigi-	cinctipes, 385; 74.
nella, Zeller.	nigricalceatus, 385; 77.
Geometridæ.	picticoxus, 385; 82.
Aplodes venustus (Hipparchiscus), 40; 31**=	rhomboidalis, 385; 74.
mimosaria, Guen.	rufifrons, 385; 75.
Hipparchiscus, 40; 300 = Aplodes, Guen.	Cteniscus albilineatus, 385; 107.
Bombycidæ.	ornatus (Exeuteron), 385; 105.
Halesidota antiphola, 40; 288 = tessellata, S. & A	Echthrus annulicoruis, 385; 159.
harrisii,45;430**=tessellata,S.&A.	Ephialtes gigas 385; 110.
Limacodes? tetradactylus, 40; 300.	pusio, 385; 111.
Phobetron hyalinum (Limacodes?), 40; 300.**	pygmæus, 385; 111.**
Sphingicampa, 40; 290.	Exetastes illinoiensis (Leptobatus), 385; 148
distigma, 40; 290 = bicolor,	suaveolens, 3.5; 146.
Harris.	Exochiscus, 385; 96 = Orthocentrus, Grav.
Ægeriadæ.	Exochus albiceps, 385; 96.
Ægeria hospcs (Trochilium), 197; 270.	annulicrus, 385; 95.
zingoria noopoo (zroomitam), roo, roo	atriceps, 385; 95.
HYMENOPTERA.	Glypta alboscutellaris, 385; 127.
Tenthredinidæ.	diversipes, 385; 125.
Euura perturbans, 197; 254.	ruficornis, 385; 129.
s.·gemma; 197; 250 = orbitalis, Nort.	rufipleuralis, 385; 127.
snodus, 197; 253.**	Hemiletes fuscatus, 380; 12 = var. of nema
sovum, 197; 251.	tivorus, Walsh.
Nematus fur, 197; 263.	nemativorus, 380; 11.
hospes, 197; 261.	Joppidium, 385; 67.
inquilinus, 197; 260.	ruficeps, 385; 70.
mendicus, 197; 261.	Lampronota amphimilæna, 385; 117
sdesmodioides, 197; 257.**	brevivontris, 385; 120.
spisum, 197; 258.**	imitatrix, 385; 121.
s.·pomum, 197; 255.** Pristiphora grossulariæ, 140; 123.**	interpellata, 385; 118.
	pictiventris, 385; 119.
sycophanta, 197; 263.	Mesochorus vitreus, 6; 368.**
Cynipidæ.	Orthocentrus pusillus (Exochiscus), 385; 97.
Acraspis qerinacei (Cynips), 41; 483.	stigmaticus, 385; 101.
Amphibolips pruuus (Cynips) W. & R., 518;	trifasciatus, 385; 100.
104.**	Pararhyssa, 385; 109 = Rhyssa Grav.
Ancistrophus, 821; 74.	Pezomachus minimus, 6; 368.**
l.·pisum, 821; 74.**	Pimpla cœlebs, 385; 141.
Andricus qflocci (Cynips), 41; 482.**	investigatrix, 385; 142.
qpodagræ (Cynips), 41; 491.*1	pictipes, 385; 135.
Biorhiza q. forticornis (Cynips), 41; 490.*	vidua, 385; 140.
Ceroptres ensiger (Amblynotus), 41; 496*=	Polyrhabdus, 385; 98 = Chorinæus Holmgr
petiolicola, O. S.	Polysphincta nigriceps, 385; 144.
inermis (Amblynotus), 41; 498.	nigrita, 385; 144.
Holcaspis mamma (Cynips), 518; 102.**	pimploidos, 385; 144.
Synergus albipes (Syuophrus), 41; 496*=lana,	Tryphon atricoxus, 385; 104.
· Fitch.	Braconidæ.
mendax, 41; 498.	Apanteles militaris, 6; 369.**
rhoditiformis, 41; 499* = lignicola, O. S.	Chalcididæ. Antigaster, 384; 368 = Eupelmus, Dalm.
Tribalia, 41; 470.	Decatoma dubia, 384; 300 = var. of varians
hatatorum, 41 · 471.	Walsh.

Chalcididæ—Continued.

excrucians, 384; 300 = var. of nigriceps, Walsh. hyalipennis, 384; 300. nigriceps, 384; 300. nubilistigma, 384; 300. simplicistigma, 384; 300.**

Eupelmus mirabilis (Antigaster), 384; 369.**
• Eurytoma abnormicornis, 384; 299.

a abnormicornis, 384; 299.

auriceps, 384; 299.**
bicolor, 384; 298.
diastrophi, 384; 299.
gigantea. 384; 300.
globulicola, 384; 299 = var of prunicola, Walsh.
prunicola, 384; 298.**
punctiventris, 384; 299.

Chalcididæ—Continued.

seminatrix, 384; 299 = var. of auriceps, Walsh.

Glyphe viridascens, 6; 370.**

Haltichella perpulchra (Hockeria), 11; 258.

Hockeria, 11; 258 = Haltichella, Spin.

Semiotellus chalcidephagus, 384; 369.**

Smicra albifrons (Chalcis), 6; 369.**

Formicidæ.

Formica aphidicola, 27; 310. Lasius latipes (Formica), 27; 311. Pompilidæ.

Agenia subcorticalis, 375; 162. Ceropales rufiventris, 375; 163. Sphecidæ.

Ammophila pictipennis, 375; 164.**

Bembecidæ.

Megastizus brevipennis (Stizus), 375; 162.**

SYSTEMATIC INDEX OF THE NEW NAMES PROPOSED BY C. V. RILEY.

[The types, without exception, are preserved in the collection of the National Museum, Washington, D. C.]

ARACHNIDA

ACARINA.

Trombidiida.

Tetranychus? americanus (Leptus), 1326; 17. ? irritans (Leptus), 1326; 18.

Trombidium giganteum, 1632; 143.

locustarnm, 1632; 142.

muscarum, 1632; 144.

Hydrachnidæ.

Hydrachna belostomæ, 1632; 146.

Gamasidæ.

Uropoda americana, 1626; 275.

Ixodidæ.

Ixodes bovis 1309a; 118.

Oribatidæ.

Hoplophora arctata, 1363; 216.

Tyroglyphidæ.

Tyroglyphus phylloxeræ, 1363; 215.

Acarida

Acarus aceris-crumena, 1265; 339.

HEXAPODA.

ORTHOPTERA.

Acridiidæ.

Caloptenns atlanis, 1423; 169.

Dendrotettix, 2410; 86.

quércus, 2410; 86.

Gryllidæ.

Œcanthus latipennis, 2026; 61.

HEMIPTERA.

Coccidæ.

Kermes galliformis, 1972; 482.

Mytilaspis pomicorticis, 1329; 96. = pomorum,

Bonché.

Aphididæ.

Lachnus platanicola, 2138; 198.

Pemphigus acerifolii, 1678; 16.

fraximfolii, 1678; 17.

populi-monilis, 1678; 13.

populi-ramulorum, 1678; 16.

populi-transversus, 1678; 15.

Phylloxera caryæ-avellana, 1901; 230.

caryæ-fallax, 1423; 118.

caryæ-gummosa, 1423; 118. caryæ-ren, 1423; 118.

caryæ-scissa, 1901; 230.

Schizoneura americana, 1678; 4.

ulmi (Eriosoma), 1059; 124.

Psyllidæ.

Blastophysa, 2272; 75.

celtidis-gemma, 2272; 74.

Calophya nigripennis, 2272; 69.

Ceropsylla, 2272; 76.

sideroxyli, 2272; 76.

Pachypsylla, 2208; 157: 2272; 71.

celtidis-mamma, 2208; 157. 2272;

73.

Rhinopsylla, 2272; 78.

schwarzii, 2272; 78.

Jassidæ.

Diedrocephala flaviceps, 1767; 78.

Lygæidæ.

Nysius destructor, 1329; 113. = angustatus,

Uhler.

COLEOPTERA.

Scolytidæ.

Scolytus caryæ, 938; 69. = 4-spinosns, Say.

Curculionidae.

Ampeloglypter vitis (Madarus), 1059; 132.

= sesostris, Lec.

Podapion, 2231; 62.

gallicola, 2231; 62.

Tyloderma fragariæ (Analcis), 1301; 44.

Meloidæ.

Hornia, 1601.

minutipennis, 1601.

Bruchidæ.

Bruchus fabæ, 1301; 55.

DIPTERA.

Oscinidæ.

Oscinis brassicæ, 2291; 322.

Anthomyidæ.

Anthomyia var. calopteni, 1557. = angus-

tifrons, Meig.

zeas, 1059; 155.

 $S ar cophagid \alpha.$

Sarcophaga sarraceniæ, 1390; 238.

Tachinidæ.

Exorista cecropiæ, 1112; 101.

doryphoræ (Lydella), 1059; 111.

flavicauda, 1127; 51.

Masicera archippivora, 1301; 150.

Tachina aletiæ, 1712; 162.

anonyma, 1311; 129.

Asilidæ.

Proctacanthus missouriensis (Asilus), 1127; 122 = milberti, Macq.

Simulidæ.

Simulium meridionale, 2394; 513. pecuarum, 2394; 512.

piscicidium, 1174; 228

Mycetophilidæ.

Mycetophila persica, 963; 397.

Cecidomyidæ.

Cecidomyia cupressi-ananassa, 1180; 244.

Diplosis pyrivora, 2392.

LEPIDOPTERA.

Pterophoridæ.

Platyptilus carduidactylus (Pterophorus), 1059; 180.

Tineidæ.

Blastobasis iceryæella, 2394; 485.

Coleophora malivorella, 1721; 254.

Gelechia gallæsolidaginis, 1059; 175.

Holocera glandulella, 1310; 18:

Prodoxus, 1830; 155.

ænescens, 2000; 636.

cinereus, 2000; 636.

decipiens, 1830; 155.

intermedius, 2000; 635.

marginatus, 2000; 635.

Pronuba, 1329; 150: 1336; 55.

maculata, 2000; 633.

yuccasella, 1329; 151: 1336; 56.

Tortricidæ.

Conchylis erigeronana, 1968; 316.

œnotherana, 1968; 316.

Eccopsis ferrugineana (Exartema), 1968; 317.

monetiferana (Exartema), 1968; 317.

Grapholitha gallæ-saliciana, 1968; 320.

ninana, 2176; 661.

olivaceana, 1968; 320.

Mellisopus, 1969; 322.

aurichalceana 1969; 323.

Pædisca celtisana, 1968; 319.

giganteana, 1968; 318.

Penthina fullerea.* = hebesana, Walk.

Phoxopteris cornifoliana, 1968; 324.

murtfeldtiana, 1968; 323.

Proteoteras, 1968; 321.

æsculana, 1968; 321.

Semasia helianthana, 1968; 319.

Steganoptycha claypoleana (Sericoris), 2114;

Teras cinderella (Tortrix), 1311; 47.

Pyralidæ.

Acrobasis nebulella (Phycita), 1311; 42. = var. of indiginella Zeller.

vaccinii, 2291.

Chilo oryzællus, 2119; 133.

Clydonopteron, 1929; 287.

tecomæ, 1929; 288.

Pempelia hammondi, 1311; 46.

Geometridæ.

Paleacrita, 1438; 278.

Synchlora rubivora (Aplodes), 1059; 140.

 $Noctuid \alpha.$

Acronycta betulæ, 2280; 2.

populi, 1127; 120.

Agrotis cochranii, 964; 414.

morrisoniana, 1409; 286.

scandens, 1059; 78.

Anomis texana, 2343; 350.

Exyra ridingsii (Xanthoptera), 1411; 240.

Laphygma autumnalis (Prodenia), 1301; 116.

fulvosa (Prodenia), 1301; 117.

obscura (Prodenia), 1301; 117.

Lithophane cinerea (Xylina), 1301; 135.

Plusia brassicæ, 1127; 111.

Pyrophila conspersa (Amphipyra), 1301; 74.

Bombycidæ.

Cerura multiscripta, 1411; 241.

Nola sorghiella, 2119; 187.

Ægeriadæ.

Bembecia rubi (Ægeria), 1363; 113. = margi-

nata, Harris.

HYMENOPTERA.

Cynipidæ.

Cynips q.-glandulus, 1606; 578.

q.-mellaria, 1942; 298.

Didactyum, 1749; 52 = Hexaplasta, Forst.

Hexaplasta zigzag (Didictyum), 1749; 52.

Ichneumonidæ.

Hemiteles cressonii, 1059; 177.

thyridopterygis, 1059; 150.

Ichneumon obsoletus, 1570; 55. = var. of

brevipennis, Cress. Limneria lophyri, 1570; 32.

Thersilochus conotracheli (Porizon), 1301; 28.

Braconidæ.

Apanteles acronyctæ, 1960; 312.

aletiæ, 1960; 306.

cacœcia, 1960; 305.

cassianus, 1960; 307.

flaviconchæ, 1960; 308. hemileucæ, 1960; 309.

hyphantriæ, 2394; 533.

limenitidis (Microgaster), 1301;

158.

megathymi, 1960; 304.

paleacritæ, 1960; 313.

pieridivora, 2097; 679. = var. of

congregatus, Say.

politus, 1960; 307.

rufocoxalis, 1960; 310.

scitulus, 1960; 310.

smerinthi, 1960; 311.

theclæ, 1960; 308.

Bracon charus, 1423; 75.

Exothecus prodoxi, 1831; 156.

Meteorus hyphantriæ, 2394; 532.

Microgaster gelechiæ, 1059; 178.

Microplitis ceratominæ, 1960; 303.

gortynæ, 1960; 304.

Perilitus indagator, 1311; 43.

Sigalphus rufus, 1301; 27. = var. of curcu-

lionis, Fitch.

Spathius trifasciatus, 1329; 106.

^{*}Described and figured in A. S. Fuller's "Injurious insects." <Tilton's Journ. of Hortic. and Florist's Companion, October, 1868, v. 4, pp. 207-209.

Chalcididæ.

Cirrospilus flavicinctus.1

Eurytoma bolteri, 1059; 177. = var. of diastrophi, Walsh.

Isosoma grande, 2288; 111. 2291; 358. tritici, 2060; 247.

Merisus subapterus, 2332; 416.

Pentarthron 2 = Trichogramma.

Pleurotropis phyllotretæ, 2291; 308.

Spilochalcis mariæ (Chalcis), 1112; 101.

Stictonotus isosomatis, 2119; 186.

Chalcididæ—Continued.

Tetrastichus esurus (Cirrospilus), 1712; 162. productus, 2332; 419.

Trichogramma minutum, 1301; 158. 2026; 68. pretiosa, 1712; 161.

$Proctotrupid\alpha$.

Caloptenobia, 1643; 306 = Scelio, Latr. Scelio ovivora (Caloptenobia), 1643; 306 = famelicus, Say.

Telenomus bifidus, 2378; 48. 2394; 531.

Described and figured in Lintner's First Report N. Y. State Entomologist, p. 159.

² Proposed (Record of Amer. Ent., 1871, p. 8) for Trichogramma minutum. Pentarthron has, however, been used for a genns of beetles.



INDEX.

Acrobasis oonsociella, 1311.

Ægeriidæ, 302, 576, 2341.

lifo habits, 2410.

Ægiale cofaqui = Megathymus yuccæ.

381

Ægerian maple borer. A new, 1360.

Abbot Sphinx, 1127.

Acridium amoricanum, 1413, 1549, 1570,

obscurnm, 2119.

perigrinum, 1959,

eaten out by ants, 1413.

larva, 1248, 1277. hammondi = Pempelia hammondi. Parasites on its larva, 1277. indiginella, 2, 341, 373, 511, 574, 810, 949, 975, 998, 1012, 1069, 1151, 1232, 1311, Abia caprifolia, 2360. 1580, 1962. Abnormal cocoon, 1797. juglandis, 1311. Abroad. Colorado potato-beetle, 1364. nebulella n. v., 1311=A. indiginella. Abutilon. Aletia feeding on, 2343. nebulo=A. indiginolla. Acacia filiaria. Galls on, 2176. ACACIA. INSECTS AFFECTING. vaccinii n. sp., 2291, 2349. Dorthesia characias, 1730. Acroneura rufinsulonsis n. sp., 24. Grapholitha ninana, 2176. Acronycta acericola = A. americana. Lytta cinerea = Macrobasis unicolor. americana, 571, 841. Macrobasis unicolor, 1290. betulæ n. sp., 2280. Acacias. Large white scale on, 1730. lepusculina, 1127. Academy of Science. Address of president St. oblinita, 1208, 1276, 1301. Louis, 1564, 1629. populi n. sp., 1127. Acænitus rupinsulensis n. sp., 385. xylinoides, 1329. Acanthia lectularia, 497, 1502, 1741. Acrosoma spinea, 813. Acaridous gall on Cratægus, 376. stellata, 1869. Acarina, 412, 1444, 1639, 1710, 1739, 1992. Acrydii, 1959, 2267. Acarus aceris-crumona n. sp., 1265. Actias luna, 776, 1311. pruni-crumena n. sp., 373. selene, 1311. salicis-ænigma, 46. Ad interim committees, 394. salicis-semen, 46. Adalia bipunctata, 2452. scabiei, 497. Address at Atlanta Cotton Convention, 2038. Accusation. Unjust, 1537. on entomology, 1454. Acer dasycarpum. Scale on, 1575. St. Louis Academy Science. President's, sp. Supposed tineid larva on leaf of fossil, 1564, 1629. Washington Entomological Society. Achatodes zeæ, 311. President's, 2355. Achemon Sphinx, 1127. Adelops, 2033. Acheta abbreviatus = Gryllus abbreviatus. Adimonia rufosanguinea, 156. Achetidæ == Gryllidæ. Æcidium borberidis, 1605. Achorutes nivicola, 608. Ægeria acerni, 743, 1063, 1360, 1363. Acoloithns falsarius, 124. albicornis, 2410. Acorn cnps. New oak gall on, 1606. candatum = Alcathæ candatum. gall, 1944. oxitiosa = Sannina exitiosa. moth, 1310, 1311. hospes, 197. Acraspis q.-erinacei, 41. impropria, 2385, 2410. Acrididæ, 391, 431, 433, 929, 1446, 1452, 1548, 1645, polistiformis = Sciapteron polistiformis. pyri, 2410. 1947, 1959, 2002, 2118, 2267, 2311, 2331, 2363. rubi n.sp., 1363 = Bembecia marginata. Dimorphism in, 1889. syringæ = Podosesia syringæ. Effects of winter on the eggs of, 1550. Egg parasitos of, 2355. tipuliformis, 55, 340, 481, 974, 1642, 2238.

Æschna constricta, 759.

Agassiz, L. Criticism of entomological state. ments, 44.

Agave. New use of American, 1498.

Agenia architecta, 375.

bombycina, 375, 543.

cupida, 375.

subcorticalis n. sp., 375.

Agnomia anilis, 785.

Agonoderus comma = A. pallipes.

pallipes, 3, 1825, 2317.

Agraulis vanillæ, 2410.

Agricultural advancement in the United States, 1750.

editorial excursion, 1349.

Agriculture and its advancement. Entomology and its relation to, 1329.

> American writers on insects in relation to, 2238.

> Entomological work at the Department of, 1892.

Insects in relation to, 2238.

Legislation in regard to insects injurious to, 1468.

New insects injurious to, 2055.

Agrilus ruficollis, 821, 1124, 1771.

Agrion binotatum n.sp., 24.

dentiferarum n. sp., 39.

hageni n. sp., 39.

Agrotidæ, 3, 868, 901, 964, 1504, 1522, 2310. Agrotis, 2238.

annexa, 2291.

clandestina, 901, 2291.

cochranii n. sp., 964=A. messoria, Harris.

Description of a new species of, 1409.

devastatrix=Hadena devastatrix.

fennica, 2355.

herilis, 964, 1059.

inermis = A. saucia.

lycarum, 2156.

malefida, 2291, 2322.

messoria, 901, 964, 1059, 1873, 2156, 2291, 2331, 2363.

> Harr. vs. Agrotis scandens Riley, 2156.

morrisoniana n. sp., 1409.

repentis = A. cochranii.

saucia, 657, 1059, 1095, 1775, 1826, 1941, 2291. scandens n. sp., 1059, 1408, 2156.

Riley vs. Agrotis messoria Harr., 2156.

sp., 281.

subgothica, 964, 1059.

telifera = A. ypsilon.

tricosa, 964, 2324.

ypsilon, 626, 658, 964, 1059, 1504, 2291.

AILANTHUS. INSECTS AFFECTING, 1343.

Attacus cynthia, 899, 1179, 1343, 1460, 1718.

Œta compta = Œ. punctella.

punctella, 1059, 1343.

Samia cythia = Attacus cynthia.

Ailanthus silk-worm, 899, 1311, 1718.

in Missouri, 1460.

naturalized, 1179.

Air-holes not needed in sending insects, 655.

Alabama. May beetles swarming in, 1812.

Moths caught in, 1774.

Spread of Picris rapæ into, 1720.

Alaria florida = Rhodophora florida.

Alcathæ caudatum, 481.

Alcohol. Bugs in, 605.

Aleochara anthomyia = A. nitida.

nitida, 2353.

Aleocharini, 1729.

Aletia argillacea = A. xylina.

Aletia xylina vs., 2339.

Migrations and hibernation of,

chrysalides. Not, 1826.

Chrysalides supposed to be those of, 1775.

Moths mistaken for, 1976.

xylina, 328, 636, 1127, 1338, 1353, 1363, 1369,

1649, 1657, 1689, 1702, 1712, 1719, 1721, 1722,

1728, 1736, 1749, 1758, 1769, 1814, 1826, 1831,

1845, 1852, 1882, 1888, 1903, 1914, 1932, 1953,

1976, 1997, 2038, 2064, 2069, 2102, 2119, 2130,

2141, 2155, 2164, 2295, 2343.

Aletia xylina. (See also Cotton-worm.)

Anatomy, 2343.

Annual generations, 1736.

Bibliography, 2343.

Bulletin No. 3, U.S. Ent. Com., 1763.

Characters, 1736, 2343.

Classification, 2343.

Destruction of the moth, 1763.

Destructiveness, 1763, 2343.

Devices for the destruction of, 2343.

Flight (powers of), 1763.

Habits, 2343.

Hibernation, 1763.

History of literature of, 2343.

Insects liable to be mistaken for, 2343.

Invertebrate enemies, 1763.

Machinery for the destruction of, 2343.

Mechanical means of killing, 1763. Meteorological influence upon,

2343.

Migrations, 1763.

Natural enemies, 1763, 2343.

Natural history, 2343.

Nomenclature, 2343.

in other countries, 2343.

Past history in the United States, 2343.

Poisoning the worm, 1763.

Preventive measures, 1763, 2343.

Remedies, 1763, 2343.

Stages, 1763.

Terrestrial influences, 2343.

in the United States. Hibernation

of, 2141.

Vertebrate enemies of, 1763.

vs. Aletia argillacea, 2339.

Wet weather (effects of), 1763.

Aleurodes, 221.

on Oxalis, 1791.

Allorhina nitida, 54, 718.

INDEX. 383

Alpine flowers. Fertilizers of, 1838.

Alwood, W. B. Report on Ohio insects, 2382.

Tests with insecticides on garden insects, 2382.

Alypia octomaculata, 343, 1059,1127, 1130, 1208, 1363. Amara, 1643.

Ambiguous Hippodamia, 2119.

Amblycorypha oblongifolia, 569, 1060, 1157, 1329, 1363.

Amblynotus ensiger n. sp., 41 = Ceroptres petiolicola.

inermis n. sp., 41 = Ceroptres inermis.

Amelanchier. Lixus parvus on, 2404.

America. Problem of the hop-plant louse in Europe and, 2400.

American Agave. New use of, 1498.

A. A. S. Entomology at the 1880 meeting of, 1893.

Permanent subsection of entomology, 2027.

bean Bruchus, 1301.

Cimbex, 2291, 2418.

Entomological Society, 621.

Entomologist, 1309.

Salutatory, 386.

insects. Imported and native, 1115, 1127. lackey moth, 1301.

Meromyza, 1058, 1059, 2291, 2394.

Procris, 1127.

Staphylinidæ wanted, 1786.

tent caterpillar. Eggs of the, 1329.

Ames, Iowa, Osborn, H. Experiments at, 2344.

Ammophila pictipennis n. sp., 375, 543.

Amorbia. Pine, 2363.

Amorpha fruticosa. Walshia amorphella on, 1127, 2356.

Ampeloglypter ater, 821.

sesostris, 821, 1059.

vitis = A. sesostris.

Ampelophaga myron, 837, 1056, 1086, 1127, 1247, 1290, 1960.

Ampelopsis. Swellings in roots of, 1428.

Amphibolips aciculata = A. q.-spongifica.

q.-inanis, 45, 518, 1448.

q.-prunus, 518, 1606.

q.-spongifica, 41, 45, 518, 1448.

Amphicerus, 1747.

bicaudatus, 2, 55, 249, 485, 517,633, 839, 893, 1163, 1185, 1311, 1329, 1747, 1776, 1820.

Amphipyra conspersa n. sp., 1301 == Pyrophila conspersa.

pyramidea = Pyrophila pyramidea.
pyramidoides = Pyrophila pyramidoides.

Hibernation of, 1471.

Amphiscepa bivittata, 2116.

Amputating insects, 1556.

Anabrus, 2267.

purpurascens, 1959.

simplex, 1959.

Analeis fragariæ n. sp., 1301 = Tyloderma fragariæ.

Anametis grisea, 2117.

Anaphora, 2536.

Anarsia lineatella, 2235.

Anasa tristis, 409, 825, 867, 1858, 1948.

Anatis 15-punctata, 959, 1274, 1450, 1672.

Anax junius, 1851, 2455.

Anchylopera fragariæ n. sp., 499 == Phoxopteris fragariæ.

Ancistrophus n.g., 821.

1. pisum n. sp., 821.

Anderson, E. H. Cotton-worm in south Texas in 1883, 2253.

Report on cotton insects, 2343.

Andrena, 800.

Andricus q.-californica, 1967.

q.·flocci, 41.

q.-operatola, 1352.

q.-operator, 1352.

q. podagræ, 41, 45.

q.-punctata, 45, 1822.

seminator, 739, 821, 1037.

Angoumois grain-moth, 2291.

Angular winged katydid, 1363.

Eggs of, 1518.

Animal kingdom, 393.

Aniseed and grain-weevil, 1742.

Anisodactylus baltimorensis, 627.

confusus, 2042.

Anisoplia austriaca, 1935.

Anisopteryx, 91, 96, 274, 717, 902, 910, 1391.

æscularia, 1604.

Differences between Anisopteryx pometaria and, 1604.

pometaria, 1127, 1363, 1423, 1438, 1482, 1539, 1540, 1604, 2149, 2238, 2267.

Variation in images, 1540.

Venation of wings, 1540.

vernata = Paleacrita vernata.

Anisota = Dryocampa.

rubicunda = Dryocampa rubicunda.

Anomala lucicola, 113.

Varying, 2291.

Anomis erosa, 2343.

exacta, 1651, 2343.

texana n. sp., 2343.

urena, 2119.

xylina = Aletia xylina.

Anonymous Tachina-fly, 1423, 1625.

Anophthalmus, 2033.

Ant-hill. Hemispherical larva at bottom of, 1789. Antheræa pernyi, 1311.

yama-mai, 1311.

as a silk-producer, 1346.

Anthicus, 2105.

Anthocoris insidiosus = Triphleps insidiosus.

Anthomyia, 1643.

angustifrons, 1541, 1557.

brassicæ, 2291.

Cabbage, 2291.

calopteni n. sp., 1541, 1557 = angusti-

frons, Meig.

ceparum, 225.

Egg parasite, 1570, 1625, 1643.

zeas n. sp., 657, 1059, 1065.

Anthomyidæ. North American, 1966.

Anthonomus cratægi n. sp., 197, 376, 1244.

musculus, 2322, 2331, 2363.

avenæ = Nectarophora granaria,

Anthonomus prunicida n. sp., 33 =Coccotorns Aphis, Bad work of the grain, 1638. bella n. sp., 27. scutellaris. quadrigibbus, 42, 254, 1088, 1173, 1176, brassicæ, 1365, 2291. 1228, 1301, 1358, 2232. carduella n. sp., 27. suturalis, 2363. cerasi = Myzus cerasi. sycophanta n. sp., 197, 821. Cherry, 960. tessellata n. sp., 197. Grain, 2394. Anthophora abrupta, 372, 1601, 1651. Grape-vine, 170. sponsa = A. abrupta.Hop, 1001, 2291, 2394, 2418. maidis, 23, 27. Anthreni, 483. mali, 59, 198, 201, 270, 291, 325, 413, 507, 539, Anthrenus musæorum, 492, 572. scrophulariæ, 1659, 1663, 1699, 1752. 550, 579, 591, 883, 952, 996, 1056, 1139, 1429, varius, 5772. Antigaster n.g., 384 = Eupelmus. prunifoliæ, 678. mirabilis n. sp., 384 = Eupelmus miraquercifoliæ n. sp., 27. ribis = Myzus ribis. vs. Eupelmus, 1932, 1978. Root, 990, 1038. Antiopa butterfly, 1234. rudbeckiæ = Nectarophora rudbeckiæ. Autistrophus n. g., 821, 1131. solani, 1679. vitis, 27, 102, 170. lygodesmiæ-pisum n. sp., 821.Ants, 1278, 1736, 2418. vs. rust. Grain, 1806. Acridium eaten out by, 1413. Aphodius inquinatus, 1445. and aphides, 520, 2300. Aphorista vittata, 2406. in Arizona. Leaf-stripping, 2532. gardens. Destroying black, 438. Aphrophora, 93. quadrangularis, 691. horticulture. Utilization of, 2089, 2137. Apiaries. Floating, 1785. Apiculture. McLain, N. W. Experiments in, 2394. injurious in Arizona, 1987. nests in gardens, 496. Report on experi-Red, 2309. ments in, 2363, 2418. Apidæ, 2098. Apanteles acronyctæ n. sp., 1960. Apion languinosum n. sp., 197 = A. walshii. aletiæ n. sp., 1960, 2343. cacœciæ n. sp., 1960. Apioninæ. Gall-making genus of, 2231. Apis mellifica, 13, 372, 441, 782, 833, 1455, 1774. cassianus n.sp., 1960. congregatus, 1247. Aplodes, 40. flaviconchæ n. v., 1960. mimosaria, 40. glomeratus, 2291. rubivora n. sp., 1059 = Synchlora rubihemileucæ n. sp., 2378, 2394. limenitidis n. sp., 1301.venustus = A. mimosaria. megathymi n. sp., 1960.Apple bark-lice on pears, 973. militaris n. sp., 6, 12, 1059, 1357, 1442, louse, 1552. Scurfy, 2305. paleacritæ n. sp., 1960. borer and root Aphis, 1038. remedy, 1023. pieridivora n. v., 2097. borers, 870, 1583. politus n. sp., 1960. rufocoxalis n.v., 1960. Coleophora, 1721. scitulus n. v., 1960. Curculio, 1176, 1301, 1441, 2238. smerinthi n. sp., 1960. go under ground to transform! theclæ n. sp., 1960. Does the, 1228. Apatura alicia, 1363, 1977. gall. Grape-vine, 1329, 1898. celtis, 1356, 1363. growing on a grape-vine, 403, 436. clyton, 1356, 1363. APPLE. INSECTS AFFECTING. herse = A. elyton.Acrobasis hammondi = Pempelia hammondi. lycaon = A. celtis. Amphicerus bicaudatus, 249, 485, 517, 633, 839, Apnelinus mytilaspis, 1329. 893, 1163, 1311, 1329, 1747, 1776, 1820. Aphid. Bitten by an, 2205. Anametis grisea, 2117. Aphides, 952, 1736. Anthonomus cratægi, 1244. Ants and, 520, 2300. quadrigibbus, 42, 1088, 1176, 1228, Aphididæ, 58, 198, 265, 518, 520, 1883, 2017, 2072, 2119, 1301, 2238. 2230, 2359, 2407. Aphis mali, 201, 270, 335, 507, 539, 550, 579, 591, Means against, 80, 102, 198, 508, 644, 678, 883, 996, 1139, 1429. Arctia virginica = Spilosoma virginica. 1148, 1184. of the United States. Genera of the, 27. Aspidiotus conchiformis = Mytilaspis pomo-Notes on the, 1678. harrisii = Chionaspis furfurus. Aphis, Apple borer and root, 1038, rapax, 2235.

Attacus cecropia, 139, 1028, 1076, 1215,

APPLE. INSECTS AFFECTING -- Continued.

Bostrichus bicaudatus=Amphicerus bicaudatus.

Brochymena annulata, 684.

Bucculatrix pomifoliella, 92, 1311, 1733.

Buprestis femorata = Chrysobothris femorata.

Capsus oblinoatus = Lygus pratensis.

Carpocapsa pomonella, 330, 373, 467, 523, 565, 769, 1010, 1022, 1059, 1311, 1320, 1334, 1373, 1398, 1666, 1677, 1682, 1721, 2037.

Caterpillars, 1054.

Catydid, 29.

Cecidomyia sp., 963.

Ceresa bubalus, 2195.

Chætochilus pomonellus, 874.

Chilocorus bivulnerus, 731.

Chionaspis furfurus, 69, 345, 373, 579, 951, 1061, 1152, 2305.

Chloroneura malefica = Empoasca viridescens.

maligna = Empoasca obtusa.

Chrysobothris femorata, 20, 379, 537, 555, 634, 870, 1059, 1423, 1475, 1513, 1640, 2238, 2267.

Clisiocampa americana, 66, 77, 336, 1301, 2238.

Coccus harrisii - Chionaspis furfurus.

Cœlodasys unicornis, 749.

Coleophora malivorella, 1721.

Conotrachelus nenuphar, 1214.

Corycia vestaliata, 744.

Cybocephalus californicus, 2079.

Datana ministra, 467, 922, 2222.

Diaspis harrisii = Chionaspis furfurus.

Dipteron, 548.

Elaphidion parallelum = E. villosum.

villosum, 839.

Empoasca malefica = E. viridescens.

maligna = E. obtusa.

obtusa, 22.

viridescens, 22, 30.

Epicærus imbricatus, 296.

Eriosoma pyri=Schizoneura lanigera.

Euphoria melancholica, 842.

Euryomia melancholica = Euphoria melancholica.

Galls, 552.

Gastropachna americana, 812.

velleda = Tolype velleda.

Haltica punctipennis, 2383.

Helops micans, 963.

Homipteron, 243.

Hyphantria cunea, 112, 2238.

textor = H. cunea.

Ithycerus noveboracensis, 869, 1033, 1085, 1188.

Iulus marginatus = Spirobolus marginatus.

Ixodes unipunctata, 1133.

Lagoa opercularis, 145, 796.

Larva, 758, 1054.

Lepidopteron, 243, 602.

Limacodes pithecium = Phobetron pithecium.

sp., 847.

Limenitis ursula, 145.

Lygus pratensis, 2.

Lytta cinerea = Macrobasis unicolor.

Macrobasis unicolor, 38.

Macrodactylus subspinosus, 748.

Minings, 532.

Mycetophila persicæ, 963.

25 ENT

APPLE. INSECTS AFFECTING—Continued.

Mytilaspis pomicorticis = M. pomorum.

pomorum, 69, 201, 345, 373, 423, 565, 951, 973, 982, 1059, 1061, 1329, 1552, 1772, 2238.

Notodonta concinna = Œdemasia concinna. unicornis == Cœlodasys unicornis.

Oberea sp., 1850.

Œcanthus niveus, 1691.

Œdemasia concinna, 922.

Orgyia leucostigma, 1000, 1151, 2238.

Papilio glaucus, 1048.

tarnus, 1048.

Pempelia hammondi, 810.

Pemphigus pyri = Schizoneura lanigera.

Phobetron pithecium, 1272.

Phoxopteris nubeculana, 1721.

Phycis indiginella, 511, 574, 810, 949, 975, 998, 1069, 1151.

Phycita nebulo = Phycis indiginella.

Polycaon confertus, 2103.

Prionus laticollis, 2287.

Psenocerus supernotatus, 1458.

Psocus venosus, 1263, 1526.

Reduvius raptatorius = Sinea diadema.

Saperda bivittata = S. candida.

calcarata, 963.

candida, 20, 67, 379, 421, 870, 962, 1023, 1038, 1059, 1213, 1332, 1441, 1513, 1608, 1630, 2238, 2266.

Schizoneura lanigera, 373, 467, 495, 963, 997, 1038, 1059, 1187.

Sciara mali, 211.

Sinea diadema, 638, 731.

Solenobia, 38.

Spilonota oculana = Tmetocera ocollana.

Spilosoma virginica, 296.

Spirobolus marginatus, 963.

Teras cinderella, 1311.

malivorana, 1311.

Tettigonia sp., 951.

Tineid, 77.

Tmetocera ocellana, 747.

Tolypo velleda, 972.

Tortricid, 847.

Tremex columba, 1126.

Trypeta pomonella, 177, 373, 1320, 1553, 2119, 2238.

Urocerid, 893.

Xyleborus obesus, 1583.

Apple. Lappet caterpillars on the, 972.

Large green caterpillars on the, 1076.

leaf Bucculatrix, 1311.

crumpler, 574, 949, 975, 998.

mistaken for Curculio, 1069.

folder. Tho lesser, 1311.

sewer. Chapin's, 1721.

skeletonizer, 1311, 1322.

tyer. The green, 1311.

worm, 1322.

Small, 747.

maggot, 1553, 2119, 2238.

fly, 373, 1320.

and mountain ash. Plant-louse eggs on, 579.

Apple. Oak, 1448. New York woevil on, 1085. Apple trees. Oyster-shell bark-louse of the, 1329, 2238. Scarred, 951. and peach borers, 1475, 1513. Seed ticks under bark of, 1133. pear. Two new foes of the, 22. South. Muscle-shaped bark louse pest. New, 2383. on, 1774. Pigeon Tremex in, 1126. Borer in, 839. twig. plant-louse, 883. Plnm Curculio breeds in, 1214. root-blight, 997. borer, 2287. plant-louse, 373, 495, 1059. scions. Swellings on, 548. Small reddish snout-beetle in, 1244. snout-beetle or four-humped Curculio, treo bark-lice, 345, 1059. Native, 1152. louse, 255. in, 177. borer, 1332, 1640. Flat-headed, 1059, 1423, 2238, horse chestnut. Flatworms, 523, 1010. headed, 1316. Insect foes of the, 421. New facts about the roundheaded, 1630. Oviposition of the round-Rose bug on, 748. headed, 2266. Round-headed, 1608, 1059, 2238. 2292. Variations in the two-striped Apply soap, 1367. 1213. borers, 20, 67, 379, 962, 1059. Commission, 1843. on south side of trees, 634. Aquatic larvæ, 1851. bugs, 685. Sphinx larva, 1951. caterpillar. Unicorn, 749. Arachnida, 1955. caterpillars, 66, 874, 922, 1054. Characters, 59. cocoons, 139. Poisonous, 2399. Egg-masses and cocoons on, 1000. Aragnomus griseus, 2428. Insect focs of the, 467. insects, 77, 145, 201, 276, 296, 514, 731, 1056, 1151, 1733. Archippus butterfly, 1301. Lappet caterpillar on, 812. Large moth on, 1028. worm on, 1048. leaf crumpler, 574, 1311. plant lice, 270, 335, 591, 996, 1429. virgo, 674. in Oregon, 1899. Arctic insects, 2347. Scab in apple vs., 1139. Arctiidæ, 2242. louse, 1127. Argiope riparia, 830, 1299. Argynnis, 1301. again. Eggs of, 550. Eggs of the, 507, 539. bellona, 1410. roots caused by root-lice. Knots in, Argyramæba, 1959. 1187. Arhopalus = Cylleno. Insects affecting, 963. Tent caterpillars of the, 336, 1301, 2238, 2291. tineid, 92. web-caterpillar, 112. worms, 511, 744, 810. trees. Bark lice on, 423. Eggs on, 638. Injured, 537.

Insects clustered on, 1263.

Native bark-lice on, 1061.

borers, 249, 485, 517, 633, 1163, 1311, 1329, 1747, 1776, 1820. Flat-headed, 555. Small borer in, 1458. twigs. Catydid eggs on, 29. Eggs on grape canes and, 1323. Larva boring along the axis of, 1850. Small galls and minings in, 552. vs. apple-tree plant lice. Scab in, 1139. Woolly, slug-like worm on, 796. worm, 211, 330, 376, 1022, 1059, 1666. natural history, remedies, 1373. Notes on, 1682, 1721. trap. Thomas Wier's, 1312. by machinery. Killing, 769. Severe cold and hibernating, 769. Apples and codling-moth. Michigan, 1677. Melancholy chafer in, 842. Appliances. Remedies and, 2375. Applied entomology. General truths in, 2291, Appropriations for United States Entomological Aramigus fullcri, 1708, 1721, 1740, 2117. Habits of, 1708. Pupa of, 1235. Architects. Galls and their, 518, 821. Arctia isabella = Pyrrharctia isabella. virginica = Spilosoma virginica. infaustus = Cyllene decorus. pictus = Cyllene pictus. robiniæ = Cyllene robiniæ. speciosus = Plagionotus speciosus. Aristolochia. Papilio philenor feeding on, 1774. Arizona. Ants injurious in, 1987. Arma spinosa = Podisus spinosus... Army-worm, 11, 17, 120, 670, 876, 906, 1127, 1442, 1482, 1551, 1570, 2119, 2239, 2267, 2269. Additional notes, 2001. and canker-worm, 910. Complete life-history of the, 1670.

INDEX.

387

Army-worm. Eggs of the, 1484. Asb root-borer, 1876. and its enemies, 10. saw-fly, 2291. The fall, 1256, 1267, 1282, 1301, 1482. trees in the nursery. Holes around the roots Further notes on the, 2253. Ashmead, W. H. Report on insects injurious to Further notes and observations on the, 1856. garden crops in Florida, 2388. Hibernation of, 7, 8, 2086. Ashy-gray lady-bird, 2119. Asia Minor. Bombylid larvæ destroying locust How it comes and goes, 1442. Important point yet to ascertain eggs in, 2118. in its bistory, 1442. Asilid larva, 1643. Ichneumon fly, 1127. Asilidæ, 1543. and its insect foes, 12. Asilus, 660, 668, 1127. Natural enemies, 1442. flies, 1127, 1543, 1558, 1643, 1959. bistory complete, 1551. fly. Large, 668, 1269. missouriensis n. sp., 1127 =: Proctacanthusin New York and other Eastern States. Supposed, 1990. milberti. Northern, 1835. sericeus, 668, 1543. notes and inquiries; its work in Asopia costalis, 83, 675, 948, 993, 1132, 1363, 2090. clover, 1877. farinalis, 980, 2337. in 1882, 2122. olinalis, 1363. Parasites, 6, 10, 1670. Asparagus beetle, 524. Prospects. Chinch-bug and, 2087. ASPARAGUS. INSECTS AFFECTING. question, 8, 9. Crioceris asparagi, 524, 2135. Recent outbreaks, 2262. 12-punctata, 2135. remedies, 2090. Aspidiotus, 1, 389. Supplementary notes, 1482. anrantii, 2394. True, 647. conchiformis = Mytilaspis pomorum. vs. the clover hav worm, 2090. harrisii n. sp., 2=Cbionaspis furfurus. in the West. Genuine, 2009. pinifolii - Chionaspis pinifolii. Wbeat-head, 1570, 2418. rapax, 2235. Canker-worms not, 1304. salicis-nigræ n. sp. = Chionaspis saliworms. Prof. Riley on, 1800. Three so-called, 328. Aspidoglossa subangulata, 372. What are, 1400. Aspila virescens, 1689, 2343. Arrow-weed. Carpoeapsa saltitans on, 2173. ASTER. INSECTS AFFECTING. ARTEMESIA. INSECTS AFFECTING. Diabrotica vittata, 148. Cochylis bilarana, 1059. Epicauta peunsylvanica, 38. Galls, 1347. Gortyna nitela, 75, 940. Lytta atrata = Epicauta pennsylvanica. Articulates, Yersin. Function of the nervous system of, 1959. Aster stalk borer. Dahlia and, 940. Astoma gryllarium = Trombidium locustarum. Arzama densa, 2357. obliquata, 2223. locustarum = Trombidium locustarum, Asaphes memnonius, 1874. Trombidium developed from, 2071. Aselepias. Larva on, 1535. Athysanus maritima, 2116. Asemum mæstum, 2267. Atlanta cotton convention address, 2038. Asexual females. Lichtenstein's theory as to di-Atomizers. Sprinklers and, 1857. morphic, 2072. Atlantic migratory locust, 1423, 1625. Ash. Borers in black, 1733. Atta fervens, 1964, 2532. Cocoons on the flowering, 950. Attacus angulifera, 1311. colored locust, 2363. cecropia, 139, 209, 238, 277, 287, 297, 542, 629, destroyers. Linden and, 1847. 635, 786, 832, 841, 911, 1028, 1029, 1076, gray blister beetle, 1209, 1558. 1112, 1215, 1249, 1311, 1533. leaf-bug, 1127, 1423, 2418. cynthia, 542, 899, 1179, 1311, 1343, 1460, 1718, pinion, 1301. ASH. INSECTS AFFECTING. luna = Actias luna. Attacus promethea, 950. polyphemus = Telea polyphemus. Fatua denudatum, 1733. promethea, 259, 268, 604, 629, 950, 1311, 1486, Hypbantria cunea, 1733. 1797, 2204. textor = H. cunea. ricini, 1718. Neoclytus capræa? 1849. Attagenus injuring featbers, 56. Parandra brunnea, 1733, 1876. megatoma, 2113. Trochilium denudatum = Fatua denudatum. Attractive but untrue, 1644. Xyloryctes satyrus, 471. Attus, 2302.

Aulax, 318.

Austrian pines. False caterpillars on Scotch and,

Ash larva. Prickly, 1914.

and mountain ash, 705.

Plant-louse eggs on apple and mountain, 579.

BARLEY. INSECTS AFFECTING. Azalea. Beetles on, 156. Laehnosterna fusca, 2394. Prasocuris varipes on, 156. Bacillus alvei, 2418. Nectarophora granaria, 2394. Schizoneura sp., 2394. Back-rolling wonder, 1363. Bacterial disease of the imported cabbage-worm Siphonophora avenæ = Nectarophora granaria. Barley-root louse, 2394. F. M. Webster. Insects affecting, 2394. Bacterium, 2251. Bad packing, 837. Barnard, W.S. Machinery for destroying Aletia, Bætis femorata = Silphlurus femoratus. 2343. interlineata n. sp., 39 = Silphlurus femora-Tests of machinery for destroying the cotton-worm, 2253. sicca n. sp., 24. Basket-worm, 271, 1059, 1090, 1650. Bietisca n. g., 24. Bass. Insects from stomach of rock, 1793. obesa. Pupa of, 43. Larvæ in stomach of black, 1792. Bag-worm, 1059, 1090, 1189, 1650, 2378. Bassus bicapillaris n. sp., 385. New, 1441. ruficrus n. sp., 385. at South Pass, Ill., 1156. semifasciatus n. sp., 385. Thyridopteryx ephemeræformis. Retripictierus n. sp., 385. marks on, 2272. Bass-wood. Dipteron on, 548. worms, 182, 266, 538, 641, 738, 1036. Batrachedra salicipomonella, 821. again, 815. Bazille, Louis. Obituary, 2370. and borers, 1472. Be on the guard, 708. Ways of, 1424. BEAN. INSECTS AFFECTING. Bahia, Brazil. Cotton culture and the insects af-Bruchus fabæ, 1301, 1631, 2127, 2238. granarius, 1120. feeting the plant at, 1788a. Bailey, J. S. Tree borers of the family Cossidæ, obsoletus, 1155, 1232, 1296. 2253. Epilachna corrupta, 2235. Balaninus rectus, 1310, 1311. Lytta cinerea = Macrobasis unicolor. Balsam fir. Lachnus strobi on, 1039. Macrobasis unicolor, 38. Bamboo Sinoxylon, 2454. Bean. A new enemy to, 2418. weevil, 1155, 1296, 2127, 2238. Banded Ips in calyx of pear, 1239. Baridius sesostris = Ampeloglypter sesostris. American, 1301. trinotata = Trichobaris trinotata. Beans. Buggy, 1631. Bark beetles, 295. Beautiful wood nymph, 1127, 1363. borer. Hickory, 938, 1329. Bed bugs, 1502. Peach-tree, 2047. Lepidium vs., 1741. borers, 220. Bedeguar of the rose, 1939. boring beetles. Packard on the develop-Bee bread devoured by worms, 1293. ment of, 2267. Bug preying on honey, 13. lice, 1, 264, 935, 944, 989, 994, 1007. disease. Possible cause of the, 704. again, 1040. enemy, 1186. on apple-trees, 423, 1059. fly, 1959. Native, 1061. killer. Nebraska, 1127, 1311. Curculio and, 487. killers, 1543. eggs in Missouri. Supposed, 1084. Bcetle named, 627. on grape-vine, 1212. Remarkable felting caused by a, 2113. Maple, 1004. in Russia. Excessive injury by, 1935. Means against, 104. on sugar maple, 1014. Native apple-tree, 1152. in thistle, 169. on the pear, 982. Beetles, 1736. pears. Apple, 973. around peach-trees, 751. Preventive of, 1012. Food-habits of ground, 1760. remedy, 896. in dried English currants, 1260. · on rose bushes, 1303. in flouring-mills, 1261. louse, 987, 1397, 1583. injuring cabbages and fuchsias, 1868. Apple, 1552. in stomach of meadow lark, 1013. on apple - trees South. in the corn-field, 2317. Muscleshaped, 1774. Mites parasitic on, 315. enemy, 154. named, 732, 802, 806, 1172, 1187. Imported apple-trec, 255. on buttercup and azalea, 156. Insect focs of the, 417. Plant-feeding habits of predaceous, 1956. in Missouri. Oyster-shell, 1169. supposed to be feeding on wheat, 1746. on Osage orange, 389. swarming about the lawn, 718.

Oyster-shell apple, 1127, 1329, 2238.

Scurfy apple, 2305. Tulip-tree, 271.

under dead fish, 817.

working in wheat, oats, and rye, 1259.

Belated individuals of the periodical Cicada, 648.

Belfrage, G. W. Obituary, 2161.

Belostoma americanum, 473, 534, 733, 1570, 1584.

grande = B. americanum.

Belvoisia bifasciata, 1329, 2343.

Bembecia marginata, 576, 1363, 1454.

rubi = B. marginata.

Bembex fasciata, 543.

marginata, 2230.

Bembidium 4-maculatum, 3.

Bennett, T. Report of experiments at Trenton, N.J., 2344.

Berberry. Aphididæ on, 644.

Berne. Phylloxera convontion at, 2147.

Berry and cherry twigs, 1503.

Beschreihung einer den Birnen schädlichen Gallmücke, 2392.

Best's fruit-tree invigorator again, 530.

invigorator once more, 545.

Betula nigra. Acronycta on, 2280.

Bibio alhipennis, 257, 683.

Bibionidæ, 2076.

Bihliography of gall literature, 2059.

Bidens chrysanthemoides. Syrphus sp. caught by, 1761.

Bill bug. Corn, 2291, 2418.

Biorhiza forticornis, 41, 745.

Birch. Clytus sp. ? on, 2267.

Bird question dispassionately considered, 1768.

Birds. Professor Riley in favor of the, 1675.

Relation of Carahidæ and Coccinellidæ to, 2152.

vs. insects, 199, 369.

Bittacus apterus. Hahits of, 2092.

Bitten by an aphid, 2205.

Bitter-sweet. Enchenopa binotata on, 1913.

White waxy secretion on stems of, 1913.

Black ants in gardens. Destroying, 438.

ash. Borers in, 1733.

bass. Larva in stomach of, 1792.

blister beetle, 1059, 1558.

beetles on potatoes, 1381.

breeze fly, 1127.

bug, 314.

currant. Currant-worms and, 1079.

grape-vine caterpillars, 1018.

knot onco more, 930.

legged tortoise beetle, 1127.

potato heetles. Large, 1206.

rat blister beetle, 1059, 1558.

walnut. Insects feeding on sap of, 1195. and willow. Enemies of the, 2333.

Blackberry borers, 302.

cane borers, 576.

gall insects, 318.

Pithy, 1131.

Galls on, 263.

BLACKBERRY. INSECTS AFFECTING.

Ægeria rubi = Bembecia marginata.

sp., 302.

Agrilus ruficollis, 1771.

Bembecia marginata, 576, 1364, 1454.

Diastrophus cuscutæformis, 609.

nebulosus, 263, 610, 1131. sp., 318.

Eudemis botrana, 1232,

BLACKBERRY. INSECTS AFFECTING—Continued.

Lecanium sp., 117.

Lobesia botrana = Eudemis botrana.

Oherea bimaculata, 302, 783, **1363**.

perspicillata = 0. bimaculata.

tripunctata, 1503.

Œcanthus niveus, 1503.

Psylla rubi = Trioza tripunctata.

Trioza tripunctata, 669.

Blackberry. New insect foe of the, 669.

and raspberry canes. Gouty galls on, 1771.

scale, 117.

twig. Fuzzy galls on, 609. Pithy galls on, 610.

Blastobasis iceryæella n. sp., 2394.

Blastophysa n. sg., 2272.

celtidis-gemma n. sp., 2272.

Blatta, 2267.

germanica = Ectobia germanica.

Blattidæ, 573, 1858.

Blennocampa pygmæa, 1056, 1252.

Blepharida rhois, 849, 1363.

Blepharoceridæ, 1981, 2004.

Blight. Apple-root, 997.

Blind-cyed Smerinthus, 1912.

Going it, 588.

as those who shut their eyes. Nonc so, 275.

Blissus lencopterus, 6, 90, 180, 398, 548, 886, 888, 890, 891, 894, 895, 1068, 1127, 1357, 1402, 1423, 1432, 1453, 1508, 1538, 1681, 2018, 2039, 2048, 2087, 2119, 2143, 2237, 2252, 2271, 2418.

Blister beetle. Ash-gray, 1059, 1209, 1558.

Black, 1059, 1558.

rat, 1059, 1558.

Margined, 1059, 1558. on potatoes. Black, 1381.

Retarded dovelopment in, 1860.

Striped, 1059, 1558.

beetles, 134, 289, 301, 362, 1059, 1558, 2238, 2418.

Larval characters and habits of, 1600.

from New Mexico, 1796.

Notes on life-history of, 1651.

Pear leaf, 1739.

Blcod-red lady-hird, 2119.

sucker and pear slug, 1253.

sucking cone-nose, 795, 854.

Conorhinus, 2291.

Blow-fly. Cheese-fly and, 1266.

Bluebird. Ichneumonid from stomach of, 1878. Larvæ from stomach of, 1871.

Bluebirds feeding on parasitic and predaceous insects, 1885.

Blue caterpillars of the vine, 1127, 1363.

spangled peach-worm, 1301.

Bogus chinch-hug, 1127, 1423.

Colorado potato-beetle, 1558.

bug, 519.

Thrips. True Thrips and, 203.

Yucca moth. True and, 1804.

Boll-worm, 821, 1301, 1353, 2119, 2164, 2343.

Report on, 2164.

worms feeding on cabbage, 1915.

Bombardier beetle, Kansas, 1311.

Bombus, 800, 1032, 1226.

Hibernation of the queens of, 1032. pennsylvanicus, 800.

virginicus, 1774.

Bombycidæ, 1718.

Bombylid larvæ destroying locust eggs in Asia Minor, 2118.

Bombylii, 1959.

Bombyliidæ, 2118.

Larval habits, 1947, 2169.

Life-habits, 2002.

Natural history of certain, 1928.

Bombylius major, 1947, 1959.

Bombyx arrindria, 899.

mori = Sericaria mori.

Bordered soldier bug, 1059.

Borer, 47.

Borers, 55, 141, 233, 967, 1083.

Bag-worms and, 1472.

Brimstone for, 931.

Canker-worms and, 941.

Tansy for, 1016.

Boston. Entomologists at, 1894.

Bostrichus = Amphicerus.

bicaudatus = Amphicerus bicaudatus.

Botis. Cauliflower, 2232.

repctitalis, 2232.

Bot flies, 114.

fly. Cattle, 2238.

Horse, 2238.

Sheep, 2238.

Squirrel, 526.

Botrytis rileyi, 2232.

Bots, 1563.

Bowels. Larvæ in human, 382.

Box Psylla, 2291.

Brachinus, 802.

americanus, 802.

Brachyrhynchus, 553.

Brachytarsus variegatus, 18, 1461, 1479.

Bracon charus n. sp., 1423.

Cherished, 1423.

scolytivorus n. sp., 1329.

Bradynotes opimus n. sp., 1959.

Brandt on nervous system of insects, 2225.

Branner, J. C. Cotton-caterpillars in Brazil, 2343.

Preliminary report of observations upon insects injurious to cotton, orange, and sugar-cane in Brazil,

2277.

Brauer on Hirmoneura. Review of, 2275.

Braula cœca not particularly injurious to the honey-bee, 1982.

Brazil. Branner, J. C., on cotton-caterpillars in, 2343.

> insects injurious to cotton, orange, and sugar-caue in, 2277.

Lignified snake of, 2136.

Breakfast. Entomological, 2311.

Breeding-cages, 695.

silk-worms, 1648.

Breeze-fly. Black, 1127.

Brenthian. Northern, 1363.

Brimstone for borers, 931.

British pest, 2413.

Broad-necked Prionus. Eggs of, 1329.

winged katydid, 1329, 1363.

Brochymena annulata, 684.

Brown mantispian, 1243.

Brown's answer to inquiries about Colorado potato-beetle. Mayor, 1491.

Bruchus of Europe just imported. Grain, 1120.

fabæ n. sp., 1301, 1631, 2127, 2238.

Grain, 1301.

granarius, 1120, 1301.

obsoletus, 1155, 1232, 1296.

pisi, 434, 1120, 1301, 1357, 1916, 2238.

Bruner, L. Additions to the chronology of locust

ravages in 1880-'81, 2267.
List of North American locusts, 2267.

Notes on other locusts and the western cricket, 2267.

from Nebraska, 2291.

Observations in the Northwest on the Rocky Mountain lo-

angt 2165

cust, 2165.
on the Rocky Mountain

locust and other insects in the Northwest during the summer of 1883, 2277.

Report on the abundance of the Rocky Mountain locust

in 1885, 2363.

locusts in Texas in spring of 1886, 2382.

Nebraska insects, 2382.

of the season's observations

in Nebraska, 2418.

on the Rocky Mountain locust in Montana in 1880, 2267.

> the Rocky Mountain locust in Wyoming and Montana, etc., in 1881, 2267.

Bucculatrix. Apple-leaf, 1311.

pomifoliella, 92, 1311, 1733.

thuiella, 1311.

Buck fly, 595.

moth, 1329.

Buckeye leaf. Sericoris claypoleana on, 2114. stem-borer, 2114.

Buckwheat to destroy insects. Use of, 1744.

Webster on insects affecting, 2394.

Bud-eating insects, 1408.

Buffalo. Eutomology in, 2035.

gnat problem in the Lower Mississippi

Valley, 2416.

Southern, 2291, 2394, 2418.

gnats, 2394.

Webster on, 2388.

tree hopper, 415, 1329.

injurious to potatoes, 2107.

Bug. A most precious, 575.

Buggy beans, 1631.

peas, 1916.

Bug-hunter in Egypt, 372.

hunters. Cruel, 440.

Bugs, 354.

in alcohol, 605.

Bugs. Hogs vs., 387. Butterfly larva injurious to cotton squares, 1872. Our, 2374. pino-trees, 2121. Toads vs., 1103. larvæ. Summer dormancy of, 1410. Buhach. Coquillett on the production and manunamed, 666. facture of, 2364. tongue, 1815. Bulletin Div. Ent. U. S. Dept. Agric., No. 1, 2164. trees of Montercy again, 2052. 2, 2165, wings. Chemical change in color of, 1935. 3, 2253. Byrsocrypta pseudobyrsa n. sp., 27. 4, 2277. vagabunda n. sp., 27. 6, 2304. Cabbage Anthomyia, 2291. 8, 2315. bug. Harlequin, 1099, 1311, 1534, 2238, 2263, 9, 1724. 2291. 10, 2378. butterfly. Imported, 1127, 2291. 11, 2344. Larger, 2232. 12, 2364. Southern, 1127, 2232, 2238. 13, 2382. cut-worms, 2291. 14, 2388. Remedies for, 2291. 15, 2389. flea-beetle. Colorado, 2291. Michigan Agrie. College, No. 39 (Reinsects, 1297, 2291, 2418. view), 2501. CABBAGE. INSECTS AFFECTING. United States Entomological Commis-Agrotis annexa, 2291. sion, No. 1, 1577. clandestina, 2291. malefida, 2291. 2, 1578. messoria, 2291. 3, 1736. 6, 2026. saucia, 2291. tolifera = A. ypsilon.Bumble-bees vs. red-clover, 2376. What becomes of, 1032. ypsilon, 626, 2291. Bundle of entomological queries, 556. Anthomyia brassicæ, 2291. Buprestidæ, 821, 2267. Aphis brassicæ, 1365, 2291. Buprestis divaricata = Dicerea divaricata. Botis repetitalis, 2232. femorata = Chrysobothris femorata. Caloptenus spretus, 2291. Burr oak. Insects stripping the, 966. Doryphora 10-lineata, 2291. Burgess, E. Anatomy of Aletia, 2343. Elaterid, 740. Burying beetles, 315, 845. Epicærus imbricatus, 1868. Hadena devastatrix, 2291. Bushels of butterflies, 238. Butalis cerealella, 1232. Heliothis armigera, 1915. Buteo pennsylvanicus, 643. Lygus lineolaris = L. pratensis. pratensis, 2291. Buthus carolinianus, 453, 1119. Butler, A. W. The periodical Cicada in south-Mamestra chenopodi = M. trifolii. picta, 1127, 2232. eastern Indiana, 2364. Buttercup. Beetles on, 156. subjuncta, 2291. INSECTS AFFECTING. trifolii, 2232. Adimonia rufosanguinea, 156. Murgantia histrionica, 1097, 1297, 1311, 1534, Galeruca rufosanguinea = Adimonia rufosan-2238, 2263, 2291. Nysius angustatus, 2291. guinea. Oscinis, 2291. Butterflies. Bushels of, 238. caught by the tongne, 1761. Phyllotreta albionica, 2291. Controlling sex in, 1352. vittata, 2291. Effects of cold applied to the chryszimmermani, 2291. alids of, 1778. Pieris monuste, 2232. Hackberry, 1356, 1363. oleracea, 835, 1127, 2232, 2238. protodice, 1127, 2232, 2238. Imitative, 613. Migrations, 1770, 1991. rapæ, 1127, 1257, 1279, 1294, 1694, 1721, Migratory, 1622, 1635. 2221, 2232, 2238, 2251, 2298. Pionea rimosalis, 1915, 2232. named, 461. of New England. Notice of Scud-Plusia brassieæ, 1127, 2232, 2238. Plutella cruciferarum, 1291, 1297, 2232. der's, 2464. North America. Review of Ed-Strachia histrionica = Murgantia histrionica. ward's, 395, 479, 623, 827. Cabbage-lice, 1365. Odor in, 2025. Mamestra, 2232. Philosophy of the pupation of, 1711. Oscinis, 2291. Swarm of, 406, 1050. pest. Our now, 1257. Butterfly chrysalis, 1535. pests, 740.

Pionea, 2232.

plant-louse, 2291.

Goat-weed, 1117, 1127, 1329.

larva. Carnivorous, 2359.

392 INDEX.

Callosamia angulifera = Attacus angulifera.

Cabbage plants. Cut-worms severing, 626. Callosamia promethea = Attacus promethea. Callostoma fascipennis, 2075, 2118. Plusia, 1127, 2232, 2238. Calocoris chenopodi, 2458. Plutella, 1291, 2232. rapidus, 2363. worm. Bacterial disease of the imported, Calomicrns. Characters of, 1799. 2251. Calophya nigripennis n. sp., 2272.Imported, 2232, 2238, 2291. A new remedy for the imported, Caloptenobia n.g., 1643 = Scelio. ovivora n. sp., 1643 = Scelio fameliparasite, 2221. in the South. The imported, 1714. Caloptenus, 2267. Successful introduction of a paraatlanis n. sp., 1423, 1549, 1643, 2198, 2232, site of the imported, 2291. The westward progress of the im-Packard on the development of, ported, 1694, 1721. Zebra, 2232. bivittatus, 1959. devastator, 2323, 2377. worms, 835, 1127, 1279, 1915, 2232. differentialis, 1423, 1549, 2323. upon gillyflowers, 614. Remedy for, 1920, 2232. Epidemic disease of, 2240 Dimorphism in, 1889. Safe remedies for, 1948. femur-rubrum, 147, 475, 1423, 1437, 1451, Cabbages and fuchias. Beetles injuring, 1868. Cabinet pests. Means against, 1057. 1549, 1598, 1643, 2471. Cacœcia fractivittana, 2475. Mite infesting, 146. rileyana, 1059, 1192. spretus, 147, 368, 373, 475, 536, 926, 929, 1423, 1433, 1434, 1437, 1451, rosaceana, 735. 1452, 1456, 1473, 1481, 1482, v-marked, 2363. 1488, 1492, 1493, 1494, 1495, Cactus lady-bird, 2119. 1501, 1521, 1527, 1532, 1538, Caddis-worms. Carnivorous habits of, 1839. Calandra granaria, 510, 586, 1259, 1673, 1683, 1742. 1541, 1549, 1555, 1557, 1562, 1564, 1565, 1566, 1567, 1570, oryzæ, 510, 1673, 1683. 1571, 1572, 1573, 1575, 1576, remotepunctata, 15. 1577, 1578, 1582, 1585, 1588, Calaphis n. g., 27.1596, 1598, 1612, 1613, 1614, betulella n. sp., 27.Calathus gregarius, 2418. 1615, 1617, 1643, 1669, 1674, 1693, 1959, 1994, 1998, 2003, California. Chinch-bug, 2431. 2005, 2007, 2040, 2044, 2057, Fruit insects in, 2245. 2267, 2276, 2291, 2335, 2366. Fuller's rose beetle in, 1740. (See also Locust; Rocky Grape Phylloxera in, 1727. Grasshopper ravages in, 2323. Mountain locust.) Agricultural bearingsof, 1643. Injurious insects in, 2074. Aid of animals against, 1643. Interest in economic entomology in, Air-sacks of, 1959. Anatomy of, 1643. Introduction of Lestophonus iceryæ Animals as foes to, 1643. into, 2522. Locust ravages in, 1959. Autumn flights in 1877, 1643. Birds that feed on, 1643. migratory locust, 2363. Brain of, 1959. Overflow bug in, 2099. Breeding-grounds of, 1643. Phylloxera in, 2140. Promotion of silk-culture in, 2006. British-American data, 1643. red scale, 2394. Characters, 1625. Winged Phylloxera in, 1895. Chronological history, 1625. Californian locust, 1959. Chronology of ravages, 1643, orange insects, 2373. Callidium antennatum, 50, 319. Classification, 1643. Colorado data for 1877, 1643. janthinum, 50, 319. Crops as affected by, 1643. Callidryas philea, 1271. Secondary sexual characters of, 1784. Dakota data, 1643. Callimorpha fulvicosta = C. lecontei. Destruction (devices for), 1643. lecontei, 1301. (means of), 1643. Development of, 2267. sp., 1871. vestalis = C. lecontei. as influenced Calliphora, 1390. by meteorovomitoria, 1266. logical condi-Callipterus, 1678. tions, 1959. Callochlora viridis = Parrasa chloris. Devices for destruction of,

1643.

Caloptenus spretus. Distribution of, 1625, 1643, Caloptenus spretus. Remedies, 1643. Statistics of losses caused (southern limby, 1643. its of), 1959. in the temporary region af-Effects that follow severe inter the departnre of the rejury, 1643. turn swarms, 1643. Eggs, 1643. Texas data, 1957. in same ground for two for 1877, 1643. consecutive years, Transformations, 1625. 1643. travel. Direction in which Enemies (invertebrate), 1643. young, 1643. (natural), 1625, 1959. Uses of, 1643. (vertebrate), 1643. Utah data, 1643. Fires (influence of prairie), Vertebrate enemies of, 1643. 1648. Visitation of, 1643. Flight, 1959. Weather (effects of), 1643. (powers of), 1959. Wings (time of year when Flights in 1877 (autumn), 1643. acquired), 1643. from 1877 to 1879, 1959. Young in country where General considerations, 1625. they hatch, 1625. Calopteron and Photinus. Food of, 2358. Habits, 1625, 1643. (at night), 1643. reticulatum, 806. Histology, 1643, 1959. terminale, 806, 2358. Injury (effects that follow Calosoma calidum, 721, 1274, 1314, 1357, 1672. severe), 1643. scrutator, 1226, 1898. Invertebrate enemies of, 1643. Calves. Lice on, 889. Iowa data, 1643. Cambala annulata, 193, 236. for 1877, 1643. multistriatus=C. annulata. Camel cricket, 1059. Kansas data for 1877, 1643. Losses caused by, 1643. Campula atrox, 1959. pellucida, 1959, 2044, 2313, 2363. Means against, 1959. Metamorphoses, 1643. Campoplex fugitiva = Limneria fugitiva. Meteorological conditions (in-Camptonotus scudderi, 2276. fluence of, on migrations), Campyloneura vitripennis, 1301. Canadian Entomologist (Review), 395, 479, 827. Migrations, 1643, 1959. Canker-worm, 86, 172, 1021, 1066, 1127, 1363, 2012. (influence of meagain, 173. teorological Army-worm and, 910. conditions on), Cure for, 1426. 1959. Fall, 1423, 1482, 2238. in 1877, 1643. Howell, A. J. Experience with Minnesota data, 1643. the Spring, 2267. for 1877, 1643. once more, 274. Missouri data for 1877, 1643. parasite, 717. Native home, 1625. Spring, 1423, 1482, 2238. Natural enemies, 1625, 1959. worms, 91, 96, 902, 1482, 2267, 2270. Natural history, 1625, 1643. not army-worms, 1304. Nebraska data, 1643. Borers and, 941. Concerning, 2149. for 1877, 1643. New Mexico data, 1643. Remarks on, 1438. Night habits of, 1643. at the West, 1539. Nomenclature, 1643. Canna flavida. Pamphila ethlius feeding on, 1897. Parasites on eggs, 1612. Hesperid larva feeding on, 1897. Cannibal mites, 624. Permanent breeding-grounds Cantharis cinerea=Macrobasis unicolor. of, 1643. Powers of injury, 1625. vesicatoria, 2118. Cauthon chalcites, 1172. Practical considerations, 1625. Prairie fires and, 1643. hudsonias == C. lævis. lævis, 1172, 1386. Previous visitations of, 1643. Ravages of (chronology of), Cape. Grape Phylloxera not at the, 1841. Capitalizing specific names, 2170, 2257. 1643, 1959. Capnia minima, 1141. (east of the Mis-

sissippi), 1625.

(relative to agriculture), 1959.

Capsus oblineatus = Lygus pratensis.

Carabidæ, 1537, 1726, 1738, 1760, 1956.

4. vittatus = Pecilocapsus lineatus

Carabidæ and Coccinellidæ to birds. Relations Cecidomyia enpressi-ananassa n. sp., 1180. of, 2152. destructor, 6, 62, 122, 150, 157, 250, 426, Food relations of, 494, 711, 1506, 1581, 1587, 1665, 1690, 2151. 1721, 1787, 2008, 2031, 2235, 2267, 2331, Curculio-feeding habits of, 372. 2365, 2395, 2398. Horn's classification of, 2051. impatientis, 852. Mode of oviposition of, 2341, 2355. leguminicola, 1690, 1721, 2238. Care of entomological specimens. On the, 2290. orbitalis n. sp., 46. Carnivorous. Are honey-bees, 2098. oryzæ, 1949. butterfly larva, 2359, 2361. q.-pilulæ n. sp., 41, 799. habits of caddis-worms, 1839. q.-symetrica, 799. propensities of plant-feeders, 1867. s.-ænigma n. sp., 46 = Acarus s.-ænig-Carolina sphinx, 1262. Carpet-beetle. The imported, 1699. s.-batatas n. sp., 46. Trapping the, 1752. s.-brassicoides n. sp., 46, 518.bug, 1659. s.-cornu n. sp., 46, 197. pests, 1663. s.-coryloides n. sp., 46, 518. Carpets in Texas. A tineid on, 2531. s.-gnaphalioides n. sp., 46. Carpocapsa, 934. s.-hordeoides n. sp., 46. pomonella 57, 307, 330, 367, 373, 377, s.-nodulns n. sp., 46. 387, 454, 467, 523, 565, 769, 977, 1010, s.-rhodoides n. sp., 46. 1022, 1056, 1059, 1254, 1301, 1311, 1320, s.-semen n. sp., 46 = A carus s.-semen.1329, 1334, 1363, 1373, 1394, 1398, 1476, s.-siliqua n. sp., 46, 1170. 1496, 1609, 1666, 1695, 1721, 1887, 2037, s.-strobiliseus n. sp., 46, 197.2230, 2261, 2293, 2418. s.-strobiloides n. sp., 46, 518.saltitans, 1496, 1609, 2163, 2173. s.-triticoides n. sp., 46. Carrying out the law, 1993. s.-verruca n. sp., 46. Carteria laccæ, 2119. secalina, 2267. larreæ, 2119. solidaginis, 798. mexicana, 2119. sp., 799, 963. Case-bearer. Dominican, 1363. tritici = Diplosis tritici. Lepidopterous, 260. tubicola, 1232. Walnut, 1311. v.-coryloides n. sp., 518, 1329. v.-lituus = C. v.-viticola.Cassida, 94, 1238. aurichalcea = Coptocycla aurichalcea. v.-pomum n. sp., 403, 518, 1284, 1329, bivittata, 681, 993, 1127, 1510. 1898, 2136. elavata = Coptocycla elavata. v.-viticola, 791, 1077, 1116, 1329. nigripes, 1127, 1510, 1697. Cecidomyid gall on Cratægus, 376. pallida = Coptocycla aurichalcea. in wheat, 18. 5-punctata n. sp., 767 = Physonota uni-Cecidomyidæ, 294, 518, 799. punctata. on grape. Galls of, 111. texana, 2096, 2215, 2235. salix. Galls of, 46, 197. Cassididæ, 105, 767, 973, 1056, 1082, 1510. new species, 46. Castor bean. Locusts eat the, 1645. Cecil's books of natural history (Review), 506. Catalogues and monographs, 1765. Cecropia Cryptus, 1301. Catalogus coleopterorum, 1866. moth, 1112. Catalpa sphinx, 2119, 2291. caterpillars, 786. Cat-bird. Tipula eggs in stomach of the, 1735, cocoon, 635, silk-worm, 1311. Catcher. Dr. Hull's Curculio, 651. Remarkable abundance of, Caterpillar nuisance, 1995. Caterpillars named, 857, 1276. Tachina fly, 1311. Stinging, 1748. worm, 1215, 1249. Catocentrus n.g., 385. on elder, 1533. Cattle bot-fly, 2238. CEDAR. INSECTS AFFECTING. Catydid eggs, 29, 247. Callidium antennatum, 319. as an enemy to Doryphora 10-lineata, 313. janthinum, 319. Catydids. Food of, 143, 241. Cerambycid, 1324. Cauliflower Botis, 2232. Thyridopteryx ephemeræformis, 427. Cause of variation in organic forms, 2460. Cedar longicorns, 319. Cave fanna of Kentucky. Coleopterous, 2033. Celæna renigera = Hadena renigera. Cecidomyia, 1077, 1461. CELERY. INSECTS AFFECTING. albovittata n. sp., 46. , Buprestis divaricata = Dicerca divaricata. carbonifera, 1924. Dicerca divaricata, 1063. cornuta n. sp., 46. Tettigonid, 452. Cranberry, 194. Trochilium sp., 1063. cratægi-bedeguar n. sp., 376. Celery. Leaf-hoppers on, 452.

Celtis, 2208.

Centennial insects, 1611.

Those, 1511.

Central America. Prevalence of screw-worm in,

2158.

Report on cotton crop and its enemies in, 2343.

Cephalomyia ovis = Œstrus ovis.

Cerambycidæ, 1324, 2267.

Food habits of, 1902.

Cerastophorus cinctus = Chion cinctus.

Ceratina dupla, 1171.

Ceratocampa imperialis = Eacles imperialis.

regalis = Citheronia regalis.

Ceratomia catalpæ, 2119.

hageni, 2119.

Cercopididæ, 882.

Cercopsylla n. g., 2272.

sideroxyli n. sp., 2272.

Cereals. Insects injurious to, 2238.

Ceresa bubalus, 415, 1323, 1329, 2107, 2195, 2235.

Cermatia forceps, 752, 764, 1154.

Cerococcus quercus, 2119.

Ceropales rufiventris n. sp., 375, 543.

Ceroplastes cirripediformis, 2450.

n. sp., 1964.

Ceropsylla n.g., 2272.

sideroxylin. sp., 2272.

Ceroptres ensiger, 41.

ficus, 745.

inermis, 41.

petiolicola, 41.

Cerotoma caminea, 2418.

Cerura multiscripta n. sp., 1411.

Cetonia, 2105.

inda = Euphoria inda.

Chærocampa pampinatrix = Ampelophaga myron.

Chætochilus pometellus, 874.

Chætophorus, 1678.

Chætoptria hypericana, 2076.

Chafer. Melancholy, 522.

Rose, 1329.

Chalcid egg parasites. Efficacy of, 2115.

fly, 1643, 1959.

Chalcididæ, 128, 384, 2115, 2343.

Joint-worm not one of the, 49.

Chalcis albifrons n. sp., 6 =Smicra albifrons.

fly, 1059.

Inflating, 1059.

mariæ n. sp., 1112 = Spilochalcis mariæ.

Mary, 1311.

ovata, 2343.

Chalcophora virginiensis, 2267.

Chalepus trachypygus, 273, 1911, 1949, 1973, 2119.

Chambers, V. T. Obituary, 2218.

Change of habit, 2096.

in vegetation caused by locusts, 1495.

Chapin's apple-leaf sewer, 1721.

Chariesterus antennator, 847.

Chauliodes. Larval characteristics of, 1652.

pectinicornis, 712.

rastricornis, 712.

Chauliognathus pennsylvanicus, 372, 374, 1959.

Cheese-fly and blow-fly, 1266.

Other insects affecting, 1703,

skipper, 1700.

Chelifer cancroides, 1884.

Chelymorpha argus, 767.

cribraria = C. argus.

Chemical changes in the color of butterfly wings,

Chenopodium hybridum. Lixus macer bred from,

Cherished bracon, 1423.

Chernetidæ, 1884.

Cherry aphis, 960.

borer. Flat-headed, 2267.

Caterpillars nest on wild, 242.

Cocoon on wild, 259.

CHERRY. INSECTS AFFECTING.

Acarid, 1265.

Aphis ccrasi = Myzus cerasi.

Attacus polyphemus, 1164.

promethea, 257.

Dicerca divaricata, 2267.

Eriocampa cerasi, 1222.

Hyphantria cunca, 242.

textor=H. cunea.

Myzus cerasi, 960, 1067, 1207, 1251.

Œcanthus niveus, 1503.

Scolytus rugulosus, 1940.

Selandria cerasi = Eriocampa cerasi.

Telea polyphemus = Attacus polyphemus.

Cherry plant-lice. Destroying, 1207.

and their foes, 1251.

tree borers, 1063.

plant lice, 1067.

trees. Minute borers in, 1940.

Slug on pear and, 1222.

twigs. Berry and, 1503.

Chester onion pcst, 2319.

Chestnut. Gall on, 353.

Gregarious worms on horse, 1192.

Phylloxera sp. on, 353.

Chicken mite, 1308.

Chickweed. Geometer, 1059, 1154.

Hæmatopis grataria on, 1059, 1154.

Chilo oryzæellus n sp., 2119, 2120.

Chilocorus bivulnerus, 38, 188, 301, 417, 730, 731, 1554, 1798, 1883.

CHINA-TREE. INSECTS AFFECTING, 1964.

Atta fervens, 1964.

Ceroplastes sp., 1964.

Lecanium sp., 1964.

Chinch-bug, 584, 886, 891, 894, 1127, 1423, 1508, 1681,

2018, 2039, 2048, 2271, 2363, 2418.

and army-worm prospects, 2087.

Bogus, 1127, 1423.

Bug allied to, 90.

Bugs on grape-vines mistaken for, 398.

Diseases of, 2143.

False, 1329, 1423, 2291, 2418.

in New York, 2237, 2252, 2271.

not in seed grain, 888.

notes, 2119, 2291.

once more, 890, 894.

bugs, S. A. Forbes. Experiments on, 2165.

Genuine vs. bogus, 1402.

Locusts vs., 1432.

Chion cinctus, 55.

Chionaspis euronymi, 2403.

Cicada. Song notes of the periodical, 2334. Chionapsis furfurus, 2, 188, 264, 345, 372, 373, 579, 951, 1008, 1040, 1061, 1152, 1554, 1968, in southeastern Indiana, Butler, A. W. The periodical, 2304. 2305. Massachusetts. Periodpinifolii, 97, 1329, 2079. ical, 2216. salicis, 373. nigræ n. sp., 373 = C. salicis. Sting of the, 371, 372, 407. Chipman, A. J. Locust notes in 1880, 2267. tibicen, 920, 1034, 1546. Chiro omus, 167, 625. tredecim = Tibicen tredecim. Chlænius impunctifrons, 2340. Twigs punctured by periodical, 1055. Chlamys plicata, 1363. years, 51. Chlorion cæruleum, 543. Cicadas, 26, 2311. Chloroneura n, g, 22 = Empoasca. Cotton-worms and, 1997. abnormis n. sp., 22 = Empoasca ab-Destroying, 2305. and walking-sticks, 920. normis. malefica n. sp., 22 = Empoasca viri-Cicadidæ, 431. Cicadula exitiosa, 1766. descens. maligna n. sp., 22 =Empoasca obtusa. Cicindela repanda, 1763. Chloroperla brunnipennis n. sp., 24.6-guttata, 347. fumipennis, 24. vulgaris, 628. uana n. sp., 24. Cicindelidæ, 719. Chlorops, 1461, 2060. Cimbex americana, 699, 1380, 1514, 2291, 2333, 2418. Chorinæus cariniger, 385. laportei = C. americana. Chortophaga viridifasciata, 1566. Cioidæ, 1937. Chronology of locust ravages in 1880-'81, 2267. Circotettix maculatus n. sp., 1959. Chrysididæ, 375. Circular No. 1, United States Entomological Com-Chrysobothris, 934. mission, 1575. femorata, 2, 6, 55, 352, 377, 379, 537, 2, United States Entomological Com-555, 578, 634, 870, 1059, 1227, 1250, mission, 1576. 1316, 1367, 1423, 1454, 1472, 1475, 9, U.S. Dept. Agric., Div. Ent., 2303. 1513, 1640, 2267. Cirrhophanus triangulifer, 2189. Chrysochus auratus, 735, 790. Cirrospilus esurus n. sp., 1712 = Tetrastichus csu-Chrysomela bigsbyana, 323. rus. cyanea = Gastroidea cyanea flavicinctus n. sp., in Lintner's first philadelphica, 323. report. (See foot-note 1, p. 379.) scalaris, 45. Cis fuscipes, 1737. Chrysomelidæ, 323, 2229. Citheronia regalis, 425, 629, 702, 775, 803, 860, 1087, Chrysopa plorabuuda, 584, 1127. 1275. sp., 372, 533. Clemens, B. Obituary, 223. Cicada, 189, 213, 372, 431. Cleonus sp., 1755. alias the seventeen-year locust. Period-Cleora. Evergreen, 2363. ical, 1159, 1971, 1979. Cleridæ, 1363. Belated individuals of the periodical, 648. Clerus nigripes = C. 4-guttatus. canicularis = C. tibicen. 4-guttatus, 200. cassini = Tibicen cassini. Click-beetle larvæ, 1643. Fungus in, 1809. beetles, 84, 351, 1625. in Iowa. Seventeen year, 1737. Climate on Cicada septeudecim. The influence of Little, 1242. **2**329. in Massachusetts. Periodical, 2321. for Doryphora, 1412. Note on the periodical, 2318. on prolongation of life of insects. Effects our first brood established. Periodical, of, 1129. 707. Climbing cut-worms, 1059. parvula = Melampsalta parvula. Clisiocampa americana, 66, 77, 112, 240, 277, 336, 350, Periodical, 474, 527, 619, 1059, 1159, 1489, 363, 642, 819, 1181, 1301, 1352, 2238. 1624, 1836, 2014, 2216, 2312, 2314, 2315, 2318, disstria, 240, 291, 328, 363, 645, 646, 688, 2320, 2363. 1127, 1181, 1200, 1301, 1331. Premature appearance of the periodical, sylvatica=C. disstria. 2320, 2326. Cloantha derupta, 2410. pruinosa = C. tibicen. Clæ dubia n. sp., 24. in savin-twigs. Eggs of the periodical, ferruginea n. sp., 24. fluctuans n. sp., 24.septendecim = Tibiceu septendecim. mendax n.sp., 24. Influence of climate on 2329. Close of the first volume, 701. Instinct of, 2144. Clostera americana = Ichthyura inclusa. Some popular fallacies re-Clothes moths, 70, 1633.

observed in the United States,

2146.

garding, 2367.

Seventeen-year, 1034, 1093, 1698, 2312, 2314.

397

Clover. Army-worm notes and inquiries, its work on, 1877. beetle. Flavescent, 2394. Bumble-becs vs. red, 2376. enemy. New imported, 2028. hay-worm, 1363. Army-worm vs., 2090. CLOVER. INSECTS AFFECTING, 1690, 2050. Asopia costalis, 83, 675, 948, 1132, 1363, 2090. Cecidomyia leguminicola, 1690, 1721, 2238. Drasteria ercchtea, 1459. Hylesinus trifolii, 1690, 1721, 1777, 1846. Leucania nnipuncta, 1877. Oscinis sp., 2394, Phytouomus punctatus, 2010, 2028, 2061, 2119. Pyralis olinalis = Asopia costalis. Sitones flavescens, 2394. Clover-leaf beetle, 2119. weevil. Further uotes on, 2061. New imported enemy to, 2010. root borer, 1721, 1846, 2512. secd midge, 1721, 2238, 2291. stem maggot, 2394. Webster, F.M. Insects affecting white, weevil, 1777. worms, 83, 675, 948, 1132. Clubbed tortoise beetle, 1127, 1558. Larva of, 1238. Cluster flies. Notes on, 2174. fly, 2128. Clydonopteron n.g., 1929.

tecomæ n. sp., 1929.

Clytus capræa = Neoclytus capræa.

Oak, 2267.

pictus = Cyllene pictus. robiniæ = Cyllene robiniæ.

speciosus = Plagionotus speciosus.

Coccid mistaken for a gall, 1972.

Coccidæ, 1329, 1883, 1968, 2088, 2137, 2235, 2520.

Means against, 104, 218, 264, 301, 487, 508, 944, 2372, 2373, 2403.

Monograph of, 1828.

Coccinella mnnda = C. sanguinca.

sanguinea, 6.

Coccinellidæ, 69, 322, 323, 639, 824, 1883, 1956, 2119.

to birds. Relations of Carabidæ and, 2152.

Food relations of Carabidæ and, 2151.

Habits of, 125.

Coccotorus sentellaris, 33, 34, 42, 254, 373, 377, 1301, 1378.

Coccus, 159.

harrisii = Chionaspis furfurus.

Cochylis hilarana, 1059.

Cocklebnr. Rhodobænus 13-punctatus on, 1301. Sphenophorus, 1301.

Cockroach eggs, 573.

Cockroaches. Means against, 1197.

Cockscomb elm gall, 1384.

Codling moth, 57, 307, 373, 1022, 1059, 1127, 1254, 1329, 1334, 1363, 1476, 1677, 2119, 2293, 2418.

again, 1301, 1311. heresies, 1398. Codling moth, jarriug down infested fruit, 1318.

' Michigau apples and, 1677.

Wier's trap, 1334.

Cœliodes inæqualis = Craponius inæqualis. Cœlodasys unicornis, 749, 880, 1280.

Coffee-borer, 498.

Cohn, F. The Hessian fly in Silesia in 1869, 2267. Colaspis, 372.

barbara, 1301.

flavida, 129, 133, 135, 231, 993, 1301, 1311, 1904.

Grape-vine, 231, 1301.

Cold applied to chrysalids of butterflies. Effects of, 1778.

on Curculio. Influence of extreme, 1335. and hibernating apple-worms. Severe, 2037. on insects. Effects of severe, 1818.

the scale insects of the orange in Florida. Voyle, J. Effects of, 2277.

Coleophora. Apple, 1721.

malivorella n. sp., 1721.

Coleoptera, 1329, 1904, 2267.

Classification of North American, 2085.

2085. Myrmicophilous, 2105.

of North America. The new classiffcation of, 2175.

Parasitic, 2353.

Willow-galls made by, 46.

Coleopterorum. Catalogus, 1866.

Coleopterous cave fauna of Kentucky, 2033. larvæ. Remarkable, 40.

Colias. The genus, 2259. philodice, 461.

Collecting and prescrying insects, 881, 1057. Colonies. Silk-culture in the, 2278.

Colopha ulmicola, 27, 137, 151, 518, 1384, 1678.

Colorado beetle poisonous? Is the, 1405.

cabbage flea-beetle, 2291.

grasshopper, 502.

potato-beetle, 21, 991, 1059, 1311, 1329, 1363, 1423, 1482, 1558, 1570, 2291, 2294, 2418.

abroad, 1364, 1407, 1431.

again, 1301.

Bogus, 1558.

in Europe, 1597.

Experience with the, 1518.

Inquiries coucerning the, 1491.

Insect euemies of the, 411.

Mite parasite of the, 1505.

Native home of the, 1462.

in New Hampshire, 1859. New York, 1379.

Onward march of the, 1218.

poisonous? Is the, 1483.
Poisonous qualities of
the, 1425.
Specific for, 1490.

potato-bug, 293, 334, 366. Bogus, 519. 398 INDEX.

Colorado potato-bug. Poisonous qualities of the, Cordulia? molesta n. sp., 39. 1101. Cordyceps, 332, 478. bugs. To destroy, 650. ravenelii, 594, 640. Coloration in insects. Unity of, 50. Coreus tristis = Anasa tristis. Come. Still they, 908. Corimelæna pulicaria, 354, 637, 741, 1046. Common Curculio and its allies, 254. Corn bill-bug, 2119, 2291, 2418. flesh-fly, 1423, 1570, 1625, 1643. borer, 754. longicorn pine-borer, 2291. Confessing the, 256. Curculio, 1042. May beetle. Eggs of, 1329. Commoner insects. Notes on our, 1802. cut-worm, 1059. Companion wheat-fly, 2394. ear-worm, 2238. feeding Syrphus-fly, 2422. Complimentary, 504. Compositæ. Antistrophus confined to the, 1131. fields. Beetles in, 2317. Compsomyia macellaria, 2256. CORN. INSECTS AFFECTING, 2119. Agonoderus pallipes, 2317. Comstock on the classification of insects, 2390. Anthomyia zeas, 657, 1059, 1066. Diaspinæ, 2227. Aphis maidis, 23. J. H. Report on miscellaneous insects, 2119. Butalis cerealella, 1232. Calandra granaria, 1683. Conchylis erigeronana n. sp., 1968. oryzæ, 1683. cenotherana n. sp., 1968. Cetonia inda = Euphoria inda. Conclusions. Jumping at, 253. Cut-worm, 1073. Cone nose. Blood-sucking, 795, 854. Diabrotica longicornis, 1905, 2187. Confessing the corn, 256. Empretia stimulea, 754. Confounding friend with foe, 1378. Euphoria inda, 1655. Conocephalus, 2267. Gortyna nitela, 754. Conorhinus. Blood.sucking, 2291. sp., 138. sanguisugus, 497, 795, 854. Conotrackelus, 934. Hadena sp., 1047. anaglypticus, 372. Heliothis armigera, 192, 1301, 1664, 2119, 2238. cratægi n. sp., 35, 358, 1301, 1358. Ips fasciatus, 4. geminatus, 34. 4-signatus = I. fasciatus. nenuphar, 6, 33, 42, 50, 98, 188, 237, 254, Lepidopteron, 162. 290, 329, 367, 372, 373, 377, 387, 439, Mesograpta polita, 2422. 487, 501, 566, 589, 629, 649, 680, 703, Noctuid, 181. 873, 930, 958, 978, 1056, 1059, 1062, Pempelia lignosella, 2119. 1069, 1071, 1129, 1135, 1173, 1177, Sphenophori, 2030. 1201, 1214, 1258, 1301, 1325, 1357, Sphenophorus robustus, 2119. 1358, 1372, 1477, 1685, 2230, 2296. sculptilis, 337, 1042, 1301. puncticollis n. sp., 34 = C. geminazeas = S. sculptilis. tus. Corn insects. Potato and, 138. Controlling sex in butterflies, 1352. Larval habits of Sphenophori that attack, Convolulateæ, Insects, 1238. 2030. Cook on insecticides. Review of, 2501. A new enemy to, 1905. Cooke's injurious insects of the orchard. Review foe of the, 337. of, 2245. insect foe of green, 1655. Copidryas gloveri, 2486. pest. Unknown, 1073. Copper underwing. Spattered, 1301. plant louse, 2418. Copris carolina, 416. root Diabrotica, 2187. Coptocycla, 1238. louse, a new enemy to the corn, 23. aurichaleca, 681, 736, 802, 1127, 1510. rustic, 1059. bivittata = Cassida bivittata. Serious injury to stored and cribbed, 1683. clavata, 693, 742, 767, 1238, 1558. Sphenophorus, 1301. guttata, 1127, 1510. stalk borer. Smaller, 2119. nigripes = Cassida nigripes. Webster, F. M. Insects affecting, 2418. Coquillett, D. W. Remedies for cottony cushion. worm, 192, 820, 1301. scale, 2394. Worm in 181. Report on the gas treatment for worms, 162, 1047, 1664, 2119. scale insects, The war on, 1522. 2418. Corrected. A few errors, 977. locusts of the San Correction, 939. Joaquin Val-Entomological, 1328. ley, California, Phylloxera, 1355. 2363. Corrodentia, 2267. production and

manufacture of

buhach, 2364.

Corticaria pumila, 2418.

Corycia vestaliata, 744,

Corydalus cornutus, 130, 308, 473, 1329, 1570, 1584, 1627, 1652.

Larval characters of, 1652.

Corynetes rufipes = Necrobia rufipes.

Corythuca ciliata, 200, 400.

Coscinoptera dominica, 1363, 2093.

Cossidæ, Bailey, J. S. The tree borers of the family, 2253.

Cossus robiniæ, 37, 245, 967, 1003, 1121.

Cotalpa lanigera, 40.

Cotton belt. Smith, E. A., 2343.

boll-worm, 2238.

bolls. Euryonia melancholica vs., 2451. in Brazil, Branner, J. C. Insects injurious to, 2277.

caterpillar, 1353, 1801.

caterpillars in Brazil, Branner, J. C., 2343. convention. Address at Atlanta, 2038.

crop and its enemies in Mexico, Central and South America, and the West Indies. Report on the, 2343.

culture and the insects affecting the plant at Bahia, Brazil, 1788a.

destroyers, 1886.

field mistaken for those of the cottonworm. Chrysalids dug up in, 1790. insects, 636.

COTTON. INSECTS AFFECTING.

Aletia argillacea = A. xylina.

xylina, 646, 1127, 1338, 1353, 1363, 1369, 1649, 1657, 1702, 1719, 1721, 1722, 1736, 1758, 1769, 1814, 1845, 1852, 1882, 1886, 1997, 2038, 2064, 2067, 2119, 2130, 2155, 2164, 2295, 2343.

Anomis xylina = Aletia xylina.

Euphoria melancholica, 2451.

Euryomia melancholica = Euphoria melancholica.

Gryllus sp., 2384.

Heliothis armigera, 646, 1353, 2238, 2343

Hyperchiria io, 2343.

Saturnia io = Hyperchiria io.

Spilosoma acræa, 2343.

Theclapeas ? 1872.

Cotton insects, Jones, R. W. Report on, 2343.

W.J. Report on, 2343. Kcrosene as a means against, 2164.

moth. Habits of, 1758.

Hibernation, 1728.

plant. Insects affecting, 1721.

planters. Suggestions to, 1832.

Plums and, 1477.

seeds in hot-beds. Sowing, 1772.

squares. Butterfly larvæ injurious to, 1872.

Cottonwood borers, 1525.

dagger, 1127.

A foe to, 1834.

COTTONWOOD. INSECTS AFFECTING.

Acronycta lepusculina, 1127.

Aphid, 495.

Drasteria ercchtea, 1459.

Lina scripta, 1654, 1834, 2291.

Pemphigus populicaulis, 446, 713.

COTTONWOOD. INSECTS AFFECTING-Continued.

· Pemphigus vagabundus, 446.

Plectrodera scalator, 1525.

Saperda calcarata, 71.

Cottonwood killer, 1654.

leaf beetle. Streaked, 2291,

galls, 713.

Unsightly galls on, 446.

Worms on, 1459.

Cotton-worm, 1127, 1363, 1649, 1702, 1719, 2077, 2119,

2130, 2164, 2295.

articles in the News, 1719.

Barnard, W.S. Tests of machinery for destroying, 2253.

Bulletin on the, 1736.

Chrysalids dug up in corn-fields mistaken for those of the, 1790.

damage in 1881, 2119.

To destroy the, 1338.

destroyer. Best, 1882.

Early appearance of, 184.

Facts about, 1657.

How to manage the, 1831.

use Paris green for the, 1544.

injury in regions overflowed by Mississippi. Protection from, 2119.

investigation, 1833.

Jones, R. W. Observations and experiments on, 2164.

moth hibernation, 1953.

Not the, 1773.

notes from Vera Cruz, Mexico. Interesting, 1845.

Orange insects and, 2067.

parasites, 1712.

Parasites bred from, 1722.

Poisoning devices for the, 2119.

Possible food-plants for, 2064, 2119, 2155.

in prairie belt. First appearance of, 1852.

question of hibernation settled, 2119.

A remedy for the, 1369.

Report on, 2164.

in south Texas in 1883. Anderson, E. H., 2253.

Sowing cotton-seeds in hot-beds as a means against the, 1772.

in the United States, 1769.

Cotton worms, 2291.

and Cicadas, 1997.

Cottony cushion-scale, 2394, 2418.

Coquillett, D. W. Remedies for, 2394.

Koebele, A. Remedies for, 2394.

maple-scale, 1816, 2291.

scale insect on maples, 1515.

Couper's thorn-leaf gall. Mr., 376.

Crab-apple borers, 602.

Crack-jaw names, 592.

Crambus. Vagabond, 2119.

vulgivagellus, 1990, 2029, 2119.

Cranberry fruit worm, 2291, 2349.

galls, 194.

CRANBERRY. INSECTS AFFECTING, Ctenucha latreilliana = C. virginica. Acrohasis vaccinii, 2291. virginica, 1241. Cecidomyia sp., 194. Cucumber beetle. Striped, 773, 1127, 2238. Phytopus sp., 2493. Cranberry leaf-folder, 2291. Smith, J. B. Report on insects affecting the hop and the, 2277, 2291. Crane-flies, 1278. fly larva, 512. Craponius inæqualis, 367, 373, 1059. Cratægus. Insects on, 376. Cratonychus incertus = Melanotus incertus. Cratypedes putnami, 1959. Culex, 625. Cremastochilus, 2105. Creophilus villosus, 805. Crepidodera brevis, 1782. Culicidæ, 603, 2162. cucumeris, 357, 401, 1056, 1059, 1558, Distribution of species of, 1782. helxines, 358. parvula, 1782. Crescent. The little Turk and his, 329. Cribbed corn. Serious injury to, 1683. Cricket. Grapes cut by a tree, 414. in Louisiana. Destructive, 2384. in raspberry canes. Eggs of tree, 2195. Tree, 251, 953, 961, 999. Snowy tree, 2291. Western, 1959, 2267. Bruner, L. Report on, 2267. Histology of, 1959. Crickets, 143, 1363. Mole, 1270. Crioceris asparagi, 524, 2135. 12-punctata, 2135. Crippled moths, 729. Critic criticised, 222, 942. Criticism. Friendly, 1307. Crop gall-gnat, Wagner, B. Observations on the new, 2267. Croton-bug as a library pest, 1717. Cruel bug.hunters, 440. Crumpler. Apple-leaf, 949. in Georgia. Rascal, 1962. mistaken for Curculio. Apple-leaf Rascal leaf, 311, 373, 1311, 1580 Cryptocentrus n.g., 385 = Mesoleius. Cryptorhopalum, 1746. Cryptus albicaligatus n. sp., 385. alhisoleatus n. sp., 385. annulicornis = Pimpla annulicornis. cecropia, 1311. cinctipes n. sp., 385. conquisitor = Pimpla conquisitor. inquisitor = Pimpla inquisitor. junceus = Linoceras junceus. nigrical ceatus n.sp., 385. nuncius, 1112. picticoxus, 385. pleurivinctus = Pimpla conquisitor. rhomhoidalis n. sp., 385. rufifrons n. sp., 385. samiæ, 2343. Cteniscus alhilineatus n. sp., 385.

ornatus, 385.

bug. Striped, 148. flea-beetle, 1059, 1558. CUCUMBER. INSECTS AFFECTING. Diabrotica vittata, 148, 773. Eudioptis nitidalis, 808, 843. Phakellura nitidalis = Eudioptis nitidalis. Cucumber. Worm boring in, 808, 843. Cucnrbitaceous vines. Insects affecting, 1127. mosquito, 2185. pipiens, 2071. Cupuliferæ. Cynips confined to, 1131. Curculio, 958, 978, 1062, 1080, 1127. again. Comparative scarcity of the, 703. and its allies. Common, 254. Apple, 1176, 1301. leaf crumpler mistaken for the, snout-beetle or four-humped, and bark-lice, 487. hreeds in apple. Plum, 1214. Carabidæ feeding on, 372. catcher, 875. Hull's, 372, 651, 875. Corn, 1042. extermination possible. Great discovery, 1173. Fighting the, 237. Flea-beetles and, 873. flies by night. How the, 706. Four-humped, 1088. Gas-waste vs., 549. go underground to transform? Does the apple, 1228. humbug. Another new, 589. A new, 680. Influence of extreme cold on the, 1335. injurious to grape, 267. Means against the, 290. natural history and how to catch him. Plum, 1372. parasite. Porizon, 1301. Sigalphus, 1301. Parasites of the plum, 1751, 1795. Paris green for the, 1258. Plum, 1059, 1129, 1301, 2296. To protect plums from the, 1071. Quince, 1301. remedies, 566. remedy. The new, 1177. Ransom, 1201. scarcer than last year, 620. trap. Smith's patent, 969. Use of the word, 1069. Curculionidæ. Monograph of, 1828. Curculios on pears, 1358. Currant borers, 481. bush borer, 974.

Currant, Currant-worms and black, 1079. Four-lined leaf-bug on, 715. and gooseberry worms, 1570.

CURRANT. INSECTS AFFECTING.

Ægeria tipuliformis, 340, 481, 974, 2238.

Alcathæ caudatum, 481.

Amblycorypha oblongifolia, 1157.

Aphis ribis = Myzus ribis.

Capsus 4-vittatus = Pœcilocapsus lineatus.

Ellopia ribearia = Eufitchia ribearia.

Eufitchia ribearia, 176, 364, 772, 877, 1224.

Mytilaspis pomicorticis=M. pomorum.

pomorum, 973.

Myzus ribis, 304.

Nematus ribesii, 78, 340, 364, 380, 381, 772, 1031,

1224, 1570, 1696, 2238.

ribis = N. ribesii.

ventricosus=N. ribesii.

Phylloptera oblongifolia = Amblycorypha oblongifolia.

Pecilocapsus lineatus, 715.

Pristiphora grossulariæ, 176, 364, 772, 1570,

Prosopis affinis, 1171.

Psenocerus supernotatus, 481.

Silvanus surinamensis, 1260.

Tenthredinid, 136.

Trochilium tipuliformis = Ægeria tipuli-

formis.

caudatum = Alcathe caudatum.

Currant plant-lice, 304.

louse, 322.

stalk-borer, 2238.

worm, 877, 1204.

Imported, 1127, 1570, 2238.

Native, 1570, 2238.

Pear-slug and, 1031.

worms, 364, 882, 1696.

Destructive, 78.

Gooseberry and, 772.

Currants. Beetle in dried English, 1260.

Green worms on gooseberries and,

Cushion-scale. Cottony, 2394, 2418.

Coquillett, D.W. Remedies for, 2394.

Koebele, A. Remedies for, 2394.

Cushions. Ravages of moths in, 1731.

Cussed red-leg, 1598.

Cuterebra buccata, 526.

Cut-worm, 868,992.

Climbing, 1059.

Corn, 1059.

Dark-sided, 281, 1059, 2291, 2363.

Dingy, 1059.

eggs, 1095.

Glassy, 1059, 2291, 2394.

Granulated, 2291.

Greasy, 1059, 2291, 2418.

lion, 1059, 1314.

moth. Eggs of, 607.

Shagreened, 2291.

Small white bristly, 1059,

Speckled, 1059, 2291,

Cnt-worm. Variegated, 1059, 2291.

W-marked, 1059, 2291.

Western striped, 1059.

Wheat, 1059.

Yellow-headed, 1059.

Cut-worms, 3, 85, 321, 658, 914, 1059, 1315, 2238, 2291.

Cabbage, 2291.

A chapter on, 964.

Climbing, 321, 359.

destroying recently sown wheat, 455.

Means against, 226, 312, 1059, 2387.

Novel facts about, 901.

originate? How, 787.

and other insects? Can land be in-

sured against, 801.

parasites, 349.

Remedy for cabbage, 2291.

Ridding the ground of, 2310.

severing cabbage-plants, 626.

Smilax injured by, 1941.

from the stomach of a robin, 1873.

Tree, 228, 281.

Cybister fimbriolatus, 750, 816.

Cybocephalus californicus, 2079.

Habits of, 2079.

nigritulus, 2079.

Cyclocephala immaculata, 1236.

Cyclone nozzle, 2327.

Cylindrical Orthosoma, 1059.

Cyllene, 560.

decorus, 50.

pictus, 45, 50, 55, 89, 269, 308, 560, 828, 941,

1516

robiniæ, 37, 45, 50, 55, 141, 308, 472, 560, 828,

921, 941, 967, 1196, 1250, 1288, 1516.

Cynipid gall on oak twigs, 1822.

Cynipidæ, 518, 799, 821, 2072, 2343.

Dimorphism in, 41, 1980.

Illinois, 41.

Important work on, 2125.

List of, 41.

New classification, 41.

Synoptic table of genera, 821.

Cynips, 1131, 1559.

acienlata = Amphibolips spongifica.

bicolor = Rhodites bicolor.

confined to Cupuliferæ, 1131.

Dimorphism in, 41.

fecundatrix, 1606.

gallæ-tinctoriæ, 1560.

n. sp., 1822.

q.-californica = Andricus q.-californica.

q.-decidua, 1925.

q.-erinacei n. sp., 41 = A craspis q.-erinacei.

q.-flocci n. sp., 41 = Andricus q.-flocci.

q.-forticornis n. sp., 41 = Biorhiza forticor-

q.-frondosa, 821, 774, 1666.

q.-glandulus n. sp., 1606.

q.-inanis = Amphibolips q.-inanis.

q.·mamma n. sp., 518 = Holcaspis mamma.

q.-mellaria n. sp., 1942.

q.-operator = Audricus operator.

q.-operatola, 1352.

402 INDEX.

Delicate long-sting, 1329.

Deloyala = Coptocycla.

clavata = Coptocycla clavata.

Cynips, q. pilulæ n. sp., 41 =Cecidomyia q.-pilulæ. Deltocephalus xanthoneurus, 2116. q.-podagræ n. sp., 41 = Andricus q.-poda Delusion. A popular, 525. Dendrotettix n.g., 2410. q.-prunus n. sp., 518 = Amphibolips q.-pruquercus n. sp., 2410. Deræocoris rapidus = Calocoris rapidus. nus. Dermanyssus avium, 1308. q.-punctata = Andricus q.-punctata. gallinæ, 1308. q.-saltatorius=Neureterus q.-saltatorius. q.-spongifica=Amphibilops q.-spongifica. Dermatobia hominis, 497. seminator = Andricus seminator. Dermatoptera, 2267. sp., 799. Dermestes lardarius, 72, 726, 1190, 1240, 1703. terminalis, 1448. Toothed, 2363. Cynthia atalanta = Pyrameis atalanta. vulpinus, 2363. cardui = Pyrameis cardui. Dermestidæ, 1746, 2133, 2180. Cypress. Cecidomyia cupressi-ananassa on, 1180. Means against, 56, 246, 462. galls, 1180. Deserved honor, 2124. Cyrtoneura a parasite or a scavenger? Is, 2102. Desmia maculalis, 468, 993, 1056, 1158, 1301, 1569, stabulans, 2102, 2343. 1579. Cytoleichus sarcoptoides, 2135, 2157. Desmocerus palliatus, 358. Cytophyllus concavus, 516, 569, 1005, 1157, 1363, Destructive larvæ, 722. locusts. Literature of, 1959. Dactylosphæra caryæ-semen n. sp., 373 = Phyllox-Devastating locust, 2363. era caryæ-semen. Devil's darning-needle, 1709. vitifoliæ = Phylloxera vastatrix. riding-horse, 457, 1059. Dexia rustica, 2260. Dactylosphæridæ, 1301. Dexidæ. Larval habits of, 2260. Dagger. Cottonwood, 1127. Smeared, 13.1. Diabrotica. Corn-root, 2187. Dahlia and aster stalk-borer, 949. 12-punctata, 123, 687, 2418. Gortyna nitela injuring, 862, 940, 1009. Long-horned, 1905. stalk-borer, 862, 1009. longicornis, 169, 1905, 2187. Daisy as an insecticide. Ox-eye, 1861. Twelve-spotted, 687, 1127, 2418. Dakota. Locust flights in, 2007. vittata, 123, 148, 175, 355, 772, 897, Dakruma convolutella, 1059. 2236, 2238, 2260. Dana, J. D. Criticism of the eutomological views Diapheromera, 2267. femorata, 144, 448, 920, 1111, 1395, of, 44. and his entomological speculations. Pro-1701, 1709, 1721, 2276. fessor, 82. velii n. sp., 45. Danaidæ, 613, 1301. Diaspinæ. Comstock on the, 2227. Danais archippus, 406, 461, 613, 1235, 1301, 1340, 1535, Diaspis harrisii = Chionaspis furfurus. 1622, 1635, 1711, 1756, 1770, 2052. rosæ, 1303. Daugerous looking, 814. Diastrophus, 318, 1131. Dapsilia rutilana, 1713, 1721. cuscutæformis, 609. Darapsa myron = Ampelophaga myron. nebulosus, 263, 610, 1121. Dardanelles. Sarcophaga lineata destructive to Dicerca divaricata, 871, 1063, 2267. locusts in, 2075. Didictyum n.g., 1749 = Hexaplasta. zigzag n. sp. = Hexaplasta zigzag. Dark grape-worm, 763. sided cut-worm, 1059, 2291, 2363. Diedrocephala flavipes n. sp., 1766, 1767, 2363. Darning-needle. Devil's, 1709. Differential locust, 1423, 2363. Darwin's work in entomology, 2129. Digger wasps, 309, 1643, 1959. Dascyllid in stomach of black bass, 1792. Dilar in North America, 2023. Dasyllis thoracica, 1226. Dilophus, 2076. Datana contracta, 50. Diminished Pezomachus, 1127. ministra, 50, 132, 151, 165, 467, 922, 1045, Dimmock's inaugural dissertation, 2082. 1229, 2222, 2333. Dimorphic asexual females. Lichtenstein's theory Decatoma dubia n. sp., 384 = var. of varians.as to, 2072. excrucians n. sp., 384 = var. of nigri-Dimorphism in Cynipidæ, 1980. ceps. Cynips, 41. hyalipennis n. sp., 384. Dytiscidæ, 28. nigriceps n. sp., 384. locusts, 1889. nubilistigma n. sp., 384. Dingy cut-worm, 1059. simplicistigma n. sp., 384. Dinoderus pusillus, 2104. sp., 536. as a museum pest, 2104, varians n. sp., 384. Dionea, 1499, 1688. Defense. Inexpert, 2476. Diphryx prolatella, 2120.

Diplax rubicundula, 2007.

Diplosis annulipes n. sp., 46.

Diplodus luridus, 372.

Diplosis atricornis n. sp., 46.atrocularis n.sp., 46.decem-maculata n. sp., 46. found in phylloxera galls, 1901. grassator, 2150. helianthi-bulla n. sp., 197. nigra, 2363, 2393. Pear, 2363. pyrivora n. sp., 2393. septem-maculata n. sp., 46. tritici, 109, 110, 142, 216, 280, 292, 372, 428, 711, 1512, 1690, 1721, 2238. Diplotaxis not a vesicant, 1796. sordida, 1926. Diptera, 1329, 1736, 2267. Parasitic, 2076. Preparation of, 2041. Swarming of, 2128, 2174. Willow-galls made by, 46. Dipterous enemies of the honey-bce, 704. Phylloxera vastatrix, 2150. galls of the willow, 46. larva on swallow, 153. larvæ in human body, 2256. Remarkable, 40. Discontinuance of publication [of Am. Ent.], 2106. Disease. Infesting Phylloxera with fungus, 1808. Diseases of beneficial insects. Fungus, 1813. chinch-bug, 2143. Disippus butterfly, 1301. Pupa of, 1193. Microgaster, 1301. Disonycha alternata, 45. flaviventris, 1506. Dissosteria carolina, 667. Ditching for young locusts, 1488. Dobson, 780. Dock. Galls on supposed, 1165. DOCK. INSECTS AFFECTING. Gastroidea cyanea, 1165. Gastrophysa cyanea = Gastroidea cyanea. Doctoring fruit-trees again, 230. Doctors differ, 98. Dog-day harvest-fly, 1546. Dolerus arvensis, 358. unicolor, 1989. Dollars. That hundred and fifty million, 1647. Domesticated katydids, 1536. Dominican case-bearer, 1363. Dorthesia celastri, 1988. characias, 1686, 1730. Covering of egg-puncture mistakeu for, 1988. viburni, 1988. Doryphora, 1639. Climate for, 1412. 10-lineata, 21, 32, 48, 50, 88, 94, 119, 171,

275, 276, 293, 313, 334, 347,

365, 401, 411, 565, 650, 708, 709, 760, 866, 905, 908, 919,

925, 946, 968, 977, 983, 991,

995, 1019, 1027, 1044, 1056,

1059, 1068, 1096, 1101, 1218,

1301, 1311, 1325, 1329, 1363,

1364, 1379, 1391, 1405, 1407,

Doryphora, 10-lineata, 1412, 1423, 1425, 1431, 1447, 1450, 1462, 1469, 1480, 1482, 1490, 1491, 1505, 1519, 1524, 1528, 1558, 1564, 1570, 1593, 1597, 1610, 1639, 1672, 1714, 1859, 1948, 1993, 2081, 2235, 2291, 2294. Fire cure for, 121. juncta, 48, 50, 372, 401, 519, 1056, 1059, 1431, 1450, 1558, 1672, 2096, 2215. Dotted-legged plant bug 1311. Double cocoons. The shedding of tracheæ, 1715. Dragon-flies, 1707. Dragon-fly. Large, 759. Drasteria erechtea, 631, 1459, 2343. Drasterius amabilis, 1798, 1905. dorsalis, 2418. Drepanosiphum, 1678. Dried leaves as food for lepidopterous larvæ, 2159. Driving nails into fruit-trees, 87. potato-beetles, 1044. Drones to death? Do worker bees sting, 1145. Drop of gold, 775. worm, 1059, 1090, 1650. Drosera, 1499, 1688. Drosophila amœna, 2119. ampelophila, 2119. nigricornis, 1607. sp., 1607. Drought. Connection of locust invasion with the occurrence of, 1422. on Hessian-fly. Effects of, 2031. Insects and, 2100. Drug-store pests, 510, 551. Dryobins sexfasciatus, 583. Dryocampa, 1352. bicolor = Sphingicampa bicolor. pellucida = D. virginiensis. rubicunda, 244, 841, 915, 1329, 1352, 1357, 2490. senatoria, 151, 783. stigma, 488. virginiensis, 488. Dutchman's pipo. Papilio philenor on, 1321, 1774. Worms on, 1321. Dynastes tityrus, 580, 1216, 1292. Dysdercus suturellus, 2343. Dytiscid in stomach of black bass, 1792. Dytiscidæ. Dimorphic, 28. Dytiscus. Modo of feeding of the larva of, 2065. Eacles imperialis, 1087, 1268. Earthquake commotion and locust multiplication and migration, Swinton, A. H., 2267. Eastern States. Supposed army-worm in New York and, 1990. Eburia 4-geminata, 1014. Eccopsis ferrugineana, 1968. Hickory, 2363. monetiferana, 1968. permundana, 2324. Variegated, 2363. Echthrus annulic rnis n. sp., 385. Economic entomology, 1357, 1362. in California. Interest in,

2053,

Colopha ulmicola, 1384,

Economic entomology in England, 1983. ELM. INSECTS AFFECTING—Continued. Dryobius sexfasciatus, 583. Fostering the study of, Galeruca calmariensis = G. xanthomelæna. 2142. xanthomelæna, 1350, 1520, 1931, 2188, Importance of, 2238. 2232, 2304, 2325, 2378, 2394. Iowa, 2197. Recent advances in, 2281. Monocesta coryli, 1721. investigations in the South and West, Pemphiginæ, 1653. Physocnemum brevilineum, 696. 1864. notes, 2220, 2248. Saperda lateralis, 583. Ecpantheria scribonaria, 1140, 1153, 1311, 1802. Schizoneura rileyi, 1059. Ectobia germanica, 573, 1717, 2276. Tremex columba, 928. Elm-leaf beetle, 2188. Edema albifrons, 422, 913. Edinburgh International Forestry Exhibit. Re-Great, 1721. Imported, 2232, 2304, 2325, 2378, port of, 2342. Edwards' butterflics of North America (Review), Notes on the imported, 1931. 395, 479, 623, 827. and pear-tree borer, 928. Egg-feeding mites, 1637. parasite. Anthomyia, 1570, 1625, 1643. tree borer, 583, 696. parasites of Acrididæ, 2355. depredators, 2414. Efficacy of chalcid, 2115. louse. Woolly, 1059. plant. Enemies of, 2096, 2215. saw-fly, 699. EGG-PLANT. INSECTS AFFECTING. Emmon's New York report, 84. Cassida texana, 2096, 2215, 2235. Emperor. Eyed, 1363. Doryphora 10-lineata, 21. Tawny, 1363. juneta, 2096, 2215, 2235. Emphytns maculatus = Harpiphorus maculatus. Egg puncture mistaken for Dorthesia. Covering Empoa albicans n. sp., 22.Empoasca n.g., 22. punctures in raspberry and grape-vines, 2195. abnormis, 22. sack of some unknown spider, 1144. consobrina n.sp., 22.slits made by Homoptera, 79. malefica = E. viridescens. Eggs in or on canes or twigs, 1329. maligna = E. obtusa. Experiments with locust, 1572. obtusa n. sp., 22. growth of insect, 2241. viridescens n. sp., 22, 30, 79. 'hatch? Will unimpregnated, 1029. Empretia stimulea, 60, 161, 424, 811, 829, 837, 1092. in maple twigs. Rows of, 665. Emulsions. Kerosene, 2291. Mass of, 65, 240. of petroleum as insecticides, 2134. of moth, 611. and their value as inbut parasitic cocoons. Not, 1290. secticides, 2126, 2200. in peach twigs, 554. Enchenopa binotata, 725, 1913, 1988. pear twigs. Rows of, 577. Enchophyllum binotatum = Enchenopa binotata. on sugar-maple, 350. Endrosa quercus = Lachnosterna quercus. in sumach, 18. England. Doryphora 10-lineata in, 2081. Egypt. Bug-hunter in, 372. Economic entomology in, 1983. Eight-spotted forester, 343, 1059, 1127, 1363. Hessian-fly in, 2395. Eighth Report State Entomologist of Missouri, Introduction of the Hessian fly into, 1482. 2398. Emige unserer schädlicherer Insekten, 1325. English currants. Beetles in, 1260. Elachistus euplectri, 2343. sparrow. Anent the, 1667. Elaphidion parallelum = E. villosum. Ennomos alniaria, 75. putator = E. villosum. magnaria = E. alniaria. tectum, 1602. subsignaria = Eudalimia subsignaria. villosum, 288, 793, 839, 1556, 2267. Entilia concava = Publilia concava. Elateridæ, 3, 224, 430, 740, 846, 892, 932, 1030, 1504, sinuata, 163. 1522, 1866, 2238. Entomography of Hirmoneura, 2275. Entomological, 371, 869, 893, 1411. Monograph of, 1828. Elder. Attacus eccropia on, 1533. breakfast, 2311. Cecropia worm on, 1533. cabinet. Walsh, 1107. Elephantiasis or Filama disease, 2185. circular. A recent British, 2509. Ellopia ribearia = Eufitchia ribearia. collection, 1328. Elm. Enemics of the, 1350. commission. Appropriation for, 1843. enemy, 1520. Bulletin No. 1, 1577. gall. Cockscomb, 1384. 2, 1578. ELM. INSECTS AFFECTING. 3, 1736. Cimbex americana, 699. 6, 2026. laportei = C. américana, Circular No. 1, 1575.

2, 1576,

Entomological commission. Report. First, 1643. Entomoligists in Europe. Number of, 1865. Second, 1959. Self-taught, 276. Third, 2267. Address on, 1454. Entomology. Fourth, 2343. all a humbug, 63. Reports of the U.S. Best works on, 418. Books useful for the study of, 418. 1757. correction, 1328. in Buffalo, N. Y., 2035. ignorance in the North, 431. California. Interest of economic, South, 390. 2053. information, 1351. Darwin's work in, 2129. notes, 2, 1396, 1554, 1609, 1610, 1681, Descriptive, 1199. 1958, 2015, 2058, 2066, 2135, Discussion on, 1391. 2147, 2168, 2184, 2192, 2219, Economic, 1362. 2228, 2247, 2261. in England. Economic, 1983. of the year, 2235. Fostering the study of economic, papers read at A. A. A. S., 1893. 2142. periodicals. New, 2056. General truths in applied, 2291, 2292. quackery, 439. Importance of economic, 2238. queries. A bundle of, 556. indeed run mad, 227, 1224. question, 1463. in Iowa. Economic, 2197. reports. Index to Ontario, 2228. Lecture on, 1394. Missouri, 1680. at Minneapolis, 2212. Society. American, 621. in Missouri, 1361. Address of president be-New York, 2244, 2254. fore Washington, 2335. Permanent subsection of, 2027. specimens. Care of, 2200. and the phonograph, 2481. speculations of the N. E. school of Professor Riley to Dr. Schaffer, 2360. naturalists, 44. in reality. Practical, 914. Professor Dana and Recent advances in economic, 2281. his, 82. horticultural, work at the Department of Agri-2230. culture, 1892. its relation to agriculture and its works, 831. advancement, 1329. wanted, 1529. Report Illinois State Horticultural Entomologist. American, 1309. Society on, 1056. Salutatory, 386. Missouri, 1113, 1127. Canadian (Review), 395, 479, 827. Study of, 2345. caught napping, 1110. Work in, 202. of Illinois. State, 327. Entomophthora, 1813. for Minnesota. A State, 1108. calopteni, 2240. of Missouri. First report of State, Epeira, 1847. 1059. riparia = Argiope riparia. Second. 1127. sp., 1144. Third, 1301. spinea, 813. Fourth, 1311. Ephemera flaveola n. sp., 24.Fifth, 1329. flies. A hard story, 1043. Sixth, 1363. myops n. sp., 39. Seventh, 1423. Ephemerella n.g., 24.Eighth, 1482. consimilis n.sp., 24.Ninth, 1570. excrucians n. sp., 24. Index to reports of Ephemeridæ, 283, 525, 1043, 1205. State, 2026. Ephemerina, 2267. for New York. State, 1863. Ephemerinous genus Bætisca. Pupa of, 43. New York without an, 1330. Ephemeron, 283. One day's journal of a State, 383. Ephestía interpunctella, 325, 1293, 1896. for the Pacific coast, 2022. zeas = E. interpunctella. Report U.S., 1878, 1721. Ephialtes gigas n. sp., 385. 1881-'82, 2119. pusio n. sp., 385. 1883, 2232. pygmæns n. sp., 385. 1884, 2291. Ephydra californica, 2203. 1885, 2363. gracilis, 2203. 1886, 2394. hians, 2203. 1887, 2418. Epicærus imbricatus, 35, 296, 1301, 1868, 2291. in the South. Field for the, 1109. Epicauta, 1600, 1643, 1651, 2238, 2267. Entomologists at Boston, 1894. atrata = E. pennsylvanica. Death of noted, 1285. cinerea, 134, 185, 401, 1056, 1558, 1600.

Epicauta corvina, 1206. Euplectrus comstockii, 2343. Larval characters and habits of, 1600. platyhypenæ, 2343. Eupsalis minuta, 1363. habits, 1387. pennsylvanica, 38, 134, 185, 284, 362, 401, Europe and America. Problem of the hop-plant 732, 1056, 1558, 1600. louse in, 2400. puncticollis, 1381. The Colorado potato-beetle in, 1597. Grain Bruchus just imported from, 1120. vittata, 38, 134, 185, 401, 470, 794, 1056, 1230, 1558, 1600, 1860, 1959. Hagen, H. A. The Hessian-fly not im-Epidemic disease of Caloptenus differentialis, ported from, 2267. 2240. Number of entomologists in, 1865. Epilachna borealis, 125, 417, 1289. European natural encmies of the asparagus beetle. corrupta, 2135, 2235. Mew, 2458. Epimeris. Grape-vine, 1301, 1363. parasites. Importing, 208. Epipocus punctatus, 2406. Eurycreon rantalis, 2322, 2331, 2363. Epitrix brevis = Crepidodera brevis. Eurvomia melancholica = Euphoria melancholica. cucumeris = Crepidodera cucumeris. Euryptychia saligneana, 180, 1127, 2356. Eurytoma abnormicornis n. sp., 384. Distribution of species of, 1782. hirtipennis = Crepidodera parvula. auriceps n. sp., 384. Epizeuxis. Spruce, 2363. bicolor n. sp., 384. bolteri n. sp., 1059 = var. of E. diastro-Erax, 1127. bastardi, 1127, 1269. diastrophi n. sp., 384, 1131. Eriocampa cerasi, 1031, 1222, 1253, 1382, 2243. gigantea n. sp., 384. Eriosoma cornicola n. sp., 27 = Schizoneura corniglobulicola n.sp.,384=var. of prunicola fungicola n. sp., 27 = Schizoneura funprunicida n. sp., 384.punctiventris n. sp., 384. lanigera = Schizoneura lanigera. seminatrix n. sp., 384 = var. of auriceps.pyri = Schizoneura lanigera. sp., 62, 563. tessellata = Pemphigus tessellata. Eurytomid from stomach of Ohio shad, 1853. ulmi n. sp., 1059 = Schizoneura ulmi.Eurytomides, 384, 1223. Euschistus fissilis, 2363. Eristalis, 986. Errors corrected. A few, 977. Euthyrhynchus floridanus, 2472. Erynnis alceæ boring in stems of Malva sylves-Euura orbitalis, 821. perturbans n. sp., 197. tris, 1602. s.-gemma n. sp., 197, 821 = E. orbitalis. Erythroneura, 164. australis n. sp., 22. s.-nodus n. sp., 197. octonotata n. sp., 22. s.-ovum n. sp., 197, 821. tricineta = Typhlocyba tricineta. Evagoras viridis = Diplodus luridus. vitifex = Typhlocyba vitis. Evarthrus orbatus, 872. vitis = Typhlocyba vitis. Evergreen cleora, 2363. ziczac n. sp., 22. and other forest trees in New England and New York. Packard, A. S. Eucalyptus. Galls on, 1965. Enchætes egle, 413. Causes of destruction of, 2232, 2291, Euclemensia bassettella, 1972. 2363, 2364, 2382. Eudalimia subsignaria, 75. plant-lice, 1039. Eudamus tityrus, 787, 857. Evergreens. Borers in, 1324. Eudemis botrana, 557, 585, 792, 1059, 1232. Insects affecting, 271, 527, 1039, 1324, Eudioptis nitidalis, 808, 843, 1127. 2232, 2291, 2363. Exartema ferruginearum n. sp., 1968 = Eccopsis Eudryas grata, 1127, 1363. unio, 831, 1127, 1363. ferruginearum. montiferarum n. sp., 1968 = Eccopsis Eufitchia ribearia, 176, 364, 772, 877, 1068, 1224, 1570. Eumenes fraterna, 543, 1357, 1847. montiferarum. Eumenia atala, 2410. Excursion. Agricultural editorial, 1349. Eumenidæ, 770. to Rocky Mountains, 988. Eunomia eagrus, 2242. Exenteron ornatus n. sp., 385 =Cteniscus ornatus· Euonymus latifolia? Chionaspis euonymii on, 2403. Exetastes illinoiensis, 385. Scale on, 2403. suavcolens n. sp., 385.Euparia castanea, 2105. Exochiscus n.g., 385 = Orthocentrus. Eupelmus, 384, 1932, 2343. pusillus n. sp. = Orthocentrus pusilallynii, 2060, 2063, 2332. lus. mirabilis, 384, 563, 1363. Exochus albiceps n.sp., 385.

annulierus n. sp., 385.

doryphoræ, 1059, 1431, 1450, 1672.

flavicauda n. sp., 1127, 1442, 1670.

atriceps n.sp., 385.

Exorista cecropiæ n. sp., 1112.

vs. Antigaster, 1978.

melancholica, 372, 522, 842.

Euphanessa mendica, 2340.

inda, 447, 1655.

Euphoria hirtipes, 2105.

Exorista leucaniæ = Nemoræa leucaniæ. militaris = Nemoræa leucaniæ.

ostep-sackeni = Nemoræa Ieucaniæ.

Exothecus prodoxi n. sp., 1831.

Expected advent of the locust, 2307.

Extensile penetrating organ in a gamasid mite,

Exterminator. Fruit-pest, 2372.

Extinguisher. Treat's insect, 531.

Extreme cold on the Curculio. Influence of, 1335.

Exuviation in flight, 1961.

Exyra ridingsii, 1411.

semicrocea, 1385, 1390.

Eyed emperor, 1363.

Eyes and luminosity of fire-flies. Development of, 1840.

None so blind as thoso who shut their, 275.

Fagopyrum. Acrididæ that eat, 1645.

Failure of tea-roses, 1708.

Fall army-worm, 1256, 1267, 1282, 1301, 1482.

canker-worm, 1423, 1482, 2238.

web-worm, 1301, 1733, 2238, 2378, 2394.

on hickory, 460.

Tent caterpillars and, 819.

wheat, Webster, F. M. Insects affecting, 2291, 2363, 2394.

Falling water. Moths attracted to, 2108.

False caterpillars on the pine, 985.

Scotch and Austrian

pines, 1011.

chinch-bug, 1329, 1423, 2291, 2418.

indigo gall-moth, 1127.

Walshia amorphella on, 1127.

Fatherless and motherless race. That, 1650.

race. Some further facts regarding that, 1658.

Fatua denudata, 1733.

Fear. A groundless, 284.

Foathers. Attagenus injuring, 56.

Folting caused by a beetle. Remarkable, 2113.

Feniseca tarquinius, 2359, 2361.

Food-habits of, 2407.

Notes on, 2369.

Plant-feeding habit of, 2361.

- Fertilization of Yncca, 1329, 1336, 2171.

Fertilizers of alpine flowers, 1838.

Fertilizing plants, 324.

Few errors corrected, 977.

Fidia. Grape-vine, 1231.

longipes, 339.

sp., 102.

viticida n. sp., 272, 339, 1059, 1231.

Field cricket, 2291.

crops. Insects injurious to, 2238.

for the entomologist in the South, 1109.

Fiory ground-beetle, 1059.

Fifteen-spotted lady-bird, 959, 1311,

Fifth report State Entomologist of Missouri, 1329.

Fifty million dollars. That hundred and, 1647.

Fig insects, 2194.

Fighting the Curculio, 237.

Hessian-fly, 1587.

Fignier's Insect World (review), 408.

Filaria disease. Elephantiasis or, 2185.

sanguinis-hominis, 2185.

Filbert grape. Grape-vine, 1329.

Fir Paraphia, 2363.

Fire blight, 22, 30, 149, 285.

cure for potato-beetles, 121.

flies, 396, 1705.

Development of eyes and luminosity of,

Intermittence of phosphorescence, 1805. worm, 2291.

First report noxious insects of the State of Illinois, 373.

> State entomologist of Missonri, 1059. U. S. Entomological Commission, 1643.

volume. Close of the, 701.

Fischer on naphthaline as an insocticide. Review of, 2274.

Fish-fly. Large, 712, 903.

Fitch. Dr. Asa, 1781.

Flat-headed apple-tree borer, 1059, 2238, 2267.

in horse-chestnut,

1316.

twig borer, 555, 1423.

borer in soft maples, 1250.

cherry-borer, 2267.

peach-borer, 2267.

spruce-borer, 2267.

Flavescent clover-beetle, 2394.

Flea-beetle, 1583.

Colorado cabbage, 2291.

Cncumber, 1558.

Grape-vine, 298, 1252, 1301.

Larvæ of grape-vine, 1041, 1074.

Striped, 2291.

Wavy-striped, 348, 564, 2291.

in young tobacco-plants, 1782

Zimmerman's, 2291.

beetles, 636, 2418.

and Curculio, 873.

Moans against, 133.

Satisfactory remedy for, 2236.

like negro-bug, 1127, 1423, 2291, 2418. Fleas feeding on lepidopterons larvæ, 2110.

Flesh-flies, 1643, 1736.

fly. Common, 1423, 1570, 1625, 1643.

Flesh-worms, 459.

Flics in rooms. Swarms of minute, 596.

Sucking organs of, 2182.

Flight. Exuviation in, 1961.

in insects is directed. How, 1891.

Flights east of the Mississippi. Locust, 1549, 1613. Locust, 1590.

Floating apiaries, 1785.

Floods. One effect of the Mississippi, 2080.

Florida, Ashmead, W. H. Report on insocts injurious to garden crops in, 2388.

Voyle, J. Effects of cold on the scale ininsects of the orango in, 2277.

Flour. Poisonous, 586.

and rye. Worms in, 72.

Flouring-mills. Beetles in, 1261.

Flower-bug. Insidious, 1127, 1423, 2418.

Flower-garden. Insects in, 936. Flowering ash. Cocoons on, 950.

Flowers. Fertilizers of alpine, 1838.

Fluted-scale, 2389, 2418.

Recent California work against the,

Flying bug, 157.

locusts in Illinois, 1437, 1443.

Foe. Confounding friend with, 1378.

to corn. A new, 337.

cottonwood, 1834.

green corn. A new insect, 1655.

Scavenger mistaken for a, 416.

Cherry plant-lico and their, 1251.

Some friends and, 1274.

Folsom, Cal., Koebele, A. Notes on locusts about,

Food of Calopteron and Photinus, 2358.

habits of Feniseca tarquinius, 2407.

ground-beetles, 1760. longicorn beetles, 1902. Megilla maculata, 2145. thrushes, 1726.

of insectivorous plants, 1499.

for lepidopterous larvæ. Dried leaves as, 2159.

man. Locusts as, 1481.

Number of molts and length of larval life as influenced by, 2167.

plants for the cotton-worm. Possible, 2064, 2119, 2155.

of Samia cynthia, 2204.

relations of Carabidæ and Coccinellidæ, 2151. Salt-water insects used as, 2203.

for silk-worms, 1319.

trout, 1142.

Forage crops. Insects injurious to, 2238.

Forbes, S. A. Experiments on chinch-bugs, 2165. Forbes' investigations on the food of fresh-water fishes. Professor, 2515.

Forest insects. Larvæ of injurious, 2267.

Tent caterpillar of the, 645, 688, 1181, 1200, 1301, 1331.

tree insects, 2418.

Packard, A. S. Notes on, 2253.

trees in northern New England and New York, Packard, A. S. Causes of the destruction of evergreen and other, 2232, 2291, 2363, 2364, 2382.

and weeds of Germany. Ratzeburg. Review, 1110.

Forester. Eight-spotted, 343, 1059, 1127, 1363.

Forestry exhibition. Report on the Edinburgh international, 2342.

Forficula, 1798, 2267.

Formica aphidicola n. sp., 27.

fusca, 2105, 2226.

latipes n. sp., 27 = Lasius latipes. rufa, 2105, 2114,

Formicidæ, 438, 1278, 1594, 1987.

Fossil tineids, 2084. .

Four-humped Curculio, 1088.

Apple-snout beetle or, 1088.

lined leaf-bug on currant, 715.

Fourth report State entomologist of Missouri, 1311.

Fourth report U. S. Entomological Commission, 2343.

Fowls. Internal mite on, 2157.

Parasites on, 1308.

France affected by Phylloxora. Half the vine area of, 2020.

Grape Phylloxera in, 2193.

Fraternal potter-wasp, 1127.

Friend. Another insect, 879.

with foc. Confounding, 1378.

' unmasked, 374.

Friendly criticism, 1307.

notes, 1306.

Friends and foes. Some, 1274.

Inquiring, 1642, 1656.

and insect foes. Insect, 38.

Frog-spittle insects, 691.

Frogs, 217.

Frosted lightning hopper, 1329.

Fruit culture in the South, 2265.

foe. A new, 35.

and fruit trees. Insects injurious to, 2238.

Trimble's enemies of, 187.

growers. Insects of interest to, 367.

in northern Illinois. Six worst enemics of, 377.

insects in California, 2245.

invigorator again. Best's, 530.

Jarring down infested, 1318.

pest exterminators, 2372.

Snout-beetles injurious to, 1302.

trees again. Doctoring, 230.

Driving nails into, 87.

Icerya purchasi, an insect injurious to, 2401.

FRUIT TREES. INSECTS AFFECTING.

Epicærus imbricatus, 35.

Ithycerus noveboracensis, 16.

Lytta ænea = Pomphopæa ænea.

Oncideres cingulata? 442, 443.

Pomphopæa ænea, 1056.

Fruit trees. Insects injurious to, 16, 31.

Tarred paper for, 1568.

worm. Cranberry, 2291.

Tomato, 1136.

Fruits and grains, Webster, F. M. Experiments on the effect of punctures of Hemiptera on shrubs, 2382.

Fuchsia beetles, 133.

Fuchsias. Beetles injuring cabbages and, 1868.

FUCHSIAS. INSECTS AFFECTING.

Haltica carinata, 133, 1868.

exapta = H. carinata.

Fulgoridæ, 737.

Fuller. Retirement of Mr., 1910.

Fuller's rose-beetle, 1708, 1721, 2291.

in California, 1740. Habits of, 1708.

Not, 1825.

Fumago salicina, 2138.

Fumigation against orange scales, 2520.

Fungoid growths, 500.

Fungus in Cicada, 1809.

diseasos of beneficial insects, 1813.

Infecting Phylloxera with, 1808.

Fungus foes, 1937.

growths to destroy insects. Use of, 1808. Remarkable parasitic, 1313.

White-grub, 594.

Fuzzy galls on blackberry twigs, 609.

Gad-fly. Sheep, 887.

Galerita janus, 1824.

lecontei, 1824.

Galeruca calmariensis = G. xanthomelæna.

rufosanguinea = Adimonia rufosanguinea.

xanthomelæna, 1135, 1350, 1520, 1931, 2188, 2232, 2304, 2325, 2378, 2379, 2414.

Gall on acorn-cups. A new oak, 1606.

blackberry, 263.

and raspberry canes. Gouty, 1771.

chestnut, 353.

Coccid mistaken for a, 1972.

Cockscomb elm, 1384.

flies, 294.

gnat, Loew. Description of the rye, 2267.

Wagner. Observations on the new crop, 2267.

on grape-vine. Large compound, 720.

Grape-vine apple, 1329, 1898.

leaf, 724, 840, 1716.

insects, 1559.

Galls and, 1967.

Leafy oak, 774.

literature. Bibliography of, 2059.

louse. Hickory-stem, 685.

making genus of Apiouinæ, 2231.

moths, 1127, 2356.

Pemphiginæ. Biological notes on, 1653.

tortricid. A pretty and unique, 2176.

Mossy-rose, 570, 1166.

nuts, 1560.

Oak-fig, 745.

leaf, 799.

on oak twigs. Cynipid, 1822.

Pelargonium, 1764.

Pithy blackberry, 1131.

Pod-like willow, 1170.

Prickly rose, 1194, 1245.

Raspberry gouty, 1124.

root, 1149.

Rose, 1235.

on Solidago leaves, 1924.

spotted touch-me-not, 852.

Sugar-maple mite, 1265.

Trumpet grape, 791, 1116.

Galleria cereana, 581, 716, 904, 1059, 1293, 1357, 1508, 1723.

Gallmücke. Beschreibung einer den Birnen schädlichen, 2392.

Galls, 137, 1561.

on blackberry, 263.

cottonwood leaf, 713.

Unsightly, 446.

Cypress, 1180.

on Eucalyptus, 1965.

and gallinsects, 1967.

Golden·rod, 798.

Grape, 111, 160.

Galls on the grape-vine. Leaf, 724, 840, 1716.

growing on wild sage, 1347.

Hackberry psyllid, 2208.

Jumping seeds and, 1496, 2163, 2173.

on leaves of soft maple, 445.

wild grape-vine. Conical, 1077.

made by moths, 1127.

Phylloxera. New hickory, 1901.

and mining in apple-twigs. Small, 552.

Oak and rose, 1037.

Production of, 1056.

on sugar-berry, 762.

supposed dock, 1165.

and their architects, 518, 821.

on white oak. Woolly, 739.

Gamasid mite. An extensile penetrating organ in a, 1626.

Gamasus juloides infesting Iulus marginatus, 963.

Garden crops. Experiments with insecticides npon insects affecting, 2344.

in Florida, Ashmead, W. H. Report on insects injurious to, 2388.

insects, Alwood, W. B. Tests with insecticides ou, 2388.

Insects in the flower, 936.

vegetables. Insects injurious to, 2238. web-worm, 2363.

Gardens. Ants' nests in, 496.

Destroying black ants in, 438

Toads in, 822.

Gargaphia amorphæ, 45.

tiliæ, 45, 400.

Gas treatment for scale insects. Coquillett, D. W., 2418.

waste vs. Curculio, 549.

Gases against scale insects. Use of, 2389.

Gasteracantha cancer, 2111.

Gastroidea cyanea, 316, 1165.

Gastropacha americana, 812.

velleda = Tolype velleda.

Gastrophilus equi, 1391, 1563, 2238.

 ${\it Gastrophysa\ cyanea} = {\it Gastroidea\ cyanea}.$

Geelong, Victoria. Grape Phylloxera at, 1732.

 ${\bf Gelechia\ abietisella,\,2232.}$

cerealella, 2291.

gallæ-solidaginis n. sp., 1059, 1165, 2213.

glandulella n. sp., 1306.

Hemlock, 2232.

White-heart hickory, 2363.

Gemminger and Harold's Catalogus Coleopterorum, 1866.

General index and supplement to the Missour reports, 2026.

truths in applied entomology, 2291, 2292.

Genuine army-worm in the West, 2009.

vs. bogus chinch-bugs, 1402.

Genus grinding, 2120.

Geographical distribution of the Rocky Mountain locust, 2267.

range of species, 1614.

Geometer. Chickweed, 1059, 1154.

Knot-weed, 1059.

Geometra catenaria — Zerene catenaria. Geometridæ. Larval characters, 165.

Georgia. Rascal leaf-crumpler in, 1962. GOOSEBERRY. INSECTS AFFECTING. German asters. Diabrotica vittata on, 148. thoroughness, 1597. Gigantic rhinoceros beetle, 580, 1216, 1292. water-bug, 534, 733. Gilly-flowers. Cabbage-worms upon, 614. Plutella cruciferarum injuring, Gilt gold-beetle, 790. Girdled pear-twigs, 848. Girdler. Twig, 476, 1938. Girdling habits of Pædisca obfuscata, 2405. Glassy cut-worm, 1059, 2291, 2394. Mesochorus, 1127. winged soldier-bug, 1301. Protective device em-Glaucopid caterpillar. ployed by a, 2242. Glorified squash-bug, 262. Glossina morsitans, 1467. Glover, Townend. Obituary, 2218, 2258. Glow-worm, 1917. Nature of the phosphorescence of, That, 1097. Glyphe viridascens n. sp., 12. Glypta alboscutellaris n. sp., 385. diversipes n. sp., 385. ruficornis n. sp., 385. rufipleuralis n. sp., 385. Gnats, 625. Goat-weed butterfly, 1117, 1127, 1329. Paphia glycerium feeding on,1329. Going it blind, 588. Gold-beetle. Gilt, 790. Drop of, 775. Golden-rod galls, 798. GOLDEN-ROD. INSECTS AFFECTING. Cecidomyia carbonifera, 1924. solidaginis, 798. Euryptychia saligineana = Pædisca scudderi-Gelechia gallæ-solidaginis, 1059, 1165. Graphtholitha olivaceana, 2285. Nectarophora rudbeckiæ, 2205. Pædisca scudderiana, 180. Siphonophora rudbeckiæ =: Nectarophora rudbeckiæ. Trypeta solidaginis, 180, 798, 1059. Golden-rod tortoise-beetle, 1127. Gomphocerus shastanus n. sp., 1959. Gomphus amnicola n.sp., 24. consobrinus n. sp., 39. fluvialis n. sp., 24. grasilinellus n. sp., 24. quadricolor n. sp., 39. vastus n. sp., 24. ventricosus n.sp., 39. Gonatopus contortulus, 2116. pedestris, 2116. pilosus. Biology of, 2116. Goniocotes hologaster, 1094. Good words. More, 532. Gooseberries and currants. Green worms on, 136.

Gooseberry and current worms, 772.

fruit-worm, 1059.

Dakruma convolutella, 1059. Ellopia ribearia = Eufitchia ribearia. Epicærus imbricatus, 35. Eufitchia ribearia, 772, 1068, 1570. Nematus ribesii, 140, 228, 333, 756, 772. ventricosus = N. ribesii. Pristiphora grossulariæ, 40, 189, 772. Tenthredinid, 136. Gooseberry saw-fly, 140. Imported, 228, 333. span-worms, 1068, 1570. worm, 189. worms. Currant and, 1570. Imported, 756. Gordiacæa, 612. Gordius, 861. aquaticus, 447, 1143. varius, 1143. Gortyna nebris, 1646. nitela, 331, 401, 632, 694, 734, 754, 820, 862, 940, 976, 993, 1009, 1056, 1059, 1210, 1558, 1589, 1595, 1646, 1870, 2229. sp., 138. Gouty gall on blackberry and raspberry canes, Governor of Kansas on the grasshopper question. Report to, 1573. Governors of Western States on the Rocky Mountain locust, 1557. Gracilaria. Purple willow, 2363. Grain Aphis, 2394. Bad work of, 1638. vs. rust, 1806. Bruchus, 1301. of Europe just imported, 1120. Chinch-bug not in seed, 888. GRAIN. INSECTS AFFECTING. Aphid, 99. Asopia farinalis, 2337. Bruchus granarius, 1301. Gelcchia cerealella, 2291. . Nectarophora granaria, 126. Grain moth, 2337. Angoumois, 2291. A new leaf-hopper injurious to small, 1767. plaut louse, 99, 126, 1127. Silvanus, 1259. Sphenophorus, 2363, 2394. wcevil, 15, 2291. weevils. Aniseed vs., 1742. Grains. Calandra granaria in, 1259. and grasses, Webster, F. M. Insects affecting, 2394. Webster, F. M. Experiments on the effect of punctures of Hemiptera on shrubs, fruits, aud, 2382. Granddaddy long-legs, 838. Granulated cut-worm, 2291. Grape-bark louse, 106. berry moth, 557, 585, 792. cane-gall Curculio, 1059. canes and apple-twigs. Eggs in, 1323. Egg on, 855. punctured, 513, 1333.

411

Penthina vitivorana = Eudemis botrana.

GRAPE-VINE. INSECTS AFFECTING-Continued. Grape Curculio, 373, 1059. disease, 1311. Amphicerus bicaudatus, 1185. fruit worm, 1059. Anomala lucicola, 113. Aphis vitis, 102, 170. gall. Trumpet, 791, 1116. galls, 111, 160. Apis mellifica, 441. insects, 164, 267, 1056, 1569. Blennocampa pygmæa, 1252. Bostrichus bicaudatus - Amphicerus bicaudaleaf-folder, 1301, 1579. folders eaten by spiders, 468. Hairy, 664. Capsus oblineatus = Lygus pratensis. gall-louse, 373, 1301, 1311. Cecidomyia vitis-coryloides, 1329. galls, 127, 1295, 1435, 1531. litnus, 791, 1116. Leaf-hoppers of the, 399. pomum, 403, 1284, 1329, 1898. leaf-louse, 916. viticola, 791, 1077, 1116, 1329. Phylloxera enemy, 1530. Cecidomyid, 111. Phylloxera, 1329, 1363, 1373, 1482, 1721 Ceramhycid, 429. in California, 1727. Ceresa buhalus, 1323, 1329. France, 2193. Chærocampa pampinatrix = Ampelophaga Geelong, Victoria, 1732. myron. New hiological facts regard-Celiodes inæqualis = Craponius inæqualis. Colaspis flavida, 129, 133, 135, 231, 1301, 1311. ing, 1421. Newest facts of, 1404, Corimelæna pulicaria, 1046. not at the Cape, 1841. Craponius inæqualis, 373, 1059. permaneutly destructive, Curculio, 267. 1907. Dactylosphæra vitifoliæ = Phylloxera vasta-Notes on the, 1952, 2291. trix. natural history Darapsa myron = Ampelophaga myron. of, 1439. Desmia maculalis, 468, 1158, 1301, 1569, 1579. Use of guano for, 1837. Drosophila ampelophila, 2119. root-borer, 373, 1301. Eggs, 855. New, 1059, 1127. Encheuopa binotata, 725. rot. Phylloxera and, 1623, 1628. Enchophyllum binotatum = Enchenopa binosaw-fly, 2291. scale insect, 1706. Erythroneura sp., 164. seed Curculio, 1059. tricincta=Typhlocyha tri. Insect infesting, 482. cincta. maggot, 1127. vitis = Typhlocyba vitis. vine Aphls, 170. Eudemis botrana, 557, 585, 792, 1059. apple-gall, 1329, 1898. Endryas grata, 1127, 1363. An apple growing on a, 403, 436. unio, 1127, 1363. Fidia longipes, 339. Bark-lice on, 1212. hcetles, 113, 129, 133, 339, 354. sp., 102. borer, 429. viticida, 272, 339, 1059, 1231. caterpillars, 124. Galls, 128, 791. caterpillars. Black, 1018. Graptodera chalybea = Haltica chalybea. Colaspis, 231, 1301. Haltica chalyhea, 203, 298, 1041, 1252, 1255, 1301, again, 1311. 1821, 1855. Conical galls on leaves of wild, 1077. Harrisiana americana, 164, 213, 1127, 1136. Epimeris, 1301, 1363. Isosoma vitis, 482, 484, 1059, 1127. Fidia, 272, 1059, 1231. Lasioptera vitis, 720, 1329. filhert-gall, 1329. Leaf-hopper, 484. flea-heetle, 298, 1252, 1301, 1821, 1855. Lecanium vitis = Pulvinaria vitis. Larvæ of, 1041, 1074. Lygns pratensis, 682. and fuchsia heetles, 133. Macrodactylus suhspinosus, 373, 565, 1375. a hickory. Hybrid hetween, 1284. Membracid, 286. hoppers, 1024. Membracis ampelopsidis, 1183. Injured strawherry and, 682. Œcanthus niveus, 286, 414, 723, 1059, 1323, 1329, insects, 128, 725. 1333, 1691, 2195. GRAPE-VINE. INSECTS AFFECTING. Oncometopia undata, 36, 79, 164, 399. Acoloithus falsarius, 124. Orocharis saltator, 513, 1323, 1329. Acronycta ohlinita, 1208, 1301. Orthosoma hrunneum, 397. Ægeria polistiformis=Sciapteron polistiforcylindricum = 0. hrunneum. Oxyptilus periscelidactylus, 664, 1059, 1175, Alypia octomaculata, 1059, 1127, 1130, 1208, 1363. Pelidnota punctata, 113, 129, 354, 725, 1221, 1301 Amblycorypha oblongifolia, 1329. Pemphigus, vitifoliæ = Phylloxera vastatrix. Ampeloglypter sesostris, 1059.

Ampelophaga myron, 1086, 1127, 1247.

INDEX.

412

GRAPE-VINE. INSECTS AFFECTING-Continued. Grapes spoiled by something, 464. Grapholitha gallæ-saliciana n. sp., 1968. Philampelus achemon, 1091, 1127. Grapholitha ninana n. sp., 2176. pandorus, 1102. olivaceana n. sp., 1968. satelliatia = P. pandorus. Habits of, 2285. Phylloxera vastatrix, 127, 160, 373, 565, 724, 840, pruinivora, 367, 373, 1795. 916, 1281, 1295, 1301, 1311, 1325, 1329, 1342, Graphops pubescens, 2229. 1363, 1374, 1376, 1421, Grapta, 1301. 1423, 1435, 1439, 1482, comma, 461, 1140. interrogationis, 420, 444, 1306. 1530, 1531, 1623, 1628, Graptodera carinata = Haltica carinata. 1716, 1721, 1727, 1732, chalybea = Haltica chalybea. 1837, 1841, 1895, 1900, punctipennis = Haltica punctipennis. 1907, 1952. Grass-bug and its habits, 882. vitifoliæ = P. vastatrix. Poeciloptera pruinosa, 1329. insects, 541. Polycaon confertus, 2103. GRASS. INSECTS AFFECTING. Prionus imbricornis, 1127, 1274. Laphygma frugiperda, 1127. laticollis, 561, 1059, 1081, 1127. Leucania unipuncta, 1877. Grasses, Webster, F. M. Insects affecting small Proconia undata = Oncometopia undata. Procris americana — Harrisiana americana. grains and, 2394. falsarius = Acoloithus falsarius. Grasshopper. The, 1588. Psychomorpha epimeris, 1301, 1363. Colorado, 502. Pterophorns periscelidactylns = Oxyptiluseggs, 536, 667. Trombidium preying on, 624. periscelidactylus. Pulvinaria vitis, 106, 1212, 1706, 1716. Governors of Western States on Pyrophila pyramidoides, 671, 1301. the Rocky Mountain, 1557. Sciapteron polistiformis, 373, 1301, 1509. Hateful, 373, 475. injuries. New method of coun-Selandria vitis = Blennocampa pygmæa. Sinoxylon basilare, 1311. teracting, 2235. Smilia auriculata, 1183. injury in the near future. Prob-Spilosoma virginica, 1202, 1301. abilities of, 2335. Tettigonia coagulata, 1024. Outlook for locust or, 2461. vitis = Typhlocyba vitis. machine. A satisfactory, 1592. Thyreus abbotii, 763, 1018, 1127, 1248. A new enemy of the, 1541. Typhlocyba tricincta, 25. Parasite on hateful, 728. vitis, 203, 686. pest of the West. Important observations on, 1571. Grape-vine. Insects injurious to the, 267, 1059, 1081, 1086, 1091, 1102, 1118, 1130, 1137, prospect, 1693. 1158, 1175, 1202, 1221, 1255, 1281, 1301, question. Report to governor of 1311, 1329, 1363. Kansas, 1573. Large compound gall on, 720. ravages in California, 2323. Rocky Mountain, 1557. leaf-galls, 724, 840, 1716. hoppers, 484, 686. Western, 1998. pest. More about the, 1376. year? Is this a, 1565. plume, 1059, 1301. Grasshoppers, 368, 391, 402, 433, 1363. Procris, 213. Destructive, 2363. root-borer, 1509. and locusts, 26, 147, 929. roots. Wood-lice on, 1906. Mites ou, 146. tomato-gall, 1329. Rear-horses vs., 590. trumpet-gall, 1329. Young, 2377. worm. Dark, 763. Grass-worm, 2119. Green, 671. Gray straight-horned snout-beetle. Large, 1033. Gray. Personal reminiscences of Dr. Asa, 2419. Pyramidal, 1301. vines. Bugs on, 1046. Greasy cut-worm, 1059, 2291, 2418. Caterpillars on, 1208. Great discovery. Curculio extermination possi-On the cause of deterioration in blc, 1173. some of our native, 1342. elm-leaf beetle, 1721: Egg-puncture in raspberry and, Lebia, 1301. 2195. leopard moth, 1311. Eggs of the tree-cricket on, 723. Green apple-leaf tyer, 1311. mistaken for chinch-bug. corn. A new insect foe to, 1655. on, 398. grape-vine worm, 671. to Phylloxera in sandy soil. Rehag moth, 1233. sistance of, 2250. striped maple-worm, 1329. Rose-chafers on, 1375. worms on gooseberries and currants, 136. Grapes cut off by tree-cricket, 414. Greenhouse pests, 508. Honey-bees eating, 441. plants, 508.

INDEX. 413

Gregarious worm on horse-chestnut, 1192. Halictus sp., 800. walnut caterpillars, 1045. Halisidota antiphola n. sp., 40 = H. tessellata. , caryæ, 45. willow-worms, 856. Ground-bectle. Another herbivorous, 2042. harrisii n. sp., 45 = H. tessellata. Fiery, 1059. tessellata, 40, 45, 50. Haltica alternata = Disonycha alternata. Murky, 486. Pennsylvania, 1059. carinata, 133, 1868. Subangular, 372. chalybea, 203, 298, 1041, 1074, 1252, 1255, 1301, bcetles, 1127, 1558, 1625, 1643. 1583, 1821, 1855. Eggs of, 692. cucumeris = Crepidodera cucumeris. exapta = H. carinata. Food-habits of, 1760. Vegetal feeding, 1738. hclxines = Crepidodera helxines. punctipennis, 2383, 2468. Groundless fear, 284. striolata = Phyllotreta vittata. Growth of insect eggs, 2241. Haltichella perpulchra, 11, 12. Grub fungus. White, 594, 640, 1064, 1430, 1436, Halticidæ, 873, 1858, 2236. 1599, 1803, 1823. Ham-beetle. Red-legged, 1363. information wanted. White, 1072. Hams. Skippers injuring smoked, 1734. White, 68, 410, 1020, 1059, 2238, 2363, 2394. Hand-maid moth, 2222. worm, 54. Hard story. Ephcmera flies, 1043. Grubs and guess-work, 1440. Harlequin cabbage-bug, 1099, 1311, 1534, 2238, 2263, Gryllidæ, 929, 2267. Gryllotalpa, 2267. borealis, 562, 1270, 1798. Harmless insects, 1526. Harold's Catalogus Coleopterorum. Gemminger columbia, 562. longipennis = G. columbia. and, 1866. Gryllus, 2267, 2384. Harpactor cinctus = Milyas cinctus. abbreviatus, 143, 433. Harpalus, 1643. Guano for grape Phylloxera. Use of, 1837. caliginosus, 3, 486. Guard. Be on the, 708. crraticus, 1537. Guess-work. Grubs and, 1440. pennsylvanicus, 372. Guide to the study of insects. Review of Pack-Harpiphorus maculatus, 499, 955, 965, 1056, 1264, ard's, 395, 479, 623, 827. 1570, 1586, 2324. Gymnetis nitida = Allorhina nitida. Harris's bark-louse, 373. Gyrinus larva in stomach of shad, 1853. correspondence (review), 623. Habit. Change of, 2096. insects injurious to vegetation (review), Hackberry butterflies, 1356, 1363. HACKBERRY. INSECTS AFFECTING. Harrisina americana, 164, 213, 789, 1056, 1127, 1136. Apatura alicia, 1363. Harvest bugs, 412. celtis, 1356, 1363. fly. Dog-day, 1546. clyton, 1356, 1363. mites, 1326. Hatch pupæ. How to, 509. herse = A. clyton. lycaon = A. celtis. Will unimpregnated eggs, 1029. Galls, 791. Hatching? Are the locusts, 1566. Pachypsylla c.-mamma, 2208. Hateful grasshopper, 373, 475. c.-venusta, 2208. Parasites on, 728. Hackberry psyllid galls, 2208. locust, 1452. Hadena, 2238. locusts. Prairie fires and, 1453. chenopodii = Mamestra trifolii. Hawk's pellets, 601, 643. devastatrix, 1056, 1059, 2291, 2394, 2355. Hawthorn. Worms on, 1051. juneta, 1056. Hay. Worms under mulch, 1161. renigera, 964, 1059. Head-maggot, 450, 1059. sp., 1047. of winged insects, Packard, A. S. Number subjuncta, 901, 1059. of segments in the, 2267. Hæmatopis gratariá, 1059, 1154. Hedge-hog caterpillar, 1153. Hag-moth. Green, 1233. Helia americalis, 2414. larva, 777, 1272. Helianthus. Lixus macer bred from, 2404. Heliothinæ. Synopsis of (review), 2178. Hagen, H. A. The Hessian-fly not imported from Heliothis armigera, 192, 636, 820, 993, 1136, 1256, Europe, 2267. Observations on certain North 1301, 1353, 1664, 1695, 1886, 1915, 2119, 2238, 2343. American Neuroptera, 39. Hagen's mystery. Dr., 1943. marginidens = Pyrrhia exprimens. phlogophagus, 936, 945, 993, 1056. Hair snakes, 612, 861, 1143. worms, 861, 1643, 1959. umbrosus, 2343.

Hellgramite, 1570, 1584.

Helops æreus, 751.

fly, 473, 1329.

Hairy caterpillar, 558.

grape-leaf folders, 664. Haldeman, S. S. Death of, 1909, Helops micans, 963. Hexaplasta, 1749, 1932. pullus = H. æreus. zigzag n. sp., 1749, 1932, 2343. Hemaris thysbe, 879. Hexapoda, 1955, 2267. Hemerobidæ, 2267. Poisonous, 2399. Hemileuca maia, 595, 722, 735, 760, 1329, 1352. Hibernating Aletia chrysalids. Supposed, 1927. Hemiptera, 400, 1329, 2267. apple-worms. Severe cold and, 2037. upon shrubs, fruits, and grains. Web-Hibernation of Aletia xylina in the United States ster, F. M., 2382. a settled fact, 2141. Experiments of the ef-Amphipyra pyramidoides, 1471. fects of punctures of, army-worm, 2086. the cotton-moth, 1728. Hemipterological studies, 2034. worm moth, 1953. Hemispherical larva at bottom of ant hill, 1789. insects, 1129. Hickory-bark borer, 938. Hemiteles, 77, borer, 269, 308, 1401. cressoni n. sp., 1059. Citheronia regalis on shell-bark, 775. fuscatus n. sp., 380 = var. of nemati-Eccopsis, 2363. vorus. Fall web-worm on, 460. nemativorus n. sp., 380. galls, 360. thyridopterygis n. sp., 1059. Hemlock Gelechia, 2232. made by Phylloxera. New, 1901. HEMLOCK. INSECTS AFFECTING. Gelechia. White heart, 2363. Hybrid between a grape and a, 1284. Buprestid, 2267. HICKORY. INSECTS AFFECTING. Cerambycid, 2267. Gelechia abietisella, 2232. Arhopalus pictus = Cyllene pictus. Cecidomyia tubicola, 1232. Hen. Death of a, 1094. Citheronia regalis, 775. Henderson's experiments. Mr., 1688. Clytus pictus = Cyllene pictus. Henous, 1643. Cyllene pictus, 89, 269, 308, 1516. confertus, 1600. Datana ministra, 2222. larval habits, 1387. Eccopsis, 2363. Heptagenia n.g., 39. Galls, 791. cruentata n. sp., 39.maculipennis n. sp., 39. Gelechia, 2363. simplex n.sp., 39.Hyphantria cunea, 460. Herbivorous ground-beetle. Another, 2042. textor = H. cunea. Herpetogomphus rupinsulensis n. sp., 24.Phylloxera caryæ avellana, 1901. Hesperid larva feeding on Canna, 1897. caulis, 360, 685. Hessian-fly, 150, 250, 426, 1581, 1665, 2008, 2267, 2365. globuli, 360. Effects of drought on, 2031. scissa, 1901. in England, 2395. Scolytus caryæ = S. 4-spinosus. into England. Introduction of, 2398. 4-spinosus, 938, 1329, 1401, 1754. Fighting the, 1587. Hickory Scolytus, 938. half way around the world, 2440. stem gall-louse, 685. Koeppen's account of the, 2267. vs. locust-borer, 1516. in North America. Early references Hipparchiscus n.g., 40 =Aplodes. to the, 2267. venustus n. sp., 40 = Aplodes miNot the, 1445. mosaria. imported from Europe. Hagen, Hippodamia. Ambiguons, 2119. H. A., 2267. convergens, 639, 1251, 1431, 1672. Parasites of, 2332. glacialis 849. prior to the Revolution. Insects conmaculata = Megilla maculata. founded with the, 2470. Hirmoneura. Eutomography of, 2275. Probable parthenogenesis of the, 1787. Larval stages and habits of, 2169. in seed-wheat, 494. obscura, 2169, 2275. Silesia. Cohn, F., 2267. Hirundo americana, 1502. Wheat-rust and the, 1605. fulva, 1502. Hetærina pseudamericana n. sp., 39. Hispa scutellaris = Odontota dorsalis. rupamnensis n. sp., 39.Hitherto unknown life-habits of two genera of rupinsulensis n. sp., 24. bee flies, 2002. scelerata n. sp., 39. Hockeria n.g., 11 = Haltichella. texana n. sp., 39. perpulchra n. sp., 11 = Haltichella per. Heteronychus relictus = Ligyrus relictus. pulchra. Heteropelma datanæ n. sp., 2526. Hog caterpillar of the vine, 1127. infested with para-Heteroptera, 1736.

> Hogs vs. bugs, 387. Holcaspis mamma, 518,

sites, 1247,

Defensive odors of the, 354.

bilineata, 372, 1043, 1851,

Hexagenia n. g., 39.

Holos around the roots of young ash-trees in the nursery, 471.

Holocera glanduella n. sp., 1310, 1311.

Homalomyia leidyi n. sp., 382.

pruinivora n.sp., 382. wilsoni n. sp., 382.

Home. Colorado potato-beetlo's native, 1462.

Homoptera. Egg-slits made by, 79.
Honoy-ant. Peculiarities of the Mexican, 1417.

bee. Braula cœca not particularly injurious to, 1982.

Bug preying on, 13.

Dipterous enemies of, 704.

Discussion on, 1455.

Insect cnemies of, 1059.

bees carnivorous? Are, 2098.

eating grapes, 441.

locust seed weovil, 1026.

weevil, 1474.

producing oak gall, 1942.

Too fond of, 764.

Honor. Deserved, 2124.

Hop Aphis, 1001, 2291, 2394, 2418.

and the cranberry, Smith, J. B. Insects affecting the, 2291.

growing in the West, 235, 279.

insects, 1001.

plant louse in Europe and America. Problem of the, 2400.

fully solved. Problem of the, 2396.

Life-history of the, 2393. Problem of the, 2400.

vine caterpillars, 444, 979.

HOP-VINE. INSECTS AFFECTING.

Grapta interrogationis, 420, 444.

Hypena humuli = H. scabra.

scabra, 979, 1001.

Hyperchiria io, 420.

Phorodon humuli, 235, 279, 1001, 2393, 2394, 2396, 2400.

Saturnia io = Hyperchiria io.

Vanessa interrogationis = Grapta interrogationis.

Hop-vine. Insect foes of the, 420.

Hoplophora arctata n. sp., 1363, 1370.

Hopper in Iowa, 1585.

Horinus lævis = Merinus lævis.

Horizontal insect boxes, 1963.

Hormaphis spinosus, 1678.

Horn-bug, 1636.

Cocoon of, 784.

Horned Passalus, 1311.

Egg of, 1329.

• Hornia n. g., 1601, 1643.

minutipenuis n. sp., 1601, 1651.

Structure and development of, 1651.

Horn's classification of the Carabidæ, 2051.

Horse bot-fly, 2238.

chestnut. Flat-headed apple-tree borer in,

Gregarious worms on, 1192.

HORSE-CHESTNUT. INSECTS AFFECTING.

Cacœcia rileyana, 1192.

Chrysobothris femorata, 1316.

Tortrix rileyana = Cacœcia rileyana,

Horse-hair snakes, 612.

Horses. Lico on, 258.

INDEX.

Horticultural ontomology. Recent advances in, 2230.

Horticulture. Utilization of ants iu, 2089, 2137.

Houghton's insects in the orchard, 188.

House-fly, 864, 2078.

Proboscis, 1783.

How great wits jump together, 567.

to hatch pupæ, 509.

Howard, L.O. Chinch-bug, 2418.

Codling-moth, 2418.

Howell, M. A. Experience with the spring canker-worm, 2267.

Hubbard, H. G. Miscellaneous notes on orange insects, 2164.

Report on cotton insects, 2343.

Rust of the orange, 2291.

Scale insects of the orange, 2119.

Hudson Bay Lepidoptera, 1985.

Hull's Curculio catcher, 372, 651, 875.

Hulst's observations on Pronuba yuccasella, Mr.,

Human animal. Parasites of the, 497.

body. Larvæ in, 382.

lungs. On a larva of Scenopinus from, 1348.

Humbert on Lucilia, 2255.

Humblebees, 800.

Humbug. Another, 195.

now Curculio, 589.

Entomology all a, 63.

Now, 122.

Curculio, 680.

Humming-bird moths caught by the tongue, 1388. Hundred and fifty million dollars. That, 1647.

legged worms, 219.

Hybrid between a grape-vine and a hickory, 1284.

Hydrachna belostomæ n. sp., 1632.

Hydrophilus piceus, 2016.

triangularis, 2016.

Notes on, 2016.

Hylecetus lugubris, 575.

Hylesinus opaculus, 1656, 1721.

trifolii, 1690, 1721, 1777, 1846, 2512.

Hylobius confusus, 700, 1168.

stupidus, 845.

Hylurgops, Packard, A.S. Development of, 2267.

Hylurgus pinifex, 2267.

Hymenoptera, 543, 1329, 1736, 2267.

Description of North American, 385.

Parasitic, 310.

Willow galls made by, 46.

Workers among, 311.

Hymenorus obscurus, 2105.

rufipes, 2105, 2226.

as a myrmecophilous spocies, 2226.

Hypena humuli, 9 = H. scabra.

scabra, 979, 1001, 1976, 2343.

scabralis, 2343.

Hyporchiria io, 420, 809, 1264, 1329, 1352, 1389, 2343.

varia = H. io.

Hypermetamorphoses of Meloidæ, 2191.

Hyphantria cunea, 112, 242, 454, 456, 460, 819, 1301, 1733, 1849, 1995, 2238, 2378, 2379, 2394,

textor = H. cunea.

Hypoderma bovis, 898, 1563, 2238 Imported carpet-beetle, 1699. Hyponomeuta, 2000. enrrant-worm, 1127, 1570, 2238. malinella, 2000. fly and its parasite, 389 multipunctella, 2000. 381. 5-punctella, 1603, 1804. elm-leaf beetle, 2232, 2304, 2325, 2378, Hypopus, 1703. Icerya. Bull. No. 15 on, 2389. gooseberry saw-fly, 228, 333. Life-history of, 2386. worms, 156. in New Zealand. Enemies of, 2479. insects, 140, 225. purchasi, 2386, 2394, 2400, 2401, 2415. and native American insects, an insect injurious to fruit-1115, 1127. trees, 2401. orchard Scolytus, 2233. Original habitat of, 2415. plants and insects, 1339. Importing European parasites, 208. sacchari, 2415. Improved method of spraying trees for protection Ichneumon, 1541. against insects, 2211. brevipennis, 1570. In memoriam B. D. Walsh, 1098. cæruleus, 1802. Cocoons of, 63, 183. Index to reports State entomologist of Missouri, flies, 662, 679, 1643, 1625. Cocoons of, 851. Indian-corn insects, 23, 138. Indiana, Butler, A. W. The periodical Cicada in fly. Army-worm, 1127. fly mistaken for a wasp, 477. southeastern, 2364. Webster, F. M. The season's observaleucaniæ, 1670. obsoletus n. sp., 1570 = var. of brevitions in, 2418. Indigo. Walshia amorphella on false, 1127. pennis. signatipes, 1802. Industry in the United States. Silk, 2268. from stomach of bluebird, 1878. Inexpert defense, 2476. wing, 50, 197, 385. Inflating Chalcis, 1059. Ichneumonidæ, 2516. Information wanted, 1111. Descriptions of new, 385. Injurious caterpillars, 151. Habits, 158. insects, 126. Ichthyura inclusa, 856. in California, 2074. Inquilines in galleries of common white ant, 1729. Icy lady-bird, 1311. Identity. Mistaken, 1593. and Psenides. Relations of, 41. Ignorance in the North. Entomological, 431. Inquiries answered, 180. South. Entomological, 390. Inquiring friends, 1642, 1656. Illinois. Entomological tour in Southern, 372. Insect boxes. Vertical vs. horizontal, 1963, First report noxious insects of, 373. catching habits of Sarracenia variolaris, Flying locusts in, 1437, 1443. 1385, 1390. Natural History Society. Address bechanges, 388. collection for sale, 2046. fore, 5. New Cynipidæ of, 41. collections. Naphthaline cones for, 2073. Pseudoneuroptera, 24. Protection of, 2180. A rare capture in, 1211. damage to the corks of wine-bottles, 2477. Six worst insect enemies of the fruitdefoliators. Shade-trees and their, 2378, growers in northern, 377. State entomologist of, 327. destroyer. Paris green as an, 1447. Horticultural Society. Report of eggs, 243, 286, 291. committee on entomology of, Growth of, 2241. 1056. enemies, 1583. Imbricated snout-beetle, 1301, 2291. of the Colorado potato-beetle, 411. Imitative butterflies, 613. growing rice, 1911. Impatiens fulva. Galls on, 852. to the rice-plant, 1949. . extinguisher. Treat's (review), 531. Imperial moth. Larva of, 1087, 1268. Importation of insect parasites, 2461. foe of the apple-tree borer, 421. Lestophonus. Further on the, to green corn. A new, 1655. 2541. foes of the apple-tree, 467. Imported cabbage-butterfly, 1127, 2291. army-worm, 12. worm, 2232, 2238, 2291. bark-louse, 417. Bacterial disease of the, hop-vine, 420. 2251. pea, 14. New remedy for, 2298. potato, 1558. in the South, 1714. found about orange-trees, 1798. Successful introduction friend. Another, 879. of a parasite of the, friends and insect foes, 38. 2291,

on grape, 1569.

Insect injurious to junipers, 1713.

wheat. A new, 2288.

killer. Worthlessness of the sparrow as

life, 5.

and sun spots, 2094.

locomotion, 1955.

named, 2297.

pest. A new, 1686.

plagues, 2198.

A plant growing out of an, 332, 478.

powder. Directions for cultivating pyre-

thrum for, 1996.

Persian, 1485.

powders and their use, 1692.

ravages, 1464.

Trade in, 1986.

world, 1466, 1467.

Figuier's (review), 408.

Remarkable peculiarity in the, 622.

Insecticide. London purple as an, 1725.

Oxeye daisy as an, 1861.

Pyrethrum an important, 2131.

its use as an, 2119.

Use of naphthaline as an, 2274.

Paris green as an, 1497.

Insecticides. Emulsions of petroleum as, 2134.

and their value as,

2126, 2200.

on garden insects, Alwood, W. B.

Tests with, 2382.

upon insects affecting garden crops.

Experiments with, 2344.

Quelques mots sur les, 2283.

Two valuable, 1743.

Insectivorous plants. Food of, 1499.

Insects attracted to light, 178, 2352.

Bill providing for the extermination of,

1842.

Birds vs., 199, 369.

Bluebirds feeding on parasitic and preda-

ceous, 1885.

in California. Injurious, 2074.

Californian orange, 2373.

Catalogues and monographs of, 1765.

Centennial, 1511, 1611.

clustered on apple-trees, 1263.

coflecting and preserving, 881, 1057.

Comstock's classification of, 2390.

Damage to silver plate by, 2154.

Death of mules caused by, 1811.

is directed. How flight in, 1891.

Directions for collecting and preserving,

881.

rearing, 14.

and drought, 2100.

effects of severe cold on, 1818.

enemies of fruit and fruit-trees. Review

of Trimble's, 187.

in northern Illinois. Six

worst, 377.

the honey-bee, 1059.

rice-plant, 1949.

feeding on sap of black-walnut, 1195.

in the flower garden, 936.

27 ENT

Insects as food for man, 2166.

found on apple-trees, 731.

Fungus diseases of beneficial, 1813.

Harmless, 1526.

Imported, 140.

plants and, 1339.

Improved method of spraying trees for

protection against, 2211. Injurious, 126.

injurions to agriculture. Legislation in

regard to.

1468.

New, 2055.

cereals and forage crops,

cotton in Brazil, 2277.

field crops, 2238.

fruit and fruit trees, 2238.

trees, 16, 31.

garden vegetables, 2238.

grape-vine, 267, 1081, 1086,

1091, 1102, 1118, 1127, 1130,

1158, 1175, 1202, 1221, 1255,

1281, 1301, 1311, 1329, 1363.

live-stock, 2238.

orange in Brazil, 2277.

sugar-cane in Brazil, 2277.

vegetation. Review of Har-

ris's, 568.

in Illinois, 6, 52.

Legislation to

control, 1946.

the vine, 2238.

of interest to fruit-growers.

Introduction and spread of scale, 2232.

by malodorants. Repelling, 2091.

Maple-tree, 2279.

Mind how you pack your, 435.

most destructive to the orange. Success-

ful management of the, 2088.

named, 163, 200, 316, 358, 359, 413, 447, 463,

490, 540, 560, 562, 582, 631, 673, 714, 721, 736,

743, 746, 785, 789, 805, 832, 847, 849, 858,

1140, 1147, 1183, 1226, 1273, 1287. to be named, 488, 514, 598.

The collection in the National Museum.

of, 2299.

Report on the collection of,

2282, 2330,

2368.

and native American insects. Imported, 1115, 1127.

Nervous system of, 2225.

Notes on our commoner, 1802.

on the oleander, 730.

in the orchard. Review of Houghton's,

Packard, A.S. Larvæ of injurions forest,

2267. Notes on forest, 2253.

Number of segments in the head of winged,

2267.

around peach trees, 660.

Pitcher-plant, 1385.

hordei, 563, 923, 2060, 2238, 2316, 2394.

tritici n. sp., 2060, 2063, 2119, 2123, 2316,

Larger wheat-straw, 2291.

lineare, 2060, 2119.

2363, 2394.

nigrum = I. hordei.

INDEX. Poisoning noxious, 205. Isosoma vitis, 482, 1059, 1127. Insects. Poisonous, 2399. Wheat, 2119. straw, 2291, 2394. Popular remedies for noxious, 73. Ithycerus noveboracensis, 16, 306, 652, 659, 869, 1033, Preserving, 881, 1057. 1085, 1188, 1301. Regulation of sex in, 1415. Iulus, 430. in relation to agriculture, 2238. Remedies for various, 2284. cœruleo-cinctus, 236. marginatus = Spirobolus marginatus. Retarded development of, 2040. Salt and vinegar for, 937. infested with Gamasus juloides, Simulium feeding on other, 2177. Some interesting, 1232. multistriatus n. sp.,193 = Cambala annulata. Stings of, 116. virgatus, 261. from stomach of lark, robin, and sun-Ixodes bovis, 404. fish, 1926. sp., 834. unipunctata, 1133. rock-bass, 1792. stripping the burr-oak, 966. Jaeger's North American insects, 84. in timber, 918. Japanese mode of packing silk-worm eggs, 1616. Trade in, 1986. Jarring down infested fruit. Codling-moth, 1318. Jassidæ, 737. Transformation of, 528. Jassus, 1766. Unity in coloration of, 50. Use of buckwheat to destroy, 1744. sexnotatus, 1766. Jiggers, 412. fungus growths to destroy, 1918. Johnson, L. Report on cotton-worm, boll-worm, poisons to destroy, 1887. used as food. Salt-water, 2203. and other insects, 2164. White willow, 907. Joint-worm, 49, 62, 563, 923, 2238. of the year, 2289, 2331. Appendix to article on, 1223. Destructive, 2322. fly, 1127. Joint-worms, 2119, 2394. Insekten. Einige unserer schädlicherer, 1325. Insidious flower-bug, 1127, 1423, 2418. Notes on, 2316. Instinct of Cicada septendecim, 2144. Joints of wheat. Worm in, 1848. Instructions to agents of the U.S. Entomological Jones, R. W. Observations and experiments on cotton-worm, 2164. Commission. Supplementary, 1888. Interaction of organisms, 139, 1954. Report on cotton insects, 2343. Interest felt in economic entomology in California, W. J. Report on cotton insects, 2343. Joppidium n.g., 385. Interim committees. Ad, 394. ruficeps n. sp., 385. Intermittance of phosphorescence in fire-flies, Journal of a State entomologist. One day's, 383. Jumping to conclusions, 253. Internal mite in fowls, 2157. seeds and galls, 1496, 2163, 2173. Introduction of Phylloxera. Laws to prevent the, spiders, 2302. sumach beetle, 1363. Invigorator again. Best's fruit-tree, 530. tree-cricket, 1329. once more. Best's, 545. Juniper. Dapsilia rutilans on, 1713, 1721. Io moth, 1329, 1389. web-worm, 1721. caterpillar, 809. Junipers. Insects injurious to, 1713. Iowa. Economic entomology in, 2197. Juniperus sabina. Insects injurious to, 698. The hopper in, 1585. Junonia lavinia, 753. Osborn, H. Report upon the insects of the Kansas bombardier-beetle, 1311. season in, 2418. The locust in, 1591. The seventeen-year Cicada in, 1737. and Missouri this fall. No locust injury Ipomæa commutata. Aletia feeding, 2343. in, 1433. Ips in calyx of pear. Banded, 1239. Report on grasshopper question to the fasciatus, 4, 214, 1239. governor of, 1573. 4-signatus = I. fasciatus. Scorpion in, 1119. Isabella tiger-moth, 1311. Silk culture in, 1542. Isosoma, 2394. Kartoffel-Käfer. Nene, 919. allynii, 2060, 2063. Katydid. Angular-winged, 1363. grande n. sp., 2288, 2291, 2316, 2348, 2363, Broad-winged, 1329, 1363. Domesticated, 1536. Habits of, 2348. Eggs of the, 516, 1005.

angular, 1518.

oblong, 569.

Narrow-winged, 1363. Oblong-winged, 1329, 1363,

See Catydid.

Katydids, 1363.

Kentucky. Coleopterous cave fauna of, 2033. Kermes galliformis n. sp., 1972. Kerosene emulsion, 2291. as a means against cotton insects, 2164. orange insects, 2164. Kiesenwetter. Obituary, 1819. Killer. Cottonwood, 1654. Killers. Bee, 1543. Killing apple-worms by machinery, 769. Kingdom. Animal, 393. Klippart's wheat plant (review), 186. Knot once more. Black, 930. Knots on apple-tree roots caused by root-lice, 1187. Knotweed geometer, 1059. Hæmatopis grataria on, 1059. Knowledge useless? Is any, 1135. Koebele, A. Experiments on cottony cushionscale, 2394. the red scale, 2394. Notes on locusts about Folsom, Cal., Report of experiments against scale insects, 2418. Koeppen, F. T. Account of the Hessian-fly, 2267. Labena. Useful, 1423. Labia, 2267. Lac insects, 2119 Lace-wing fly, 533, 1423. Lace-wing larva, 1059. Weeping, 1127. Lachnosterna fusca, 3, 54, 68, 300, 332, 410, 478, 500, 594, 640, 865, 1020, 1059, 1064, 1072, 1307, 1313, 1329, 1436, 1440, 1522, 1803, 1812, 2238, 2394. pilosicollis = L. tristis. quercina = L. fusca. quercus, 372. tristis, 966. Lachnus caryæ, 27. plantanicola n. sp., 2138. strobi, 265, 320, 1039. Lackey moth. American, 1301. Lacordaire, J. T. Death of, 1285. Ladder spider, 1299. Lady-bird. Ashy-gray, 2119. Blood-red, 2119. Cactus, 2119. Fifteen-spotted, 959, 1311. Icy, 1311. and its larva. Northern, 1289. Spotted, 599. Twice-stabbed, 38, 1883. Lady-birds, 1059, 1127, 1423, 1558, 2119. Swarms of, 824. Lafayette, Ind., Webster, F. M. Experiments at, .

Lagoa opercularis, 145, 796, 1748. Lake Superior. Simulium from, 2032. Lamellicornia, 1440. Lampronota, 1878. amphimilæna n. sp., 385.breviventris n. sp., 385. imitatrix n. sp., 385. interpellata n. sp., 385. pictiventris n. sp., 385.

Lampyridæ, 1705, 1805, 2036. Revision of the, 1819. Lampyris noctiluca, 1097, 1917. Laphria thoracica = Dasyllis thoracica. Laphygma autumnalis = L. frugiperda. frugiperda, 1127, 1256, 1267, 1282, 1301, 1400, 2011, 2343. fulvosa n. v., 1301 = var. of L. frugiperda. obscura, n. v., 1301 = var. of L. frugiperda. Laporte. Obitnary, 1817. Lappet caterpillars on the apple, 972. apple-trees, 812. Larch. Nematus erichsoni on, 2232. saw-fly, 2232. Large Asilus fly, 1269. black potato-bectles, 1206. compound gall on grape-vine, 720. dragon.fly, 759. fish-fly, 712, 903. gray straight-horned snout-beetle, 1033. green caterpillars on the apple, 1076. worms in a poach, 661. moth on apple-trees, 1028. phosphorescent larva, 1874. saw-fly, 1514. silken cocoon, 604. water-bectle, 750, 816. white-scale on Acacias, 1730. willow-worm, 1380. worm on apple-trees, 1048. Larger cabbage-butterfly, 2232. wheat-straw Isosoma, 2291. Lark. Insect from stomach of a, 1926. Larva boring along the axis of apple twig, 1850. injurious to cotton squares. Butterfly, 1872. Large phosphorescent, 1874. Moth issuing from a, 1779. Larvæ. Aquatic, 1851. Habits of, 346. in the human bowels, 382. of injurious forest insects, 2267. named, 1264.

Preserving, 1300. in stomach of black-bass, 1792. blue-bird, 1871. Larval characteristics of Corydalus and Chaulio-

des, 1652. characters and habits of blister-beotles, 1600.

Epicanta, 1600. Macrobasis, 1600.

habits of boe-flies. Bombylidæ, 1947, 1970. Dexidæ, 2260. Epicauta and Henous, 1387.

Sphenophori that attack corn,

life as influenced by food. Number of molts and length of, 2167. stages and habits of bee-fly. Hirmonenra,

Larviform females in the Phengodini. Luminous, 2397, 2402.

Lasins latipes, 27.

Lasioderma serricorne, 551. Lasioptera vitis, 720, 1329.

Law. Carrying out the, 1993. Lepidoptera of the Outer Hebrides, 2058. Lawn. Beetles swarming about, 718. Lepidopterological notes, 1999, 2013, 2160. Laws to prevent the introduction of Phylloxera, Lepidopteron. A myrmecophilous, 2214. Lepidopterous case-bearer, 260. larvæ. Dried leaves as food for, Leaf-beetle. Streaked cottonwood, 2291. bug. Ash gray, 1127, 1423. bugs, 400. Fleas feeding on, 2110. crumpler. Apple, 574, 1580. Remarkable, 40. in Georgia. Rascal, 1962. Lepidosaphidæ, 1301. Rascal, 341, 373, 1311. Lepiopomus pallidus. Insects from the stomach folder. Cranberry, 2291. of, 1792. Grape, 1301, 1579. Leptobatus illinoiensis n. sp., 385 = Exetastesgalls and caterpillars on sugar berry, 762. illinoiensis. on the grape-vine, 1716. Leptostylus aculiferus, 673. Leptura capitata, 200. injurious to small grains. A new, Leptus americanus n. sp., 1326 = Tetranychus.americanus. hoppers, 2362. irritans n. sp., 1326 = Tetranychus irrion celery, 452. of the grape, 399. Lesser apple-leaf folder, 1311. Grape-vine, 484. locust, 2363. injuring wheat, 1766. migratory locust, 2232. pine-borer, 2267. miner on white-oak, 1879. Prionus, 2267. miners of the locust, 451. Lestes inæqualis n.sp., 24. roller. Strawberry, 1574. Lencania albilinea, 1507, 1570, 1610. rollers. Descriptions of new, 1969. tyer. The green, 1311. unipuncta, 6, 7, 8, 9, 10, 11, 12, 17, 52, 328, 647, 670, 879, 906, 1127, 1267, Leafy oak-gall, 774. Leather-beetle, 2363. 1282, 1400, 1442, 1482, 1484, 1551, 1570, 1670, 1800, 1835, Lebia grandis, 365, 1218. 1856, 1877, 1885, 1886, 1953, Great, 1301. 2001, 2009, 2086, 2087, 2090, Lecanium, 1. acericola n. sp., 389 = Pulvinaria innu-2119, 2122, 2235, 2239, 2253, merabilis. 2262, 2267, 2343. macluræ n. sp., 389 = Pulvinaria innu-Additional notes on, 2001. merabilis. Complete life-history of, 1670. oleæ, 1303. Hibernation, 7, 8, 2086. rosæ = L. oleæ.Parasites of, 6, 10, 17, 1670. Leucopis, 154, 160, 174, 1530. sp. on blackberry, 117. cbina-tree, 1964. Phylloxera, 2150. Libellulidæ, 1707, 1709. magnolia, 1377. Library pest. Croton bug as a, 1717. plum, 107. sugar-maple, 1004. Lice on calves, 889. vitis = Pulvinaria vitis. horses, 258. Leconte, J. L. Death of, 2246. pigs, 900. snow-balls, 1184. Tribute to the memory of, 2264. Lichtenstein, J. Obituary, 2370. Leconte's pine-worm, 1570. Lichtenstein's theory as to dimorphic asexual fc-Lederer, J. Death of, 1285. Le Duc, W. G. Letter to, 1684. males, 2072. Life. Tenacity of, 1755. Legged maple-borer, 1363. Light. Insects attracted to, 178, 2352. Legislation to control insects injurious to vege-Lightning-hopper. Frosted, 1329. tation, 1946. hoppers, 737. in regard to insects injurious to agri-Lignified snake of Brazil, 2136. culture, 1468. Lignivorous. Rhyssa not, 2286. Lema trilineata, 119, 126, 135, 136, 138, 185, 401, 565, Ligyrus relictus, 3, 54, 68, 1440. 925, 1059, 1328, 1558, 1593. Length of larval life as influenced by food. Numrugiceps, 1794, 2080. ber of molts and, 2167. Lilac-borer, 844. the thread of the silk-worm, 1359. LILAC. INSECTS AFFECTING. Leopard moth. Great, 1311. Ægeria syringæ = Podosesia syringæ. Podosesia syringæ, 844. Lepidium vs. bed-bugs, 1741. Limacodes, 847, 1150. Lepidoptera, 1329, 2267, 2340. Hudson Bay, 1985. byalinus n. sp., 40 = Phobetron, hyali-Migrations of, 1770. pithecium = Phobetron pithecium. New lists of North American, 2132. Notes on South American, 1784. scapba, 40.

Limacodes tetradactylus n. sp., 40.

Limenitis disippus, 613, 1193, 1217, 1301, 1305, 1306,

1340.

ursula, 145, 1217, 1305, 1306, 1340.

Limneria fugitiva, 413, 1311.

lophyri n. sp., 1570.

pallipes, 2378.

Limnophilus, 2267.

Lina lapponica, 1834.

populi, 1834.

scripta, 1654, 1834, 2291.

tremulæ, 1834.

Linden and ash destroyers, 1849.

LINDEN. INSECTS AFFECTING.

Hyphantria cunea, 1849.

textor = H. canea.

Odontota rubra, 1849.

Linoceras junceus, 543, 1827.

L'insectologie agricole, (review), 506, 568.

Lintner's first report, 2244, 2254.

Linum. Acrididæ that eat, 1645.

Lissothoptrus simplex, 273, 1911, 1973, 2119.

Listotrophus cingulatus, 805.

Lithocolletis cincinnatiella, 1879.

guttifinitella. Mandible of, 1934.

Lithophane antennata, 1182.

cinerea, 1301.

Little Cicada, 1242.

known facts about well-known animals, 2071.

Turk and its crescent, 329.

Live-stock. Insects injurious to, 2238.

Lives. Two useful, 2370.

Lixus. Larval habits of, 2404.

macer, 2404.

parcus on Amelanchier, 2404.

Lobesia botrana = Eudemis botrana.

Loco weed. Walshia amorphelia bred from, 1127, 2356.

Locomotion. Insect, 1955.

Locust, 2366.

Ash-colored, 2363.

Atlantic migratory, 1423, 1625.

borer, 921, 1121, 1196, 1288.

Hickory vs., 1516.

borers, 37, 472.

Bruner, L. Observations in the North-

west on the Rocky Mount-

ain, 2165.

California migratory, 2363.

Californian, 1959.

Devastating, 2363.

Differential, 1423, 2363.

eggs, 1548.

in Asia Minor. Bombylid larvæ

destroying, 2118.

Condition of, 1567.

Experiments with, 1572. Expected advent of the, 2307.

experience, 1617.

flights, 1590.

in Dakota, 2007.

east of the Mississippi, 1549, 1613.

Geographical distribution of the Rocky

Mountain, 2267.

or grasshopper. Governors of Western States on the Rocky

Mountain, 1557.

Locust, important observations on the Rocky Mountain, 1571.

injuries. New method of counteracting, 2335.

injury in Kansas and Missouri this fall, 1433.

the near future. Probabilities of, 2355.

injury next spring, 1555.

summer. Anticipated, 1615.

LOCUST. INSECTS AFFECTING.

Arhopalus robiniæ = Cylleno robiniæ.

Cossus robiniæ, 37, 1121.

Clytus robiniæ = Cyllene robiniæ.

Cyllene robiniæ, 37, 472, 827, 921, 1196, 1250, 1288, 1516.

Hispa scutellaris = Odontota dorsalis.

Nitidulidæ, 37.

Odontota dorsalis, 451.

Spermophagus robiniæ, 1026, 1474.

Xyleutes robiniæ = Cossus robiniæ.

Locust invasion with the occurrence of drought.

Connection of, 1422.

Leaf miners of the, 451.

Lecture on the Rocky Mountain, 1493.

Lesser migratory, 2232, 2363.

mite, 1423, 1625, 1643, 1959.

in Montana in 1880, Bruner, L. Rocky Mountain, 2267.

multiplication and migration, Swinton, A. H. Solar physics and earthquake

commotion applied to, 2267.

Natural history of the Rocky Mountain,
1578.

notes. Miscellaneous, 2267.

in 1876, 1521.

1880, Martin, J. Rocky Mountain, 2267. 1885, Bruner, L. Abundance of the Rocky Mountain, 2363.

and other insects in the Northwest during the summer of 1883, Bruner, L. Observations on the Rocky Mountain, 2277.

Packard, A. S. Development of the, 2267. The periodical Cicada alias the seventeenyear and the thirteen-year, 1159.

pest, 1501.

Philosophy of the movements of the Rocky Mountain, 1669.

plague, 1473.

in the United States, 1625.

probabilities for 1882, 2057.

prospects, 1527, 1562, 1582.

in southwest Missouri this fall, 1596.

 ${\bf ravages~in~California,\,1959.}$

1880 and 1881. Chronology of, 2267.

Red-legged, 2363, 1423, 1625.

report to governor of Kansas, 1573.

Rocky Mountain, 1423, 1451, 1452, 1482, 1538, 1557, 1570, 1625, 1643, 1959, 2267, 2291.

scourge. Rocky Mountain, 1629.

seed-weevil. Honey, 1026.

Seventeen-year, 370, 884, 1489.

swarms that devastate the trans-Mississippi country, 1674.

theory wanted. New, 1532.

Locust. Two-striped, 1423, 2363. Louisiana. Destructive cricket in, 2384. weevil. Honey, 1474. Lozotænia rosaceana = Cacœcia rosaceana. in Wyoming, Montana, etc., in 1881, Bru-Lubber grasshopper, 2119. ner, L. The Rocky Mountain, 2267. Lucanus dama, 784. Yellow, 2363. elaphus, 305, 755, 957, 1517. Locustidæ, 929, 1453 Lucidota atra, 358. Locusts, 926, 1349, 1363, 2418. Lucilia macellaria, 209, 1880, 1921, 2158, 2199, 2255. again, 1550. Ludius attenuatus, 224. Bruner, L. List of North American, Lumbricus, 1304. Luminosity of fire-flies, 1840. Changes in vegetation caused by, 1495. Luminous larviform females of Phengodini, 2397, in Dardanelles. Sarcophaga lineata destructive to, 2075. Luna moth, 776. Destruction of young or unfledged, 1577. silk-worm, 1311. Destructive, 2363. Lungs. Larva of Scenopinus from human, 1348. Luperus brunneus, 1799. Dimorphism of, 1889. Ditching for young, 1488. morulus, 1799. eat the castor-bean, 1645. noxius = L. brunneus. about Folsom, Cal., Koebele, A. Notes Lure for moths, 1695. Lyda sp., 656. as food for man, 1481. Lydella doryphoræ n.sp., 1059 = Exorista dory-Grasshoppers and, 26, 147, 929. phoræ. Habits of young or unfledged, 1578. Lygus lineolaris = L. pratensis. pratensis, 2, 31, 76, 682, 1127, 1213, 1219, 2235, hatching? Are the, 1566. 2291, 2363. Mistaken identity. Are the, Lymexylidæ, 575. 1593. Lymexylon navale, 575, 1135. How to destroy, 1446. Lytta ænea = Pomphopæa ænea. in Illinois. Flying, 1437, 1443. atrata = Epicauta pennsylvanica. in Kansas, 1591. cinerea = Macrobasis unicolor. Literature of destructive, 1959. fabricii = Macrobasis unicolor. and locusts, 1994. marginata = Epicauta cinerea. in Nevada, 2024. murina = Macrobasis unicolor. New remedy against destructive, 2328. next spring. Injury by, 1555. sayi = Pomphopæa sayi. Notes on, 1456. tarsalis = Pomphopæa tarsalis. 1880. Chipman, A. J. Notes on, 2267. vittata = Epicauta vittata. Prairie fires and hateful, 1453. Machine. A satisfactory grasshopper, 1592. Machinery for destroying the cotton-worm, Barnard, W. S. Tests of, 2253. Professor Riley and the, 1434. of San Joaquin Valley, Cal. Coquillett, Killing apple-worms by, 769. D. W. Report on the, 2363. sting? Do, 371. McLain, N. W. Apicultural experiments, 2382. in Texas in spring of 1886, Bruner, L. Experiments in apiculture, 2394. Report on, 2382. Report on experiments in apiin the West, 2044. culture, 2363. and the Western cricket, Brnner, L. Maclura aurantiaca as food for Sericaria, 2234. Notes on other, 2267. Silk-worms fed on, 1341, 1542, in western Missouri. Ravages of young, 1609. McMurtrie, W. Tests of silk-fiber from cocoons 1492. Loew, H. Description of the rye gall-gnat, 2267. raised at the Department, 2253. Lombardy poplars. Caterpillars on, 571. Macrobasis, 1600, 1643, 1651, 2238. London purple as an insecticide, 1725. albida, 1796. and Paris green, 2021. Macrobasis, Larval characters and habits, 1602. Long-borned Diabrotica, 1905. murina = M. unicolor. legs. Grand-daddy, 838. unicolor, 38, 134, 185, 347, 362, 401, 470, sting. Delicate, 1329. 736, 912, 1044, 1209, 1558, 2248. tailed Ophion, 1311. Macrocentrus delicatus, 1334. Longicorn beetles. Food of, 1902. Macrodactylus subspinosus, 249, 361, 373, 565, 748, borers, 95. . 1075, 1278, 1329, 1357, 1375, 1478, 1583, 2248. Longicorns in pine and cedar, 319. Macrogomphus? spiniceps n. sp., 24. Longitarsus, 636. Macromia flavipennis n. sp., 24.

sting. Delicate, 1329.
tailed Ophion, 1311.

Longicorn beetles. Food of, 1902.
borers, 95.

Longicorns in pine and cedar, 319.

Look out for the eggs of the apple-tree plant-louse, 507.

Lopha 4-maculata — Bembidium 4-maculatum.

Lophyrus abbotii, 465, 927, 956, 1057, 1570.
abietis, 115.
lecontei, 985, 1011, 1057, 1570.

The still of the eggs of the apple tree plant-louse about it is in the eggs of the apple tree plant-louse.

The still of the eggs of the apple tree plant-louse, 507.

Macromia flavipennis n. sp., 24.
Macromia carolina — Protoparce carolina.
cluentius, 1784.
5-maculatus — Protoparce celeus.
rustica — Protoparce rustlca.

Mad. Entomology indeed run, 227, 1224.

Madarus ampelopsidos = Ampeloglypter ater.

vitis n. sp., 1059 =Ampeloglypter sesos-

Madras. *Coffee-borer in, 498.

Macklin, F. W. Obituary, 2161.

Maggots in sauce, 1607. .

Magnolia. Lecanium on, 1377.

Scale-insect on, 1377.

Maia moth, 1329.

Mails. Queen bees in the, 1762.

Maine. Now potato-bug in, 119.

MAIZE. INSECTS AFFECTING, 2119.

Achatodes zeæ, 1927.

Agrotidæ, 1522.

Elateridæ, 1522.

Lachnostorna fusca, 1522.

quercina = L. fusca.

Malaria. Mosquitoes vs., 2162.

Mallodon melanopus, 2291.

Mallophaga, 258, 497.

Mallophora orcina, 1959.

Malodorants. Repelling insects by, 2091.

Malva sylvestris. Erynnis alceæ boring in stems of, 1602.

Mamestra, 2238.

Cabbage, 2232.

chenopodi = M. trifolii.

picta, 179, 1056, 1127, 2232.

subjuncta, 2291.

trifolii, 229, 281, 2232.

Man. Effect of Paris green on, 1427.

Insects as food for, 2166.

Locusts as food for, 1481.

Mautidæ, 2267.

Mantis, 2267.

carolina = Phasmomantis carolina.

Preying, 457.

or rear-horse. Eggs of, 1060.

Supposed eggs of preying, 1002.

Mantispa, 2267.

brunnea, 1243.

Mautispian. Brown, 1243.

Many-banded robber, 1059, 1423.

Maple-bark lice, 1004.

louse, 344.

Boetlo on sugar, 1014.

borer. Legged, 1363.

A new (?) ægerian, 1360.

Eggs on sugar, 350.

Maple Galls on leaves of soft, 445.

MAPLE. INSECTS AFFECTING.

Acarus aceris-crumena, 1265.

Acronycta amoricana, 841.

Ægeria accrni, 743, 1063, 1360, 1363.

Anisota rubicunda = Dryocampa rubicunda.

Arachnid, 445.

Attacus eccropia, 841.

Catydid, 665.

Ccratocampa imperialis = Eacles imperialis.

Chrysobothris femorata, 1250.

Clisiocampa americaua, 350.

Clytus speciosus = Plagionotus speciosus.

Coccid, 344.

Dryocampa rubicunda, 841, 915, 1329.

Eacles imperialis, 1268.

MAPLE. INSECTS AFFECTING-Continued.

Eburia 4-geminata, 1014.

Lecanium acericola — Pulvinaria innumerabilis.

sp., 1004.

Sp., 1004. Orgyia antiqua, 1797.

Plagionotus speciosus, 915.

Pseudococus aceris, 1890.

Pulvinaria innumerabilis, 1515, 1816, 2279.

Sciara ocellaris, 2119.

Telca polyphemus, 841.

Trochilium acericolum = Ægoria acerni.

acerni = Ægeria acerni.

Maple. Mite-gall on sugar, 1265.

Ocellate leaf-gall of the red, 2119.

scale. Cottony, 1816, 2291.

Scale insect on, 1890.

troe insects, 2279.

twigs. Rows of eggs on, 665.

worm. Green striped, 1329.

worms, 841.

Maples. Cottony scale on, 1515.

Flat-headed borer in soft, 1250.

Mare's nest. Finding a, 86.

Margined blister-beetle, 1059, 1558.

Martin, J. Report on the Rocky Mountain locust, 1959.

Roport on the Rocky Mountain locust in 1880, 2267.

Mary Chalcis-fly, 1311.

Masicera archippivora n. sp., 1301.

Mason-bee cells. Remarkable new gonus of Me-

loidæ iufesting, 1601.

Mass of eggs, 240.

mistakes, 234.

Massachusetts. Periodical Cicada in, 2321.

southeast-

ern,2216.

Massospora cicadina infesting Cicada, 1809, 2112. May-beetle, 300, 865, 1020.

Egg of the common, 1329.

beetles swarming in Alabama, 1812. fly, 283.

Meadow enemy, 1368.

lark. Boetles in stomach of, 1013. worms, 947.

Meal sack. Worm infesting, 1896.

worms, 191.

Mealy hugs. Structure of, 2119.

Measuring-worms, 75.

Megastizus brevipennis, 375.

Megathymus coloradeusis n. var., 1602.

yuccæ, 1420, 1465, 1482, 1570, 1602. Additional notes, 1602.

Note on, 1465.

Megatoma serra, 1352.

Megilla maculata, 6, 94, 188, 599.

Food-habits of, 2145.

Melampsalta parvula, 1242.

Melancholy chafer, 522.

in apples, 842.

Melanactes, 1874.

Melanippe montanata, 2224.

Melanolestes picipes, 314, 1253.

Melanophila sp., 2267.

Melanophora? diabroticæ, 2260.

Microgaster, Military, 1127.

1264.

Notes on North American, 1960.

parasitic on Protoparce celeus, 155,

sp. parasitic on Thecla (poes?), 1872.

424 Melanoplus atlanis = Caloptenus atlanis. destructor = Caloptenus destructor. devastator = Caloptenus devastator. Melanotus communis, 351, 358. incertus, 40, 846. Melasoma lapponicum = Lina lapponica. populi = Lina populi. scriptum = Lina scripta. tremulæ = Lina tremulæ. Melissopus n.g., 1969. aurichalceana n. sp., 1969latiferreana, 1969. Melittia ceto, 125, 126, 248, 377, 1083. gloriosa 2410. Meloe, 1500, 1600, 2267. angusticollis, 1387. barbarus, 1600. proscarabæus, 2083. Meloidæ, 912, 1230, 1396, 1600, 1601, 1651, 1858, 2002, 2072, 2118, 2267. Fire cure for, 121. Hypermetamorphoses of, 2191. infesting potato, 48. Means against, 794. Remarkable genus of, 1601. Remarks on, 1600. Triungulin of, 2082. Meloini, 1601. Melolontha philophaga = Lachnosterna fusca. Melon. Bug on, 897. bugs. Satisfactory remedy for, 2236. Diabrotica vittata on, 897, 2236, 2238. Membracidæ, 737. Membracis ampelopsidis, 1183. Memoriam B. D. Walsh. In, 1098. Meracantha contracta, 1871. Merinus lævis, 1153. Merisus destructor, 1581, 2332. subapterus n. sp., 2332.Mermis, 2363. Meromyza, 1461. americana, 727, 1058, 1059, 1506, 1589, 1848, 1875, 2291, 2394. Mesochorus. Glassy, 1127. vitreus n. sp., 6, 1670. Mesoleius, 385. Metamymar n.g., 2343. aleurodis n. sp., 2343. Metapodius femoratus, 373, 775. nasulus = M. femoratus. Meteorus hyphantriæ n. sp., 2378, 2394. Methods of destroying scale-insects, 2119. Mexican honey-ants. Peculiarities of, 1417. Mexico. Report on cotton crop and its enemies in, 2343.

Mezinm americanum, 2346.

Microdon globosus, 1789.

2276.

Michigan apples and codling-moth, 1677.

1290.

Disippus, 1301.

limenitidis.

gelechiæ n. sp., 1059.

Life-habits of, 64.

Microgasters. Notes on, 2097. Microlepidoptera. Works on North American, Microplitis actnosus, 1125, 1134. ceratomiæ n. sp., 1125, 1134,1960. gortynæ n. sp., 1960. Micropterus salmoides. Larvæ in stomach of, Micropus leucopterus = Blissus leucopterus. Midas fulvipes n. sp., 40.Midge, 167. Clover-seed, 2291. Pear, 2363. Wheat, 109, 110, 142, 216, 280, 292, 428, 711, 1512, 2238, 2291, 2363. Migration of butterflies, 1770, 1991. and hibernation of Aletia argillacca, of plant-lice from one plant to another, 2017. Migratory butterflies, 1622, 1635. locust. Atlantic, 1423. Lesser, 2232. Military Microgaster, 1127. Milkweed. Danais archippus on, 1535. weevil, 711. Million dollars. That hundred aud fifty, 1647. Plums for the, 501. Mills. The same in flouring 1261. Milyas cinetus, 777, 1294, 1431, 1450, 1672, 2048. Mimicry and protective resemblances, 1301, 1340. Mind how you pack your insects, 435. Minings on apple-twigs. Galls and, 552. Minneapolis. Entomology at, 2212. Minnesota. A State entomologist for, 1108. Minot, C.S. Anatomy of Aletia, 2343. Minute borers in cherry, peach, and plum trees, 1940. Miscellaneous, 735. Miscellaneous notes on orange insects, 2164. Misnamed gall-moth, 1127. Mississippi country. The locust swarms that devastate the trans, 1674. Description of Aphididæ from west of the, 1678. floods. One effect of the, 2080. Locust flights east of the, 1549, valley. The Buffalo-gnat problem in the lower, 2416. Ailanthus silk-worm in, 1460. Missonri. Microcentrum retinervis, 241, 247, 1363, 1518, 1536, bee-killer, 1127. entomological reports, 1680. Noctuidæ in Microgaster, 77, 158, 183, 490, 662, 679, 717, 851, 1217, the, 2069. Entomology in, 1361. this fall. Locust prospects in southwest, 1596. No locust injury in Kansas limenitidis n. sp., 1301 = Apantelesand, 1433. Mnrtfeldt, M. E. Notes from, 2291.

Mossy rose-gall, 570, 1166.

Missouri. Murtfeldt, M. E. Notes for 1886 from, Oyster-shell bark-louse in, 1169. . Ravages of young locusts in western, Report State entomologist of, 1, 1059. 2, 1127. 3, 1301. 4, 1311. 5, 1329. 6, 1363. 7, 1423. 8, 1482. 9, 1570. Index, 2026. State Horticultural Society. Report committee on entomology, 1113,1127. Supposed bark-lice eggs in, 1084. Mistaken identity, 1593. Mistakes. A mass of, 234. Mite. On an extensile penetrating organ in a gamasid, 1626. in fowls. Internal, 2157. gall on sugar-maple, 1265. Locust, 1423, 1625, 1643, 1959. parasites of the Colorado potato-beetle, 1505. Phylloxera, 1363. Red, 1959. Scarlet, 1470. Silky, 1423, 1625. transformations, 1618. Mites, 356. Cannibal, 624. Classification of, 1992. Descriptions of new subterranean, 1370. Egg-infesting, 1637. on grasshoppers, 146. mistaken for plant-lice, 59. parasitic on beetles, 315. Transformations of the red, 1632. Mniszech, G. V. Ohituary, 2054. Mold and Phylloxera, 1807. Mole crickets, 1270. Molts and length of larval life as influenced by food. Number of, 2167. Monell, J. Notes on Aphididæ, 1678. Monocesta coryli, 1721. Monocrepidius vespertinus, 351. Monographs again, 1765, 1828. Monohammus titillator, 1324. Monophadnus rubi, 663, 761, 1212, 1641. Monostegia rosæ, 672, 984, 1780.

Most precious bug, 575. Moth on apple-tree. Large, 1028. Bee, 716, 904, 1059, 1508, 1723. hreeding. Pedigree, 2381. eggs, 611. Handmaid, 2222. issuing from a larva, 1779. named, 807, 1241. Viviparity of a, 2153. Motherless race. That fatherless and, 1650. Moths attracted by falling water, 2108. and hutterflies caught by the tongne, 1761. caught in Alabama, 1774. in cushions. Ravages of, 1731. Descriptions of two new, 1411. Galls made by, 1127. Lure for, 1695. mistaken for Aletia, 1976. by Physianthus albens. Capture of, 1388, 1419, 1761. Sngaring for, 1300. Mottled tortoise-heetle, 1127. Moulting in Orgyia. Variable, 2379. Monnding peach-trees, 587. again, 617, 653. Mountain ash. Ash and, 705. MOUNTAIN ASH. INSECTS AFFECTING. ? Aphis mali, 579. Aspidiotus harrisii = Chionaspis fnrfurus. Chionaspis furfurns, 579. Mountain ash. Plant-louse eggs on apple and, 579. Mnd-wasp and parasites, 1827. spider egg-nest, 1847. Mulberry. Sericaria mori on, 1311, 1341, 2234. silk-worm, 1311. Osage orange for the, 1220, 1286. for the silk-worm. Osage orange vs., 2234. Mulch hay. Worms under, 1161. Mules cansed by insects. Death of, 1811. Murgantia histrionica, 849, 1099, 1297, 1311, 1534, 2238, 2263, 2291. Murky ground-beetle, 486. Murmidius. Hahits of, 2217. Murmidius ovalis, 2217. Murtfeldt, M. E. Notes from Missonri, 2291. for the season of 1886,2382. Musca, 1390. domestica, 815, 864, 2071, 2078. harpyia = M. domestica.Muscid, 459. Montana in 1880, Bruner, L. The Rocky Mounfrom stomach of Ohio shad, 1853. Muscidæ, 153, 1880, 2256. 1881, Bruner, L. The Rocky Moun-Muscle shaped bark-louse on apple-trees South, tain locust in Wyoming and, 2267. Museum pest. Dinoderus pusillas as a, 2104. A new, 2109. once more, 492. Morning-glory. Coptocycla aurichalcea on, 1510. Trogoderma tarsale as a, 2139. Sphinx. White-lined, 630, 1198, 1301, 2291. pests, 246, 462, 572, 726.

again, 483.

Mutilla coccinea = Sphærophthalma occidentalis. Mycetophila persicæ n. sp., 653, 660, 963, 1056.

Mosquitoes, 603. vs. malaria, 2162.

More good words, 532.

Monstrosities. Rare, 2224.

tain locust in, 2267.

Monterey again. Butterfly-tree of, 2052.

universal remedies, 278.

Morus as food for Sericaria, 2234.

Mycetophila sp., 419. Necrological, 1817. Mychocerus, 2217. Neerophorus marginatus, 315, 845. Mygale hentzii, 466, 493, 521, 823, 1178. Nectarophora granaria, 126, 1806, 2394. and Pepsis formosa, 1619. rudbeckiæ, 27, 2205. Myochrous denticollis, 2418. Needle. Devil's darning, 1709. Myriapoda, 219, 224. Negro bug. Flea-like, 1127, 1423, 2291, 2418. Poisonous, 2399. Nematus erichsoni, 2198, 2232, 2355. Myriapods, 1625. fur n. sp., 197. Myrmecocystus hortusdeorum, 1942. hospes n. sp., 197.mexicanus, 1417. inquilinus n. sp., 197.Myrmecophila, 2202. integer, 2232. Myrmecophilous coleoptera, 2105. mendicus n. sp., 197, 821. lepidoptera, 2214. ribesii, 78, 140, 228, 333, 340, 364, 380, 381, species. Hymenorus rufipes as 756, 772, 1031, 1224, 1570, 1696, 2238. a, 2226. ribis = N. ribesii. Myrmica minuta, 1413. s.-desmodioides n. sp., 197. Myrmicocela ochracella, 2261. s.-pisum n. sp., 197. s.-pomum n. sp., 197. Mysia 15-punctata = Anatis 15-punctata. Mystery. Dr. Hagens', 1943. Spruce, 2232. in reference to Pronuba yuccasella, vontralis, 907, 917, 924, 1211. 1933. ventricosus = N. ribesii. Mytilaspis, 1. Nemestrinidæ, 2186. conchiformis = M. pomorum. Nemobius vittatus, 1287. pomicorticis n. sp., 1329 = M. pomo-Nemoræa leucaniæ, 6, 12, 647, 1112, 1127, 1670. Neoclytus capræa, 560,582, 1849. pomorum, 188, 201, 345, 372, 373, 377, 423, erythrocephalus, 1798. 565, 896, 944, 951, 973, 982, 989, 994, Nephelodes violans, 1885, 1990. 1007, 1008, 1012, 1040, 1056, 1059, 1084, Nephila plumipes, 830. 1169, 1246, 1329, 1362, 1397, 1552, 1583, Peculiarities of, 1416. 1774, 2088, 2238. Nepidæ, 797. My zus cerasi, 678, 960, 1067, 1207, 1251. Nerium oleander. Moths caught by, 1419, 1761. ribis, 304, 322, 735, 970. Nervous system of articulates. Yersin. Function Nails into fruit-trees. Driving, 87. of the, 1959. Nanodes tamarisci, 1496. insects, 2225. Nantucket. Pine-moth of, 2183. Phylloxera, 1687. Naphthaline concs, 2133. Neuroptera, 1329, 1736, 2267. for insect collections, 2073. North American, 24, 39. as an insecticide. Use of, 2274. Neuroterus q.-saltatorius, 1496, 2163, 2173. Napping. An entomologist caught, 1100. Nevada. Locust in, 2024. Narrow-winged katydid, 1363. New Curculio humbug, 680. National Museum. The collection of insects in, New England and New York. Packard, A. S. 2299, 2390. Causes of destruction of evergreen and forest trees in, 2232, Reports on the department 2291, 2363, 2364, 2382. of insects in the, 2282, 2330, Entomolog-2368. school of naturalists. ical speculations of the, 44. Native American insects. Imported and, 1115, entomological journal, 2056. 1127.apple-tree bark-lice, 1152. Hampshire. Colorado potato-beetle in, 1859. bark-lice on apple-trees, 1061. humbug, 122. currant-worm, 1570, 2238. Mexico. Blister beetles from, 1796. philosophy. Old and, 392. grape-vines. Cause of deterioration in some of our, 1342. York. Chinch-bug in, 2237, 2252, 2271. Colorado potato-beetle in, 1379. home. Colorado potato-beetles, 1462. plums. Wier, D. B., 2388. and Eastern States. Supposed army-Naturaliste Canadien. Le (review), 546. worm in, 1990. Naturalists. Entomological speculations of the Entomology in, 2244, 2254. New England school of, 44. Packard, A. S. Causes of destruction Observations and experiments, of evergreen and forest trees in New Neal, J. C. 2164. England and, 2232, 2291, 2363, 2364, Nebraska bee-killer, 1127, 1311. Bruner, L. Notes from, 2291. State entomologist for, 1863. weevil, 306, 652, 659, 1301. Report of the season's ob-

servations in, 2418.

insects, Bruner, L. Report on, 2382.

rufipes, 1363, 1367, 1703.

Necrobia ruficollis, 1363.

without a State entomologist, 1330.

News. Recent cotton-worm articles in the, 1719.

Nicotiana tabacum. Flea-beetle eating, 1782.

in apple trees, 1085.

Night. How the Curculio flies by, 706. Nursery. Holes around the roots of young ash Ninth report State entomologist of Missouri, trees in the, 471. 1570. Nuts. Gall, 1560. Niptus hololeucus, 2154. Nymphalidæ, 1301. Nitidula bipustulata, 214. Pupation of, 1704. Nitidulidæ, 37, 214. Nysius angustatus, 1329, 1357, 1402, 1453, 2048, 2291. Noctuid. A unique and beautiful, 2189. destructor n. sp., 1329 = N. angustatus. Noctuidæ, 455, 787, 1419, 1761. n. sp., 1317. of the Missouri reports, 2069. Oak apple, 1448. North America. Notice of Grote's bark louse, 159. iliustrated essay on the, 2148. borer, 2267. Parasites of, 349. Clytus, 2267. taken at Orono, Me., 2160. coccid mistaken for a gall, 1972. Nola sorghiella n. sp., 2119. fig gall, 745. Nomenclature. Scientific, 303, 768. gall, 212, 1925. None so blind as those who shut their eyes, on acorn cups. A new, 1606. Honey-producing, 1942. Nonsense about the Phylloxera, 1457. Leafy, 774. North America. Dilar in, 2023. OAK. INSECTS AFFECTING. Early references to the Hes-Amphibolips q.-inanis, 1448. sian fly in, 2267. q.-primus, 1606. Notes on Psyllidæ of, 2272. q.-spongifica, 1448. Packard, A. S. Zoo-geograph-Andricus q.-californicus, 1967. ical map of, 2267. q.-punctatus, 1822. Americau Anthomyidæ, 1966. seminator, 739, 1037. Coleoptera. Classification of, Balaninus rectus, 1311. 2085. Biorhiza forticornis, 745. Hymenoptera. Descriptions of, Cecidomyia q.-pillulæ, 799. q.-symmetrica, 799. Lepidoptera. New lists of, 2132. Cecidomyidæ, 799. Microgasters. Notes on, 1960. Cerambycid, 2267. microlepidoptera. Works on, Ceroptres ficus, 745. 1975. Coccus, 159. Entomological ignorance in the, 431. Cossus robiniæ, 1003. Northern army-worm, 1835. Cynipid, 212. brenthian, 1363. Cynipidæ, 41, 799. Illinois. Six worst enemies of fruit-Cynips fecundatrix, 1606. growers in, 377. forticornis = Biorhiza forticornis. lady-bird and its larvæ, 1289. gallæ-tinctoriæ, 1560. Northwest in 1883. Bruner, L. Rocky Mountain q.-californicus = Andricus q.-califorlocust in the, 2277. nicus. ou the Rocky Mountain locust. Bruq. decidua, 1925. ner, L. Report of observations in q.-frondosa, 774, 1606. the, 2165. q.-glandulus, 1606. Note, 184, 954. q.-inanis = Amphibolips q.-inanis. Notes. Entomological, 2, 1396, 1554, 1609, 1610, q.-mellaria, 1942. 1681, 1958, 2015, 2058, 2066, 2135, 2147, 2168, q.-prunus = Amphibolips q.-prunus. 2184, 2192, 2219, 2228, 2247, 2261. q.-punctata = Andricus q.-punctatus. by B. D. Walsh, 174. q. spongifica = Amphibolips q. spongi-Nothrus ovivorus, 1357. · fica. Notices, 196. seminator = Andricus seminator. Notodonta concinna = Œdemasia concinna. sp., 799. unicornis = Cœlodasys unicornis. terminalis, 1448. Nottiglossa undata = Oncometopia undata. Edema albifrons, 422. Noxious insects increase upon us. Why, 766. Elaphidion parallelum = E. villosum. named, 454. putator = E. villosum. Poisoning, 205. villosum, 288, 793, 2267. Popular remedies for, 73. Euclemsis bassetella, 1972. Nozzle. Cyclone, 2327. Galls, 1269. Nuisance. Caterpillar, 1995. Holocera glanduella, 1310, 1311. made useful, 983. Ithycerus noveboracensis, 1033. Number of entomologists in Europe, 1865. Kermes galliformis, 1972. molts and length of larval life as in-Lachnosterna fusca, 1812. fluenced by food, 2167. pilosicollis = L. tristis.

quercina = L. fusca.

tristis, 966.

segments in the head of winged in-

sects. Packard, A. S., 2267.

OAK. INSECTS AFFECTING-Continued. Œdipodini, 1959. Enothera grandiflora. Moths caught by, 1419, Lithocolletis cincinnatiella, 1879. Mallodou melanopus, 2291. Œstridæ, 114, 153, 290, 2256. Myrmecocystus hortus-dcorum, 1942. Œstrus hominis = Dermatobia hominis. My tilaspis pomicorticis = M. pomorum. pomorum, 973. ovis, 450, 887, 914, 1057, 1059, 1357, 1563, 2238. Ozognathus cornutus, 1967. Œta compta = Œ. punctella. Phylloxera lichtensteini, 1421. punctella, 1059, 1343. Tenthredinid, 108. Ohio insects. Alwood, W. B. Report on, 2382. Xylotrechus colonus, 2267. Oil-beetles, 1053, 1643. Oak. Insects stripping the bur, 966. leaf gall, 799. Old-fashioned potato-bugs, 185. and new philosophy, 392. Leaf-miners on white, 1879. question of species, 2201. . Phylloxera, 1363. Oleander. Chilocorus bivulnerus on, 730. pruner, 793. Coccid on, 218. and rose galls, 1037. Insects on, 730. tree borer, 1003. Olfersia americana, 2297. caterpillars, 422. Omaha. Conference of western governors at, twigs. Cynipid galls on, 1822. Woolly galls on white, 739. Oncideres amputator, 1556. OATS. INSECTS AFFECTING. cingulata, 442, 443, 476, 489,746, 848, 1556, Aphis avenæ = Nectarophora granaria. 1938. Nectarophora granaria, 1806. putator, 1556. Silvanus surinamensis, 1259. Oats and rye. Beetles working in wheat, 1259. Oncometopia undata, 36, 79, 164, 399, 1013. One day's journal of a state entomologist, 383. Oberea bimaculata, 302, 783, 1363. effect of the Mississippi floods, 2080. perspicillata = 0. bimaculata. half the vine area of France affected by Phylsp., 1059. loxera, 2020. ? boring in apple twigs, 1850. Onion-fly, 225. tripunctatá, 1503. Obituary. Bazille, L., 2370. Onion. Insects affecting. Anthomyia ceparum, 225. Belfrage, G. W., 2161. Chambers, V. T., 2218. Cambala annulata, 236. Iulus multistriatus = Cambala annulata. Clemens, B., 223. Ortalis flexa = Tritoxa flexa. Glover, T., 2218, 2258. Tritoxa flexa, 225. • Lichtenstein, J., 2370. Onion-maggot. Means against the, 12. Maeklin, F. W., 2161. Mniszech, G. V., 2054. pest. Chester, 2319. Putnam, J. D., 2054. Ontario. Index to the entomological reports of Oblong winged katydid, 1329, 1363. the province of, 2228. Eggs of the, 569, 1157. Onward march of the Colorado potato-beetle, Ocellate leaf gall of the red-maple, 2119. Odonata. Venation of, 39. Ophiogomphus mainensis n. sp., 39. Odontota, 2267. Ophion. Long-tailed, 1311. dorsalis, 451. macrurum, 1112, 1802. rubra; 1849, 1876. purgatum, 1442, 1670. Supposed eggs of, 1876. Purged, 1127. Odor in butterflies, 2025. Opsicœtus personatus, 497. Odynerus, 770, 1827. Orange. Bark-louse on osage, 389. birenimaculatus, 1827. in Brazil. Branner, J. C., insects injuriflavipes, 770, 1827. ous to the, 2277. Œcanthus, 2267. California. Scale-insects of the, 2389. latipennis n. sp., 2026, 2276. Experiments upon scale-insects affectniveus, 143, 251, 286, 414, 554, 723, 953, ing, 2164. 961, 999, 1006, 1059, 1122, 1323, in Florida. Voyle, J., the effects of cold on the scale-insects of the, 2277. 1329, 1333, 1503, 1691, 2195, 2238, 2276. Hubbard, H. G. Rust of the, 2291. ORANGE. INSECTS AFFECTING. Habits of, 207. Coccid, 218. Œcodoma fervens = Atta fervens. Œdemasia concinna 61, 413, 454, 779, 788, 922, Mite, 2291. Phytopus oleivorus, 2308. 1545. Orange insects. Californian, 2373. Œdipoda atrox, 1959. and cotton-worm, 2067. carolina = Dissosteira carolina. cruciata, 2075. Keroscne as a means against, obliterata, 1959. pellucida, 1959.

Miscellaneous notes on, 2164.

Orange insects. Successful management of the most destructive, 2088.

for the mulberry silk-worm. Osage, 1220, 1286.

Rust of tho, 2308.

Scale insects of tho, 2119.

seed. Worms in osage, 597.

Silk-worms fed with osage, 1341, 1542.

Successful management of insects most destructive to the, 2088.

trees. Insocts found about, 1798.

Orchard giving out. An, 578.

Injured, 1594.

Scolytus. Imported, 2233.

Houghton's insects in the, 188.

Orchelimum, 118, 861.

glaberrimum, 1329.

gracile, 447.

Oregon. Apple-plant lice in, 1899.

Organisms. Interaction of, 139, 1954.

Orgyia, 872.

antiqua, 1797.

Glands in, 2412.

leucostigma, 65, 480, 535, 600, 762, 1000, 1059, 1151, 1227, 1352, 2198, 2238, 2378, 2379, 2380.

moth issning from a larva, 1779.

Variable moulting in, 2380.

Oribatidæ, 1363.

Original habitat of Icerya purchasi, 2415.

Orocharis saltator, 513, 1287, 1323, 1329, 2276.

Ortalis flexa = Tritoxa flexa.

Orthocentrus pusillus, 385.

stigmaticus n.sp., 385.

trifasciatus n. sp., 385.

Orthoptera, 1329, 2267, 2276.

Orthoptera, Packard, A. S. systematic position of the, 2267.

Orthopterous insects. Packard, A. S., embryological development of, 2267.

Orthosoma brunneum, 95, 397, 2267.

Cylindrical, 1059.

cylindricum = 0. brunnenm.

unicolor = 0. brunneum.

Osage orange. Bark-louso in. 389.

OSAGE ORANGE. INSECTS AFFECTING.

Dipteron, 597.

Lecanium acericola = Pulvinaria innumerabilis

macluræ = Pulvinaria innumerabilis.

Pulvinaria innumerabilis, 389.

Sericaria mori, 1341, 1542, 1609, 2234.

Sphinx hageni, 2119.

Osage orange for the mulberry silk-worm, 1220, 1286.

seed. Worms in, 597.

Silk-worms fed with, 1341, 1542.

Sphinx, 2119.

vs. mulberry for the silk-worm, 2234.

Osborn, H., report of experiments at Ames, Iowa,

2344.

upon the insects of the season in Towa, 2418.

Oscinis, 2394.

brassicæ n. sp., 2291.

Oscinis. Cabbage, 2291.

Other insects affecting cheese, 1703.

Otiorhynchida injurious to cultivated plants, 2117.

Otiorhynchus picipes, 1788.

Our bugs, 2374.

table. On, 395, 408, 506, 546, 568, 623, 827, 1110.

Ont of evil cometh good, 618.

Outbreaks of army-worms. Recent, 2262.

Overcrowded, 654.

Overflow bugs in California, 2099.

Oviposition in Carabidæ. Mode of, 2341.

Prodoxus decipiens, 2049.

round-headed applo-tree borer, 2266.

Saperda bivittata, 1621.

Tortricidæ, 1922.

the Yucca moth, 1354.

Owlet moth. Spider-wort, 1301.

Owls' pellets, 601.

Ox insects, 290.

Oxalis. Aleurodes on, 1791.

Ox-oye daisy as an insecticide, 1861.

Oxyopes viridans, 2343.

Oxyptilus periscelidactylus, 1059, 1175, 1301.

Oyster-shell bark-louse, 373, 1059, 1127.

of the apple, 1329, 2238.

in Missouri, 1169.

Ozognathus cornutus, 1967.

Pachylobius picivorus, 845.

Pachypsylla n. g., 2208, 2272.

c. gemma n. sp., 2272.

c.-mamma n. sp., 2208, 2272.

venusta, 2208, 2272.

Pachytylus migratorins, 1959, 2044.

Pacific coast. Entomologist for the, 2022.

Whoat-stalk worm on the, 2123.

Pack your insects. Mind how you, 435.

Packard, A. S. Additions to the third report on the causes of the destruction of the evergreen and other forest trees in northern New England, 2364.

Canses of destruction of evergreen forests in New England and New York, 2232.

Descriptions of larvæ of injurious forest insects, 2267.

Development of the bark-boring beetles Hylurgops and Xyleborus, 2267.

Embryological development of Caloptenus atlanis, 2267.

Embryological development of Caloptenus spretus, 2267.

Embryological development of the locust, 2267.

Embryological development of orthopterous insects, 2267.

Fourth report on insects injuring forest and shade trees, 2382.

The Hessian-fly, 2267.

Narrative of the first journey in 1877, 1643.

a second journey in 1877, 1643.

Notes on forest-tree insects, 2253.

INDEX. Packard, A.S. Notes of a journey to Utah and Parasites, 66. Idaho in 1878, 1959. on hees, 1500. Number of segments in the head hred from cotton-worm, 1722. of winged insects, 2267. of Colorado potato-beetle. Mite, 1505. Second report on the causes of cotton-worm, 1712. the destruction of the evercut-worms, 349. green and other forest trees in Efficacy of Chalcid egg, 2115. northern New England and on eggs of Caloptenus spretus, 161 New York, 2291. fowls, 1308. Systematic position of the Orthe hateful grasshopper, 728. thoptera in relation to the other Hessian-fly, 2332. . Hog-caterpillars of the vine infested orders of insects, 2267. Third report on the causes of the with, 1247. destruction of the evergreen of the human animal, 728. Importing European, 208. and other forest trees in northon its larva. The Ahhot Sphinx, 1277. ern New England, 2363. Packard's Guide to the study of insects (Review), of the larva of Lachnosterna fusca, 2301. Leucania unipuncta, 6, 10, 17, 1670. 395, 479, 827. Pædisca celtisana n. sp., 1968. of the plum Curculio, 1751, 1795. giganteana n. sp., 1968. Synonyms of, 1932. obfuscata. Girdling habits of, 2405. Parasitic cocoons, 158, 1125, 1134. scudderiana, 2356. Not eggs but, 1290. Notes on, 2213. coleoptera, 2353. Painted borer, 89. diptera, 2076. lady-hird, 1329. flies, 310. Pale thighed tortoise heetle, 1127. fungus. Remarkahle, 1313. Paleacrita n. g., 1438. Parasitized tomato-worm, 155. vernata, 75, 86, 172, 173, 377, 1021, 1066, Paria aterrima, 1887, 1904, 2229. 1127, 1357, 1363, 1423, 1438, 1482, 1539, Paris green absorbed? Is, 1487. 1604, 1847, 2012, 2149, 2238, 2267. for cotton-worms. How to use, 1544. Palingenia bilineata = Hexagenia bilineata. the curculio, 1258. flavescens n. sp., 24. its effects on plants, soil, and man, pulchella n. sp., 24. terminata n. sp., 24. as an insect destroyer, 1447. vittigera n. sp., 24.insecticide. Use of, 1497. Pamphila ethlius, 1897. Paris green, London purple and, 2021. purple, 2021. Paniscus geminatus, 1059. Parorgyia. Glands in, 2412. Pangus caliginosus = Harpalus caliginosus. Synonyms and food-plants, 2412. Panorpa, 2267. Parsnip caterpillars, 453, 804. Panorpidæ, 2267. PARSNIP. INSECT AFFECTING. Paper-makers, 505. Paphia glycerium, 1117, 1127, 1329, 1704, 1711. Papilio asterias, 453, 804, 863, 1273. Parsnip worm, 863. Papilio, 1352. Parthenogenesis of the Hessian-fly. Probable, asterias, 453, 804, 863, 1272, 1276, 1512. 1787. cresphontes, 1237, 1300, 1914. glaucus, 28, 1048. in insects Occurrence of, 1029. of Mytilaspis pomicorticis, 1056. marcellus, 666. Passalus cornutus, 1311, 1329, 1636. philenor, 1127, 1321, 1774, 1957. thoas = P. cresphontes. Horned, 1311. troilus, 469, 857. Egg of, 1329. Passer domesticus as an insect destroyer, 1667. turnus, 28, 361, 1048. Parandra brunnea, 1733, 1876. PEA. INSECTS AFFECTING. Bruchus pisi, 434, 1120, 1301, 1916, 2238. Paraphia. Fir, 2363. Pararhyssa n.g., 385 = Rhyssa. Cerotoma caminea, 2418. Parasa chloris, 1233. Gryllus sp., 2383. Parasite of cabbage-worm, 2221. Pea and its insect foes, 14. canker-worm, 717. weevil, 1301, 2238. the imported cabbage-worm. Intro-Peach borer, 1017, 1059. duction of a, 2291. Flat-headed, 2267. Imported currant-worm fly and its, 380, horers, 871, 1366. Apple and, 1475, 1513. Mud-wasp and, 1827. Means against, 121. PEACH. INSECTS AFFECTING, 514. Phora not a, 1923.

on Prodoxus decipiens, 1831.

Sheep, 2043.

or a scavenger! Is Cyrtoneura a, 2102.

Ægeria exitiosa = Sannina exitiosa.

Callimorpha lecontei, 1301.

Asilus sp., 660.

populicanlis, 446, 713.

populi-monilis n.sp., 1678.

PEACH. INSECTS AFFECTING-Continued. PEAR. INSECTS AFFECTING-Continued. Carpocapsa pomonella, 1334. Eriocampa cerasi, 1031, 1382. Cetonia inda = Euphoria inda. Euphoria melancholica, 372. Dicerca divaricata, 871, 2267. Euryomia melancholica = Euphoria melan-Ephestia interpunctella, 325. cholica. zeæ = E. interpunctella. Ips fasciatus, 1239. Eriocampa cerasi, 1253. Lygus pratensis, 2, 31. Euphoria inda, 447. Mytilaspis pomicorticis = M. pomorum. Gortyna nitela, 632, 1210. pomorum, 973, 982. Notodonta concinna = Œdemasia concinua. Helops æreus, 751. pullus = H. æreus. Œdemasia concinna, 779. Larva, 661. Oncideres cingulata, 848. Lepidopteron, 697, 1210. Platycerus quercus, 1162. Lithophane antennata, 1182. Polycaon confertus, 2103. Mycetophila persicæ, 653, 660. Prionus laticollis, 561. sp., 419. Rhopalus sp., 372. Œcanthus niveus, 554. Scolytus rugulosus, 2233. Phleotribus liminaris, 2047. Selandria cerasi = Eriocampa cerasi. Sannina exitiosa, 587, 617, 871, 1017, 1059, 1070, Tremex columba, 928. 1475, 1513, 2238. Typhlodromns pyri, 1739, 1759. Scolytus rugulosus, 1940. Pear leaf blister, 1739. Selandria cerasi = Eriocampa cerasi. midge, 2363. Thyridopteryx ephemeræformis, 1189. root-borer, 561. Xyliua cinerea = Lithophane antennata. shoots. Bug gathering on, 741. Peach. Large green worm in a, 661. slug. Blood-sucker and, 1253. tree bark borer, 2047. Currant worm and, 1031. borer, 1070, 2238. tree borer. Elm and, 928. borers, 871. insect. A new, 1162. trees again. Mounding, 617, 653. insects, 277. Beetle around, 751. Rows of eggs in, 577. Insects around, 660. slug, 1382. Minute borers in, 1940. worms, 779. Mounding, 587. twigs. Girdled, 848. Sulphur cure for, 176. Two new foes of the apple and, 22. Supposed cause of yellows in, 515. Pearl wood nymph, 1127, 1301, 1363. twig borer, 632, 697. Pears. Curculios on, 1358. twigs. Eggs in, 554. Peas. Buggy, 1916. worm, 325. from bugs. To keep seed, 434. Pecan trees girdled by Oncideres cingulata. Blue spangled, 1301. boring into, 1182. Young, 489. Pear. Apple-bark lice on, 973. Pediculina, 258, 497. Banded Ips in calyx of, 1239. Pediculus capitis, 497. Bark-lice on the, 982. cervicalis = P. capitis. and cherry trees. Slng on, 1222. humanus = P. vestimenti. Diplosis, 2363. pubis = Pthirius pubis. vestimenti, 497. PEAR. INSECTS AFFECTING. Pedigree moth breeding, 2381. Anametis grisea, 2117. Pelargonium. Gali on, 1764. Anthonomus quadrigibbus, 1358. Pelidnota punctata, 40, 113, 129, 354, 358, 725, 1221, Aragnomus griseus, 2428. 1301. Aspidiotus harrisii = Chionaspis furfurus. Spotted, 1301. Capsus oblineatus = Lygus pratensis. Pelopæus cementarius, 543, 1371. Carabid, 692. Habits of Polistes and, 1371. Carpocapsa pomonella, 1334. lunatus = P. cementarius. Cerambycid, 288. Pempelia hammondi n. sp., 810, 1311, 1322. Chionaspis furfurns, 372. lignosella, 2119. Chloronoura malefica = Empoasca viridescens. Pemphiginæ. Biological notes on gall-making, maligna = Empoasca obtusa. Clisiocampa americana, 363. 1653, 1678. Conotrachelus cratægi, 1358. Pemphigini, 2017. Pemphigus acerifolii n. sp., 1678. nenuphar, 1358. Corimelæna pulicaria, 741. formicarins n. sp, 27. Diplosis nigra = D. pyrivora. formicetorum n. sp., 27.pyrivora, 2363, 2392. fraxiuifolii n. sp., 1678, 2361. Dolerns unicolor, 1989. imbricator, 449, 1396, 2361.

Empoasca obtusa, 22.

viridescens, 22.

Pemphigus populi-ramulorum n. sp., 1678.

Pezomachus, 158.

populi-transversus n. sp., 1678. Diminished, 1127. pyri = Schizoneura lanigera. minimus n. sp., 12, 1442, 1670.rhois, 518. Pezotettix, some species dimorphic forms of Caltessellata, 1948. optenus, 1889. ulmifusus n. sp., 518. pacifica n. sp., 1959.Phakellura nitidalis = Eudioptis nitidalis vagabundus, 446, 518. vitifoliæ = Phylloxera vastatrix. Phalæna zea, 2343. Penetrating organ in a gamasid mite. On an ex-Phalænidæ. Description of a new genus of, 1138. Phalangidæ, 838. tensile, 1626. Pennsylvania ground beetle, 1059. Phaneroptera curvicauda = Scudderia curvicauda. soldier beetle, 1059. Phasmida, 2267. Pentagenia n.g., 39. Phasmomantis, 457, 590, 859, 971, 979, 1059, 1060, quadripunctata n. sp., 39. Pentarthron n.g. (See p. 379) = Trichogramma. Phengodes, 1874, 2397. laticollis, 2412. Penthina fullerea n. sp. (See p. 378) = P. hebeand Zarhipis. Further notes on, 2411, vitivorana = Eudemis botrana. 2412. Notes on, 2408. Pepsis formosa, 466, 521, 543, 823. Phengodina. On the luminous larviform females Mygale hentzii and, 1619. Periclistus sylvestris, 1131. iu the, 2397, 2402. Perilitus indagator, 1311. Phenomena. Vegetable, 436. Pheocyma. Pine, 2363. Perimegatoma variegatum, 2109. Periodical Cicada, 474, 527, 619, 1059, 1311, 1489, 1624, Phemonone 5-caudata, 2410. 1836, 2014, 2312, 2314, 2315, 2363. Philampelus achemon, 1991, 1127. alias the seventeen-year locust, pandorus, 1102, 1127. satellitia = P. pandorus.1971, 1979. Philenor swallow-tail, 1127. alias the seventeen-year and Philonthus, 1311. thirteen-year locust, 1159. Belated individuals of the, 648. Philosophy of the movement of the Rocky Mountain locust, 1669. in Massachusetts, 2321. Notes on the, 2318. Old and new, 392. Our first brood established, 707. of the pupation of butterflies, 1711. Phleotribus liminaris, 2047. Premature appearance of the, 2320, 2326. Phlox. Heliothis phloxiphaga injuring, 936, 945. worm, 945. insavintwigs. Eggs of the, 698. Song notes of the, 2334. Phoberia atomaris, 1976, 2343. Phobetron hyalinum, 40. Periodical Cicada in southeastern Indiana. Butpithecium, 777, 1272. ler A.W.,2364. tetradactylus, 40. Massachusetts, Phora aletiæ, 1923, 1932, 2102, 2343. 2216. a scavenger, not a parasite, 1923. Twigs punctured by, 1055. Phorodon humuli, 235, 279, 1001, 2393, 2394, 2396, or seventeen-year Cicada, 2312, 2314. Perla decipiens n. sp., 24. 2400 Phorostoma latum, 2260. dorsata, 903. Phosphorescence of fire-flies. Intermittence of elongata n. sp., 24. the, 1805. flavescens n. sp., 24.fumipennis n. sp., 24 = Chloroperla fumithe glow-worm. Nature of the, 1810. pennis. producta n. sp., 24. Phosphorescent larvæ. Large, 1874. varians n. sp., 24. Photinus. Food of Calopteron and, 2358. Perlidæ, 2267. pyralis, 396, 1097, 1705, 1917, 2358. Perny silk-worm, 1311. Photuris pennsylvanica, 1097, 1705, 1917: Persian insect powder, 1485. Phoxopteris cornifoliana n. sp., 1968. PERSIMMON. INSECTS AFFECTING. fragariæ, 499, 984, 1059, 1574, 2324. Ceratocampa regalis = Citheronia regalis. murtfeldtiana n. sp., 1968. Citheronia regalis, 1275. nubeculana, 1721. Pests. Carpet, 1663. Phryganeidæ, 1138. Drug-store, 510. Phyciodes nycteis, 1410, Museum, 246, 462, 572. Phycita nebulella n.v., 1311 = var. of Acrobasis Potato, 1554. indiginella. of the strawberry, 2324. Phycita nebulo n. sp., 2 = Acrobasis indiginella. West. Winged, 2313. Phyllobrotica longicornis = Diabrotica longi-Petroleum as insecticides. Emulsions of, 2134. cornis. and their value as insecticides. Emul-Phylloptera oblongifolia = Amblycorypha oblonsions of, 2126, 2200. gifolia.

Phyllotreta albionica, 2291. Phylloxera vastatrix, 127, 373, 565, 724, 840, 916, striolata = P. vittata.1281, 1295, 1301, 1311, 1325, vittata, 348, 564, 2291. 1329, 1342, 1353, 1363, 1373, zimmermanni, 2291. 1376, 1394, 1403, 1421, 1423, Phylloxera, 1355, 1396, 1404, 1406, 1428, 1449, 1644, 1435, 1439, 1449, 1454, 1457, 1671. 1469, 1482, 1530, 1531, 1564, in California, 2140. 1623, 1671, 1716, 1721, 1727, Grape, 1727. 1732, 1808, 1841, 1900, 1906, 1907, 1952, 2015, 2019, 2020, Winged, 1895. caryæ avellana n. sp., 1901. 2140, 2150, 2208, 2250, 2283, -caulis, 360, 685, 1439. 2355. -fallax n. sp., 1423. Dipterous enemies of, 2150. -foliæ, 1311, 1363. Impregnated egg of, 1974. -globuli n. sp., 27, 360.vitifoliæ = P. vastatrix. \cdot gummosa, n. sp., 1423. work, 1900. ' -ren n. sp., 1423. Phyloptera, 2267. -scissa n. sp., 1901. Phyloscia, 1900. -semen, 373. Phymata erosa, 778, 1660. on chestnut, 353. Physianthus albens, 1418, 1419. congress in Spain, 1907. Capture of moths by, 1388, convention at Berne, 2147. 1419, 1761. correction, 1355. Peculiarities of, 1418. enemy. Grape-leaf, 1530. Physianthus. Insects affecting, 1388. in France. Grape, 2193. Deilephila lineata, 1419. with fungus disease. Infesting, 1808. Noctnidæ, 1419, 1761. in Geelong, Victoria. Grape, 1732. Sphingidæ, 1418, 1419, 1761. Grape, 1329, 1363, 1373, 1423, 1482, Physicians. Quacks or, 282. 1721. Physocnemum brevilineum, 696. and grape-rot, 1623, 1628. Physonota quinquepunctata n. sp. 767 = P. unipunctata. Half the vine area of France affected by, 2020. unipunctata, 767. laws, 2019. Phytocoridæ, 1301. in Europe, 2248. Phytolacca not fed on by Aletia, 2343. German, 2432. Phytonomus nigrirostris, 2355. opimus = P. punctatus. lichtensteni, 1421. mite, 1363. punctatus, 2010, 2028, 2061, 2119, 2355. Further notes on, 2061. Mold and, 1807. rumicis, 2119. Nervous system and salivary glands Phytophagic species, 45, 50. of, 1687. Phytophagic varieties, 45, 50. New biological facts regarding the Phytopus, 1739. grape, 1421. oleivorus, 2308. hickory galls made by, 1901. Newest facts of grape, 1404. Pickle-worm, 1127. Nonsense about the, 1457. Pieridæ, 613, 1301. not at the Cape. Grape, 1841. Dimorphic, 28. permanently destructive. The Pieris menapia, 2121. grape, 1907. monuste, 1991, 2232. oleracea, 835, 1127, 2232, 2238. notes, 1406. protodice, 836, 1127, 2232, 2238. Notes on the grape, 1952, 2291. rapæ, 1127, 1257, 1279, 1294, 1469, 1694, 1714, in the natural history of, 1439. 1720, 1721, 1867, 1948, 2221, 2232, 2238, Nouveau remède contre, 2283. 2251, 2298, Oak, 1363. into Alabama. Spread of, 1720. One-half the vine area of France afvernalis, 1634. fected by, 2020. Piesma cinerea, 398, 2048. quercus, 1363. Pigeon Tremex, 458. Remedies for, 1403. in apple, 1126. rileyi, 1363, 1423, 1530. Pigs. Lice on, 900. in sandy soil, 2250. Pimpla annulicornis, 2343. Some recent discoveries in reference annulipes, 1334. to, 2209. atrata = Thalessa atrata. spinosa, 2209. ceelebs n.sp., 385. in Switzerland. Extermination and conquisitor, 2343. restriction of, 2207.

inquisitor, 1059.

Use of guano for the grape, 1837.

Pimpla investigatrix n. sp., 385. Plant-feeding habit of Feniseca tarquinius, 2361. pictipes n. sp., 385. habits of predaceous beetles, 1956. pleurivinctus = P. conquisitor. growing out of an insect, 332, 478. Red-leggéd, 1329. sp. on Parandra brunnea, 1876. vidua n. sp., 385. Pine Amorbia, 2363. borer. Common longicorn, 2291. Lesser, 2267. Caterpillars on the, 927. and cedar. Longicorns in, 319. False caterpillars on the, 985. PINE. INSECTS AFFECTING. Amorbia, 2363. Asemum mæstum, 2267. Aspidiotus pinifolii = Chionaspis pinifolii. Borers, 1049. Callidium antennatum, 319. janthinum, 319. Chionaspis pinifolii, 97, 1329, 2079. Cybocephalus nigritulus, 2079. Hylobius confusus, 1168. Lachnus strobi, 320. Lophyrus abbotii, 465, 927, 956, 1570. lecontei, 985, 1011, 1570. Monohammus titillator, 1324. Pheocyma, 2363. Pieris menapia, 2121. Pissodes strobi, 781, 2363. Podapion gallicola, 2231. Ptinus brunneus, 191. Therina, 2363. Pine-leaf scale insect, 1329. moth of Nantucket, 2183. Pheocyma, 2363. Therina, 2363. trees. Butterfly larvæ injurious to, 2121. killed by borers. White, 1049. worm. Leconte's, 1570. White, 465, 956. Pinion. Ash-gray, 1301, Pinus elliotii. Chionaspis pinifolii on, 2079. inops. Gall on, 2231. strobus. Insects affecting, 97, 320. Pionea, Cabbage, 2232. rimosalis, 1867, 1915, 2232. Piophila casei, 1141, 1266, 1700, 1734. radicum, 1357. Pipiza femoralis, 495. radicum n.sp., 495 = P. femoralis. Pirates biguttatus = Rasahus biguttatus. picipes = Melanolestes picipes. Pissodes strobi, 781. PITCHER-PLANT. INSECTS AFFECTING, 1385. Sarcophaga sarraceniæ, 1385, 1390. Xanthoptera semicrocea, 1385, 1390. Pithy blackberry gall, 1131. galls on blackberry-twigs, 610. Platyptera, 2267. Pityophthorus micrographis, 295. minutissimus, 295. Plagionotus speciosus, 547, 915. Plagues. Insect, 2198. Planipennia, 2267.

Plant-bug. Dotted legged, 1311.

Tarnished, 1127, 1219, 2291.

feeders. Carnivorous propensity of, 1867.

lice, 23, 80, 265. again, 1008. Apple tree, 270, 335, 591, 996. on berberry, 644. Cherry-tree, 1067. Currant, 304. To destroy, 1148. Destroying cherry, 1207. enemies, 151. Evergreen, 1039. Mites mistaken for, 59. from one plant to another. Migration of, 2017. in Oregon. Apple-tree, 1899. Plum-tree, 678. on potatoes, 1679. Scab in apple vs. apple-tree, 1139. and their foes. Cherry, 1251. their friends and foes, 198. louse again. Eggs of the apple-tree, 550. Apple, 883. root, 373, 495, 1059. tree, 1127. Cabbage, 2291. Corn-root, 23. Currant, 322. eggs on apple and mountain ash, 579. Eggs of the apple-tree, 507, 539. enemy, 301. Grain, 99, 126, 1127. Look out for the eggs of the appletree, 507. White-pine, 320. Plants. Fertilizing, 324. Food of insectivorous, 1499. and insects. Imported, 1339. Otiorhynchida injurious to cultivated, soil and man. Effects of Paris green on, 1427. Platamodes unicolor, 573. Platycerus quercus, 1162. Platygaster herricki, 2332. Platyhypena scabra = Hypena scabra. Platynus æruginosus, 1013. excavatus, 1013. maculicollis, 2099. punctiformis, 1798. Platyphyllum concavum = Cyrtophyllus con-Platypsyllus castoris, 2417. as determined by the larva. Systematic relations of, 2417. Platyptilus carduidactylus, 1059. Plectrodes scalator, 1330, 1525. Plenty of specimens. Send, 593. Pleurotropis phyllotretæ n. sp., 2291. Plug-ugly theory, 233, 1383. Plum bark-louse, 107.

Curculio, 373, 1059, 1129, 1301, 2296.

breeds in the apple, 1214.

Plum Curculio, natural history and how to eatch Podosesia syringæ, 844. him, 1372. Poduræ, 1764. Parasite for the, 1751, 1795. Podura nivicola, 608. gouger, 33, 34, 373, 1301. Poduridæ, 317. gum. Larvæ in, 419. Pœcilloptera prumosa, 737, 1329. PLUM. INSECTS AFFECTING. Pœcilocapsus lineatus, 715. Acarid, 1265. Poison. Potato pest, 1524. Ægeria exitiosa = Sannina exitiosa. Poisoning noxious insects, 205. Anthonomus prunicida = Coccotorus scutel-Poisonous Arachnida, 2399. Is the Colorado potato-beetle, 1483. Aphis cerasi = Myzus cerasi. flour, 586. prunifoliæ, 678. Hexapoda, 2399. Attacus cecropia, 1249. insects, 2399. Bibio albipennis, 683. Myriapoda, 2399. Carpocapsa pomonella, 1334. qualities of the Colorado potato-beetle, Cerambycid, 559. Coccotorus scutellaris, 33, 373, 1301, 1378. qualities of the Colorado potato-bug, Colaspis sp., 372. 1101. Conotrachelus geminatus, 34. spider, 2306. nenuphar, 33, 372, 373, 501, 566, Tomato-worm not, 771. 1059, 1071, 1301, 1372, 1477, 1685, worm, 710. Poisons to destroy insects. Use of, 1887. puncticollis in sp., 34 = C. Polistes fuscatus = P. metricus. geminatus. metricus, 1371. Dipteron, 419. and Pelopæus. Habits of, 1371, Elaphidion parallelum = E. villosum. rubiginosus, 543. villosum, 606. Pollenia rudis, 2128, 2174. Elaterid, 81. Pollination of Yucca, 1603, 2000. Grapholitha prunivora, 373. Polycaon confertus. Habits of, 2103. Hylobius stupidus = Pachylobius picivorus. Polydesmus, 261. Lecanium sp., 107. complanatus, 215. Lyda sp., 656. serratus, 342. Mytilaspis pomicorticis = M. pomorum. sp., 834. pomorum, 973. virginiensis, 219. Myzus cerasi, 678. Polymitarcys alba, 1926, 1961. Pachylobius picivorus, 845. Polyphemus moth, 542. Phorodon humuli, 2393, 2394. Caterpillar of, 765. Sannina exitiosa, 81. Cocoon of, 1164. Scolytus rugnlosus, 1940. silk-worm, 1311. Semasia prunivora = Grapholitha prunivora. Polyrhabdus n.g., 385 = Chorinæus. Telea polyphemus, 765. cariniger n. sp., 385 = Chorinæus Plum leaf worms, 656. cariniger. moth, 373. Polysphineta nigriceps n. sp., 385. saw-fly, 2291. nigrita n. sp., 385. tree insects, 683. pimploides n. sp., 385. Minute borers in, 1940. Pomace-fly. Pretty, 2119. plant lice, 678. Vine-loving, 2119. twigs. Borer in, 559, 606. Pomphopæa ænea, 289, 301, 1056. Plume. Grape vine, 1059, 1301. sayi, 188, 301. moth. Spruce, 2363. Synoptic table of, 289. Plums and cotton, 1477. tarsalis, 289. . from Curculio. To protect, 1071. Pomphilus formosus = Pepsis formosa. for the million, 501. Poplar borer, 71. Preventing rot in, 1685. POPLAR. INSECTS AFFECTING. Wier, D.B. Native, 2388. Acronycta acericola = A. americana. Plusia brassicæ n. sp., 1127, 1867, 2232, 2238, americana, 575. Cabbage, 1127, 2232, 2238. Saperda calcarata, 1849. Poplars. Caterpillars on Lombardy, 571. dyaus, 2343. Popular delusion, 525. Plutella. Cabbage, 1291, 2232. . cruciferarum, 614, 1291, 1297, 2232. names, 100, 430. Pod-like willow gall, 1170. remedies for noxious insects, 73. Populus monilifera. Insects on, 1459. Podabrus modestus, 358, 673. Porcelio, 1900. rugulosus, 353. Podapion n. g., 2231. Porizon conotracheli n. sp., 1751 =Thersilochus

conotracheli.

Curculio parasite, 1301,

gallicola n. sp., 2231.

Podisus spinosus, 372, 456, 1378, 1431, 1450, 1672.

Potamanthus? odonatus n. sp., 24. Potato beetle, 946, 968, 995, 1019, 1027. abroad. The Colorado, 1364, 1407. again. Colorado, 1301. Colorado, 21, 334, 991, 1059, 1311, 1329, 1363, 1423, 1431, 1558, 1570, 2291, 2294 2418. enemies, 313, 365. in Europe. Colorado, 1597. New Hampshire. Colorado, 1859. Experience with, 1519. Inquiries concerning, 1491 Insect enemies of the Colorado, 411. Mite parasites of the Colorado, 1505. in New York. Colorado, 1379. The onward march of the, 1218. poisonous? Is the Colorado, 1483. Poisonous qualities of the Colorado, 1425. progress, 1480. Specific for the Colorado, 1490. Sweet, 1697. Ten-lined, 925. Three-lined, 1059, 1558. beetles, 247, 470. Driving, 1044. Fire cure for, 121. Large black, 1206. nativo home. Colorado, 1462. Safe remedies for, 1948. Some notes on, 1528. Sweet, 681, 1510. Potato-bug, 742, 905, 1450, 1672. Bogus Colorado, 519. Colorado, 293, 334, 366. in Maino, 119. New, 48, 88, 171, 293. Poisonous qualities of the Colorado, bugs, 401, 794, 1096. To destroy Colorado, 650. Old-fashioned, 185. and corn insects, 138. Insect foes of the, 1558. POTATO. INSECTS AFFECTING, 767, 1059. Acarina, 1444. Aphis solani, 1679. Baridius trinotata = Trichobaris trinotata. Cassida clavata = Coptocycla clavata. Ceresa bubains, 2107, 2235.

2081, 2294.

Epicauta cinerea, 401, 1056, 1558.

corvina, 1206.

puncticollis, 1381.

pennsylvanica, 38, 401, 1056, 1558.

Melanotus incertus?, 846. Mcloidæ, 48, 121, 794, 1059, 1230. Nysius sp., 1317. Protoparce celeus, 1059, 1558. 5-maculata, 401. Sciara sp., 239. Potato. pest poison, 1524. pests, 1558. Scab on, 239. worm, 1059, 1329, 1558. Potatoes. Plant-lice on, 1679. Scabby. 1444. Wire-worms in, 846. Pot-herb butterfly, 1127, 2232, 2238. Potter wasp, 770. Fraternal, 1127. Persian insect, 1485. Practical entomology in reality, 914. in, 1852. Prasocuris. Table of species, 156. varipes, 156. Coptocycla aurichalcea, 1510. Predictions verified, 1494. clavata, 693, 742, 1558. Crepidodera brevis, 1782. 2320, 2326. Preparation of Diptera, 2041. cucumeris, 401, 1056, 1059, 1558. parvula, 1782. Preserving insects, 881, 1057. Doryphora 10-lineata, 21, 48, 121, 138, 171, 293, 334, 347, 365, 366, 401, 565, 760, 905, 925, 946, 968, 995, 1019, 1027, 1044, 1059, 1218, 1301, 1311, 1329, larvæ, 1300. President. Washington Entomological Society. 1363, 1364, 1379, 1405, 1407, 1423, 1425, 1431, 1450, 1462, 1480, 1482, 1483, 1490, 1491, 1505, Address as, 2355. 1519, 1524, 1528, 1558, 1570, 1597, 1672, 1859, Presidential address to St. Louis Academy of

Sciences, 1564, 1629.

Pretty pomace-fly, 2119.

Preying Mantis, 457.

and unique gall-making tortricid, 2176.

Supposed eggs of, 1002.

POTATO. INSECTS AFFECTING-Continuod. Epicauta vittata, 401, 470, 794, 1056, 1230, 1558. Epitrix brevis = Crepidodera brevis. hirtipennis = Crepidodera parvula. Gortyna nitela, 401, 1059, 1558, 2229. Gryllus sp., 2383. Haltica cucumeris = Crepidodera cucumoris. Lema trilineata, 138, 401, 565, 925, 1059, 1558. Lepidopteron, 794. Lytta atrata = Epicauta pennsylvanica. cinerea = Macrobasis unicolor. marginata = Epicauta cinerea. murina = Macrobasis unicolor. vittata = Epicauta vittata. Macrobasis unicolor, 38, 347, 401, 470, 1044, 1558. Sphinx 5-maculata = Protoparce 5-maculata. Trichobaris trinotata, 401, 1059, 1558, 2172. Meloidæ injurious to, 48, 121, 794, 1059, New insect foe of the, 693. stalk-borer, 1059, 2172, 2229. weevil, 1059, 1558, 2291. Black blister beetles on, 1381. Buffalo tree-hopper injurious to, 2107. Powder. Manufacture of pyrethrum, 1984. Powders and their use. Insect, 1692. Prairie belt. First appearance of the cotton-worm fires and hateful locust, 1453. Predaceous bectles. Plant-feeding habits of, 1956. Premature appearance of the periodical Cicada Directions for collecting and, Prickly ash larva, 1914.

Papilio cresphontes on, 1914.

rose gall, 1194, 1245.

.PRIMROSE. INSECTS AFFECTING.

Alaria florida = Rhodophora florida.

Rhodophora florida, 1273.

Prionidæ. Food habits, 1902.

Prionidus cristatus, 262, 491.

Prionotus novenarius = Priouidus cristatus.

Priouus. Eggs of broad-necked, 1329.

imbricornis, 95, 1127.

laticollis, 95, 561, 1059, 1081, 1127, 1329,

2287.

Lesser, 2267.

sp. 736.

Tile horned, 1127.

Prisopus sp., 2267.

Pristiphora grossulariæ n. sp., 140, 176, 189, 364, 772,

1570.

sycophanta n. sp., 197.

Probabilities of locust or grasshopper injury in the near future, 2335.

Problem of the hop-plant louse in Europe and America, 2400.

fully solved, 2396.

Proboscis of the common house-fly, 1783.

Proconia undata = Oncometopia undata.

from stomach of Sturnella magna, 1013.

Procris. American, 1127.

americana = Harrisiana americana.

falsavius = A coloithus falsarius.

Grape vine, 213.

Protacanthus milberti, 1127, 1269, 1543.

missouriensis = P. milberti.

Proctrotrupidæ, 2343.

a new genus of, 1749.

Prodenia autnmnalis n. sp., 1301 — Laphygma frugiperda.

commelinæ, 850, 1282.

fulvosa = var. of Laphygma frugiperda.

lineatella, 1775.
obscura = var. of Laphygma frugiperda.

sp., 2238.

Prodoxidæ, 2000.

Prodoxus n. g., 1830, 2000.

ænescens n.sp, 2000.

cincreus n. sp., 2000.

 ${\tt decipiens}\, n.\, sp., 1804, 1830, 1854, 1933, 1943,$

2000, 2049.

mistaken for Pronuba yucca-

sella, 1804.

Oviposition of, 2049.

A parasite on, 1831.

Evolution of, 1804.

Further notes on Pronuba and, 2000.

remarks on the difference between Pronuba and, 1854.

intermedius n. sp., 2000.

marginatus n. sp., 2000.

Pronuba vs., 1844.

Structural and anatomical peculiarities of 2171.

Profits of silk-culture, 2338.

Promachus, 1127.

apivorus, 668, 1059, 1543.

bastardii, 1269.

Promachus vertebratus, 1269.

Promethea silk-worm, 1311.

Prominent. The red-humped, 2291.

unicorn, 1280.

Pronuba n. g., 1329, 1336, 2000.

and its connection with the pollination

of Yucca, 2391.

Evolution of, 1804.

 $\mathbf{maculata}\; n.\; sp., 2000.$

A new tineid genus allied to, 1830.

and Prodoxus. Further notes on, 2000.

remarks on the differences be

tween, 1854.

Structural and anatomical peculiarities of, 2171.

vs. Prodoxus, 1844.

yuccasella n. sp., 1329, 1336, 1337, 1363, 1603, 1854, 1933, 1943, 2000,

2171.

Chrysalis of, 1414.

Further remarks on, 1603.

Mr. Hulst's observations on, 2371.

mistaken for Prodoxus decipiens, 1804.

Mystery in reference to, 1933.

Supplementary notes on, 1337.

Prophecy fulfilled, 709.

Proscopia, 2267.

Prosopis affinis, 1171.

Prospects. Locust, 1527, 1562, 1582.

Protection of insect collections, 2180.

Protective device employed by a glancopid caterpillar, 2242.

resemblances. Mimicry or, 1301, 1340.

Proteoteras n.g., 1968.

æsculana n. sp., 1968, 2114, 2206.

Protoparce carolina, 131, 933, 1089, 1262, 2238.

celeus, 131, 152, 401, 490, 710, 815, 851, 885, 993, 1059, 1089, 1264, 1329, 1558, 1661.

Prunus insects, 1503.

Psenides, 821.

and inquilines. Relations of, 41.

 ${\bf Psenocerus\ supernotatus,\,481,\,1458.}$

Pseudococcus aceris, 1890.

Pseudouenroptera, 2267.

of Illinois, 24.

New, 24, 39.

Pseudoscorpion, 1884.

Psinidia wallula n. sp., 1959.

Psoci, 2133.

Psocidæ, 1147, 2267.

Psocina. Structure of the claw in, 2045.

Psocus, 1078.

amabilis n. sp., 24.

bifasciatus n. sp., 39.

confluens n. sp., 39.

conterminus n. sp., 39.

geologus n.sp., 24.

lichenatus n. sp., 39.

madescens n. sp., 39. permadidus n. sp., 39.

perplexus n. sp., 24.

pollntr s n. sp., 24.

Psocus purus n. sp., 24. Pyrethrum willemoti, 2343. rufus n. sp., 39. Pyrophila conspersa, 1301. semistriatus n. sp., 24. pyramidea, 785, 1301. pyramidoides, 671, 785, 1301, 1471. venosus, 176, 1078, 1147, 1195, 1263, 1526, 1798. Hibernation of, 1471. Psyche confederata, 1441. Pyrrharctia isabella, 244, 558, 909, 1153, 1311, 1802. Psychidæ, 1650. Pyrrhia exprimens, 677. Psychomorpha epimenis, 1301, 1363. Quackery. Entomological, 439. Psylla. Box, 2291. Quacks and physicians, 282. rubi, 669. Quedins, 1311. Psyllid galls. Hackberry, 2208. Queen bees in the mails, 1762. Quelques mots sur le insecticides, 2283. Psyllidæ of North America. Notes on, 2272. the United States, 2210. Quercus acuminata. Cynips on, 1822. coccifera. Phylloxera on, 1421. douglassi. Gall on, 1967. Pteromalus puparum, 2221. Pteronarcys, 2267. Pterophorus carduidactylus n. sp., 1059. Galls on the prinus group of, 1606. infectoria. Gall on, 1560. periscelidactylus = Oxyptilus periscelidactylus. muhlenbergi. Cynips on, 1925. palustris. Coccid on, 1972. undulata. Galls on, 1942. Pthirius pubis, 497. Ptinus brunneus, 191, 510, 1141. Publication. Discontinuance of, 2106. Queries answered, 217. Publilia concava, 163. Bundle of entomological, 1463. Questions answered, 1246. Puccinia graminis, 1605. Pulex irritans, 497. Quick traveler, 752. Quince Curculio, 1301. penetrans = Sarcopsylla penetrans. QUINCE. INSECTS AFFECTING. sp. feeding on lepidopterous larvæ, 2110. Capsus oblineatus = Lygus pratensis. Pulvinaria acericola = P. innumerabilis. Conotrachelus cratægi, 1301. innumerabilis, 389, 1515, 1816, 2279, 2291, Eriocampa cerasi, 2243. macluræ = P. innumerabilis. Lygus pratensis, 2. vitis, 106, 1212. Selandria cerasi = Eriocampa cerasi. Punctured grape-canes, 513, 1333. Quince. Saw-fly on the, 2243. Rabbit insects, 290. Punctures of Hemiptera upon shrubs, fruits, and grains. Webster, F. M. Experiments on the Race. Some further facts regarding that fathereffects of, 2382. less, 1658. Punctures on rose twig, 1167. That fatherless and motherless, 1650. Pupa of the disippus butterfly, 1193. RAGWEED. INSECTS AFFECTING. Pupæ. Habits of, 346. Epicauta pennsylvanica, 732. How to hatch, 509. Lytta atrata == Epicauta pennsylvanica. Pupation of butterflies. Philosophy of the, 1711. Railroad worm, 2238. Nymphalidæ, 1704. Rauatra, 1709. Purged Ophion, 1127. fusca, 797, 1191. Purple willow Gracilaria, 2363. Range of the rear-horse, 859. PURSLANE. INSECT AFFECTING. Ransom chip-trap for Conotrachelus, 1177. Deilephila lineata, 1402. Ranuuculus acris. Insects affecting, 156. Putnam, J. D. Obituary, 2054. Curculio remedy, 1201. Pyralid infesting seed-pods of trumpet-vine, 1929. Rapacious soldier-bug, 1059. Pyralis farinalis = Asopia farinalis. Rape butterfly, 836, 1294. olinalis = Asopia olinalis. our new cabbage pest, 1257. Pyrameis atalanta, 540. Rare capture in Illinois, 1271. cardui, 1635, 1770. monstrosities, 2224. Pyramidal grape-vine worm, 1301. Rasahus biguttatus, 497. Pyrethrum. Additional experiments with, 1903. Rascal leaf-crumpler, 341, 373, 1311, 1580. cinerariæfolium, 1996, 2235. in Georgia, 1962. Cultivation of, 1984. Raspberries. Do bees injure, 529. Directions for cultivating, 1996. destroyed by weevils, 1788. raising, 1862. Raspberry borer, 783. Experiments with, 1948, 2353. brand, 676. an important insecticide, 2131. canes dying, 1035. powder. Manufacture of, 1984. Eggs of snowy tree-cricket in, roseum, 1996, 2119, 2343. for the screw-worm, 1921. tree-cricket in, 1006. in the United States. Additional Gouty gall on blackberry and, notes on the cultivation of, 2377. The use of, 1858. punctured by Orchelimum glaits use as an insecticide, 2119. berrimum, 2249.

Raspberry Geometer, 1059.

gouty gall, 1124.

and grape-vines. Egg punctures in, 2195.

RASPBERRY. INSECTS AFFECTING.

Ægeria rubi = Bembecia marginata.

Agrilus ruficollis, 1124, 1771.

Bembecia marginata, 1363, 1453.

Chionaspis furfurus, 1968.

Diaspis harrisii = Chionaspis furfurus.

Dipteron, 676.

Monophadnus rubi, 663, 761, 1212, 1641.

Oberca bimaculata, 783, 1363.

perspicillata = 0. bimaculata.

tripunctata, 1503.

Œcanthus niveus, 961, 1006, 1122, 1503, 1691,

Orchelimum glaberrimum, 2249.

Otiorhynchus picipes, 1788.

Rhodites radicum, 1149.

Selandria rubi = Monophadnus rubi.

Synchlora rubivoraria, 1059.

Raspberry-root borer, 1363.

gall, 1149.

saw-fly, 1212, 1641.

Scale insect on, 1968.

worms, 663, 761.

Rat-tailed larva, 168.

Ratzeburg. Forest trees and weeds of Germany (review), 1110.

Rear-horse, 1059.

Eggs of the Mantis or, 1060.

Range of the, 859.

Rear-horses vs. grasshoppers, 590.

Record of American entomology (review), 827.

Red ants, 2309.

cedar caterpillars, 427.

clover. Bumble bees and, 2376.

humped caterpillar, 788.

prominent, 61, 2291.

leg. The cussed, 1598.

legged ham-heetle, 1363.

locust, 1423, 1625, 2363.

maple. Ocellate leaf-gall of the, 2119.

mites, 1959.

Transformation of the, 1632.

scale. California, 2394.

Koehele, A., experiments on the, 2394.

shouldered Sinoxylon, 1311.

spider, 1146, 1225, 1710.

on roses, 1710.

tailed Tachina fly, 1127.

weevil, 711.

Reddish snout heetle on apple. Small, 1244.

yellow spruce bud-worm, 2291.

Reduvius personatus = Opsicætus personatus.

raptatorum = Sinea diadema.

Regal walnut caterpillar, 425.

Regulating sex in insects, 1415.

Remarks, 206, 255.

Remedies and appliances, 2375.

More universal, 278.

Universal, 204, 503, 818.

Remedy. Another universal, 232.

Report of committee of cntomology, Ill. State Hortic. Soc., 1056.

Report of committee of entomology, Mo. State Hortic. Soc., 1113, 1127.

> experiments, chiefly with kcrosene, upon insects injuriously affecting the orange-tree and cotton-plant, 2164.

to governor of Kansas on the grasshopper question, 1573.

on Lucilia macellaria, 2199.

of State entomologist of Illinois, 1, 373.

Missouri, 1, 1059.

2, 1127.

3, 1301.

4, 1311.

5, 1329.

6, 1363.

7, 1423.

8, 1482.

9, 1579.

Index, 2026.

U.S. Ent. Commission, 1, 1643.

2, 1959.

3, 2267.

4, 2343.

U. S. Entomologist for 1878, 1721.

1881-'82, 2119.

1883, 2232.

1884, 2291.

1885, 2363.

1886, 2394.

1887, 2418.

Reports. Missouri entomological, 1680.

U.S. Entomological Commission, 1757.

Resemblances. Mimicry or protective, 1301, 1340. Resistance of grape-vines to Phylloxera in sandy soil, 2250.

Retarded development in a blister-heetle, 1860.

of insects, 2040.

Remarkable case of, 2003,

Retirement of Mr. Fuller, 1910.

Revision of the Lampyridæ, 1819.

Rhagium lineatum, 2267.

Rheum rhaponticum. Asilus sericous feeding on, 1543.

Rhinoceros heetle. Gigantic, 580, 1216, 1292.

Rhinopsylla n. g., 2272.

schwarzii n. sp., 2272.

Rhipiphorus sp., 2301.

Tiphia and, 2351.

Rhizotrogus sp., 2260.

Rhodites, 1235.

hicolor, 1037, 1194, 1245.

radicum, 137, 1149.

rosæ, 570, 1166, 1939.

Rhodobænus 13-punctatus, 1301, 2119.

Rhodophora florida, 807, 1273.

Rhopalocera, 1635, 1784.

Rhopalosiphum, 1678.

Rhopalus, 372.

Rhus glabra. Insects on, 320. Rhynchophora, 254, 1301.

Rhynchophorus zimmermanni, 2119.

Rhyssa, 385.

atrata = Thalessa atrata.

not lignivorous, 2286.

lunator = Thalessa lunator.

Ribes insects, 772. Rice crop. Enemies of the, 273. grub, 2119. RICE. INSECTS AFFECTING. Acridium obscurum, 2119. Cecidomyia oryzæ, 1949.

Chalepus trachypygus, 273, 1911, 1949, 1973, 2119.

Chilo oryzæellus, 2119, 2120.

Laphygma frugiperda, 2011.

Lissorhoptrus simplex, 273, 1911, 1973, 2119.

Murmidus ovalis, 2217.

Rice. Insects affecting stored, 2190.

enemies of growing, 1911.

plant. Another enemy of the, 2011. Insect enemies of the, 1949. Insects affecting, 2119. Water weevil of the, 1973.

stalk borer, 2119.

New, 2120.

weevil, 1683.

Ricinis communis. Acrididæ feeding on, 1645. Ridding the ground of cut-worms, 2310.

1. iley to Dr. Schaffer. Entomology. Professor,

in favor of the birds. Professor, 1675. Letter from Prof. C. V., 1676. and the locusts. Professor, 1434.

Yucca moth. Professor, 2068.

Riley's report to the governor of Kansas. Professor, 1573.

researches, 2067.

Ring-banded soldier-bug, 1311.

legged Pimpla, 1329.

Robber. Many-banded, 1059, 1423.

Robin. Cut-worms from the stomach of, 1873.

Insects from the stomach of, 1926.

Rock-bass. Insects from stomach of, 1793.

Rocky Mountain grasshopper, 1557.

locust, 1423, 1451, 1452, 1482. 1538,

1557, 1570, 1625, 1643, 1959, 1998, 2267, 2291.

Bruner, L., observations in the Northwest on the, 2165.

in 1880. Martin, J., on the, 2267.

Bruner, L., re-1885. porton the abundance of the, 2363.

Geographical distribution of the, 2267.

or grasshopper. Governors of Western States on the, 1557.

Important observations on, 1575.

Lecture on the, 1493.

in Montana in 1880. Bruner, L., 2267.

Natural history of the, 1578.

and other insects in the Northwest during the summer of 1883. Bruner, L., observations on the, 2277.

Rocky Mountain locust. Philosophy of the movements of the, 1669.

scourge, 1629.

in Wyoming, Montana, etc., in 1881. Bruner, L., the, 2267.

Rocky Mountains. Excursion to, 988.

Rogas n. sp., 879.

sp., 1002.

Roller. Strawberry-leaf, 984, 1059, 1574.

Roman-nosed pupa, 1217.

Rooms. Swarms of minute flies in, 596.

Root Aphis, 990, 1038.

Apple-borer and, 1038,

blight. Apple, 997.

borer, 981.

Raspberry, 1363.

lice. Knots on apple-tree roots caused by, 1187.

louse. Barley, 2394.

Syrphus fly, 1059.

Roots of Ampelopsis. Swellings on, 1428. Insects affecting apple-tree, 963.

Rosaceæ. Diastrophus confined to, 1131.

insects, 1149.

Rose. Bedeguar of the, 1939.

beetle, 361.

in California. Fuller's, 1740. Fuller's, 1708, 1721, 2291. Habits of Fuller's, 1708. Not Fuller's, 1825.

bug, 373, 1075.

on apples, 748.

Rose bug remedy, 1478.

bugs, 1278.

bushes. Bark-lice on, 1303.

chafer, 1329, 1583.

chafers on grape-vines, 1375.

gall, 1235.

Mossy, 570, 1166.

Prickly, 1194, 1245.

galls. Oak and, 1037.

Rose. Insects affecting. Aramigus fulleri, 1708, 1721, 1740.

Cynips bicolor = Rhodites bicolor.

Diaspis rosæ, 1303.

Heliothis marginidens = Pyrrhia exprimens.

Lecanium oleæ, 1303.

rosæ, 1303.

Macrodactylus subspinosus, 361, 373, 1075, 1329, 1375, 1478, 1583.

Monostegia rosæ, 672, 1780.

Pyrrhia exprimens, 677.

Rhodites bicolor, 1037.

rosæ, 570, 1037, 1166, 1194, 1245, 1939.

sp., 1235.

Selandria rosæ = Monostegia rosæ.

Tetranychus telarius, 1710.

Rose-slug, 672, 1780.

twig. Puncture on, 1167.

worms, 677.

Roses. Failuro of tea, 1708.

The red spider on, 1710.

Rot in plums. Preventing, 1685. Rotten root, 758.

Round-headed apple-tree borer, 1059, 1608, 2238.

New facts about the, 1630.

Oviposition of the, 2266.

Rove beetle. Spotted, 1123. beetles, 1558.

Royal horned caterpiller, 702, 803, 860, 1275.

Rubus insects. 1503,

Rudbeckia. Aphid on, 2205.

RUDBECKIA. INSECTS AFFECTING.

Nectarophora rudbeckiæ, 2205.

Siphonophohra rudbeckiæ = Nectarophora rudbeckiæ.

Rue. Papilio asterias on, 1512.

worms, 1512.

Rumex insects, 1165.

Russia. Excessive injury done by a beetle in, 1935.

Rust. Grain Aphis vs., 1806.

and Hessian fig. Wheat, 1605.

of orange, 2308.

Hubbard, H. G. Report on,

red social wasp, 1329, 1558.

Rustic. Corn, 1059.

Ruta graveolens. Insects on, 1512.

Rye. Beetle working in wheat, oats, and, 1259. gall-gnat, Loew, H. Description of, 2267.

INSECTS AFFECTING. RYE.

Cccidomyia secalina, 2267.

Silvanus surinamensis, 1259.

Tenebrio molitor, 72.

Ryc. Worms in flour and, 72.

Sack bearers, 1052.

Saddle-back caterpillars, 60, 161, 424, 829, 1092.

Sago. Galls growing on wild, 1347.

St. Louis Academy of Science. Presidential address, 1564, 1629.

Sale of silk-worm oggs, 1908.

Salivary glands of Phylloxera, 1687.

Salix. Galls of Cecidomyida on, 46, 197.

nigra. Vanessa antiopa on, 1609.

Salt and vinegar for insects, 937.

water insects used as food, 2203,

Salutatory, 386.

Samia cecropia = Attacus cecropia.

columbia = Attacus columbia.

cynthia = Attacus cynthia.

The ailanthus silk-worm, 1718.

Food-plants of, 2204.

ricini = Attacus ricini.

San Joaquin Valley, California. Coquillett, D. W. Report on locusts of, 2363.

Sandy soil. The Phylloxera in, 2250.

Sannina exitiosa, 55, 587, 617, 871, 1017, 1059, 1070, 1475, 1513, 2238.

Saperda, 379, 934, 2267.

bivittata = S. candida.

Oviposition of, 1621.

calcarata, 71, 963, 1849.

candida, 2, 6, 55, 57, 277, 377, 421, 578, 714, 870, 939, 962, 1023, 1038, 1056, 1059, 1213, 1227, 1332, 1367, 1441, 1513, 1608, 1621, 1630, 2230, 2238, 2266.

lateralis, 583.

Sapenda. Variation in the two-striped, 1213. on willow, 2267.

Sarcophaga, 1390.

carnaria, 1390, 1452, 2343.

lineata destructive to locusts in Dardanelles, 2075.

sarraceniae n. sp., 1385, 1390, 2343.

Sarcophagidæ, 2256.

Sarcoptes scabiei, 497.

Sarcopsylla penetrans, 412, 497.

Sarracenia variolaris. Description and natural history of two insects which brave the dan-

gers of, 1390.

Insect-catching habits of, 1385, 1390.

Sarracenias, 1499.

Satellite Sphinx, 1127.

Saturnia io - Hyperchiria io.

maia = Hemileuca maia.

Sance. Maggots in, 1607.

Savin twigs. Eggs of periodical Cicada in, 698.

Saw-flies, 294.

Spruce-tree, 115.

Saw-fly. Ash, 2291.

eggs, 108.

Elm-tree, 699.

Gooseberry, 140.

Imported, 228, 333.

Grape, 2291.

Larch, 2232.

Large, 1514.

larvæ on the quiuce, 2243.

Plum, 2291.

Raspberry, 1212, 1641.

Strawberry, 2418.

Scab in apple vs. apple-tree plant-lice, 1139. potato, 239.

Scabby potatocs, 1444.

Scale on acacias. Large white, 1730.

Cottony maple, 1816, 2291.

ou Euonymus latifolia, 2403.

insect on maple, 1890.

Cottony, 1515.

Pine-leaf, 1329.

on raspberry, 1968.

Scale insects affecting the orange. Experiments on, 2164.

> Coquillett, D. W., gas treatment for, 2418.

Experiments on, 2232.

Introduction and spread of, 2232.

Koebele, A., experiments against, 2418.

on magnolia, 1377.

Methods of destroying, 2119.

New species of, 1919.

of the orange, 2119, 2369.

in Florida. Voyle, J., report on the effects of cold, 2277.

The use of gases against, 2389. White-pine, 97.

Scales, 218.

Apple-tree, 69.

Scarites subterraneus, 805.

442Scarlet mite, 1470. Scarred apple trees, 951. Scavenger? Is Cyrtoneura a parasite or a, 2102. mistaken for a foe, 416. not a parasite. Phora a, 1923. Scelio ovivora, 1643. Scenopinus from human lungs. Larva of, 1348. sp., 2354. Schadlicherer Inseckten. Einige unserer, 1325. Schizoneura americana n. sp., 1678. cornicola, 27. fungicola, 27. lanigera, 58, 367, 372, 373, 467, 495, 963, 990, 997, 1038, 1059, 1187. rileyi, 1059. tessellata, 2361. ulmi, 1059. Sciapteron polistiformis, 373, 1301, 1509. robiniæ, 2410. Sciara, 239, 596, 1547, 1662, 1950. mali, 211. ocellaris, 2119. Scientific names, 101, 430, 826. nomenclature, 303, 768. symbols, 405. Scolopendra castanipes, 834. Scolytus caryæ n. sp., 220, 938 = S. 4-spinosus. destructor, 938, 1329. fagi n. sp., 220. Food-habits of, 1940. Hickory, 1754. Imported orchard, 2233. pyri = Xyleborus pyri. quadrispinosus, 220, 938, 1329, 1401, 1754. rugulosus, 1940, 2233. Synoptic table of, 220. Scorpion in Kansas, 1119. Scorpions, 453. Scotch and Austriau pines. False caterpillars on the, 1011. Scrape our trees? Shall we, 1399. Screw-worm, 209. in Central America. Prevalence of the, 2158. its parentage in doubt, 1880. Pyrethrum for the, 1921. Scudderia curvicauda, 1363, 2241. Scurfy apple-bark louse, 2305. Seymnus cervicalis, 1059. Læmorrhous, 151. sp., 151. Scyphophorus yuccæ, 1602. Second report State entomologist of Missouri, 1127.

U. S. Entomological Commission, 1957. Secretion on stems of bitter-sweet. White, waxy, Secd corn maggot, 657, 1059, 1065. grain. Chinch-bug not in, 888. peas from bugs. To keep, 434. ticks under bark of apple-trees, 1133. weevil. Honey locust, 1026. Seeds and galls. Jumping, 1496, 2163, 2173. Segments in head of winged insects. Packard, A.

S. Number of, 2267.

Selandria cerasi = Eriocampa cerasi. rosæ = Monostegia rosæ. rubi = Monophadnus rubi. vitis = Blennocampa pygmæa. Self-taught entomologists, 276. Scmasia helianthana n. sp., 1968. principora n. sp., 373 = Grapholitha prunivora. Semiotellus chalcidiphagus n. sp., 384, 563. destructor, 1581. Send plenty of specimens, 592. Senometopia atropivora, 2343. militaris n. sp., 6. Serica iricolor, 74. vespertina, 1140. Sericaria mori, 45, 542, 1311, 1346, 1609, 1616, 1648, 1718, 1721, 1724, 1753, 1945, 2062, 2234, 2381. Length and weight of thread of, 1359, Sericornis claypoleana n. sp., 2114 = Steganoptycha claypo'eana. instrutana = Steganoptycha claypoleana. Serrell automatic silk-reel. Walker, P., 2388. Sesia pelasgus = Hemaris thysbc. Seventeen-year Cicada, 1034, 1093, 1698, 2312, 2314. in Iowa, 1737. Periodical or, 2312, 2314. Sting of the, 407. locust, 370, 884, 1489, 1971, 1979. or thirteen-year locust. Periodical Cicada, alias the, 1159. Seventh report State entomologist of Missouri, 1423. Severe cold on insects. Effects of, 2037. Sex in butterflies. Controlling, 1352. insects. Regulating, 1415. Shad. Terrestrial insects in stomach of, 1853. Shade trees and their insect defoliators, 2378, 2379. Shaffer. Entomology. Prof. Riley to Dr., 2360. Shagreened cut-worm, 2291. Shall we scrape our trees, 1399. Shedding of tracheæ and double cocon 1775. Sheep bot, 450. fly, 2238. gad-fly, 887. Sheldon's borer remedy, 195. Shell-bark hickory. Citheronia regalis on, 775. Shimer, H., criticised by B. D. Walsh, 174. Shrubs, truit, and grains. Webster, F. M. Ex. periments on the effect of punture of Hemiptera on, 2386. Sialidæ, 2267. Sialis infumata, 39. Sigalphus Curculio parasite, 1301. curculionis, 1751, 1795. rufus n. sp., 1301 = var. of curculionis.Silesia in 1869. Cohn, F. Hessian-fly in, 2267. Silk-culture, 1668, 1753, 2119, 2291, 2363, 2394. in California. Promotion of, 2000. the colonies, 2278.

how to dispose of cocoons, 1881.

in Kansas, 1542.

Profits of, 2338.

Silk-culture in the United States, 1668, 1945, 2070, 2196, 2336. Walker, P., 2418. fiber from coeoons raised at the Department. McMurtrie, W. Test of, 2253. industry in the United States (Review), 2268. producer. Antheræa yama maia as a, 1346. reel. Walker P. Serrell's automatie, 2386. spiders, 830. Silk-worm, 2095. Ailanthus, 899, 1311. breeding, 1648. Ceeropia, 1311. Circular, 2303. cocoons, 1486. eggs, 1114, 1753. Japanese mode of packing, 1616. prices and whore obtained, 2062. Sale of, 1908, 2303. Length and weight of the thread of the, 1359. Luna, 1311. Manual of the, 1721, 1724. in Missouri. Ailanthus, 1460. Mulberry, 1311. naturalized. Ailanthus, 1179. notes, 2232. Osage orange for the mulberry, 1220, 1286. vs. mulberry for the, Perny, 1311. Silk-worm. Polyphemus, 1311. Promethea, 1311. Tusseh, 1311. Yamamai, 1311. Silk-worms fed with osage orange, 1341. Food for, 1318. Nomenelature of American, 303. Silky mite, 1423, 1625. Silpha americana, 817, 1226. peltata = S. americana. snrinamensis, 1259, 1260, 1261. Silphurus femoratus, 39. Silvanus. Grain, 1259. in dried English currants, 1260. in flouring-mills, 1261. Silver plate by insects. Damage to, 2154. Simulium, 1160, 2291, 2416. attacking mules, 1811. feeding on other insects, 2177. from Lake Superior, 2032. meridionale n. sp., 2394. molestum, 1174. peeuarum n. sp., 2394. pietipes, 1958, 2032. piseidium n. sp., 1174, 1283, 1345. Sinea diadema, 34, 638, 731. Singular caterpillar, 880.

Sinoxylon basilare, 1311, 1747.

Red shouldered, 1311.

Siphonophora, 1678.

avenæ = Nectarophora granaria. rudbeckiæ = Nectarophora rud-

Sitaris, 1600, 1643.

Sitones flavescens, 2394.

Sitophilus granarius = Calandra granaria.

oryzæ = Calandra oryzæ.

remotepunetata = Calandra remotepunctata.

Six worst insect enemies of fruit-growers in northern Illinois, 377.

Sixth report State entomologist of Missouri, 1363.

Skeletonizer. Apple-leaf, 1311, 1322.

Skipper. Choeso, 1700.

Skippers injuring smoked ham, 1734.

Slug on pear and cherry-trees, 1222.

Pear-tree, 1382.

Rose, 672, 1780.

worm. Spined, 1150.

Small apple-leaf worm, 747.

borer in apple-twig, 1458.

galls and minings in apple-twigs, 552.

A new leaf-hopper injurious to, grains. 1767.

> and grasses. Webster, F. M. Inseets affecting, 2394.

reddish snout beetle in apple, 1244. white bristly cut-worm, 1059.

Smaller cornstalk borer, 2119.

Smart bugs. Concerning certain, 565.

Smeared dagger, 1301.

Smerinthus. Blind-oyed, 1912.

exeæeatus, 1912.

Smiera albifrons, 6.

Smilax. Agrotis saucia injuring, 1941.

injured by cut-worms, 1941.

Smilia aurieulata, 1183.

Smith, E. A. The cotton belt, 2343.
Smith, J. B. Report upon eranberry and hop in sects, 2277.

> Report upon insects affecting the hop and the evanberry, 2291.

Smith's patent Cureulio-trap, 969.

Smoked hams. Skippors injuring, 1734.

Smut in wheat, 1479.

Cause of, 1461.

Snake of Brazil. Lignified, 2136.

worms, 1547.

Suakes. Hair, 612, 861, 1143.

Horse hair, 612.

Snellen von Vollenhoven. Obituary, 1817.

Snout beetle, 700, 1168, 1302.

on apple. Small reddish, 1244.

Imbricated, 1301, 2291.

Large gray straight horned, 1033.

beetles injurious to fruits, 1302.

Aphididæ on, 1184. Snow-balls.

Lieo on, 1184.

Snowy tree-erieket, 1059, 1329, 2238, 2291.

in raspberry canes. Eggs of, 1122.

So-called army-worms. Three, 328.

web-worm of young trout, 1283.

Seap. Apply, 1367.

against borers, 47.

Social wasp. Rust-red, 1329, 1559.

Social wasps, 616.

Sod-worm, 2418.

Soft maples. Flat-headed borer in, 1250.

Species. Theory of, 384. Soil. The Phylloxera in sandy, 2250. Specific for the Colorado potato heetle, 1490. Soils. Effects of Paris green on, 1427 names. Capitalizing, 2170, 2257. Solanaccæ insects, 1238. value of Apatura aliciæ, 1977. Solanum carolinonse. Cassida texana on, 2096. Specimens lost, 1210. Doryphora juncta on, 2096. elæagnifolium. Cassida texana on, 2096. Send plenty of, 593. Flea-beetle eating, 1782. Speckled eut-worm, 1059, 2291. insects, 1238. Specters. Walking-sticks or, 1395. Speetrum bivittatum, 1298. melongena. Cassida texana on, 2215, 2235. femoratum = Diapheromera femorata. Doryphora juncta on, 2096, Speculations of the New England school of nat-2215, 2235. Solar physics and locust multiplication and miuralists. Entomological, 44. gration. Swinton, A. H., 2267. Professor Dana and his ontomolog-Soldier-beetle larvæ, 1643. ical, 82. Pennsylvania, 1059. Spermophagus robiniæ, 1026, 1474. bug, 1558, 2291. Sphæria morhosa, 930. Bordered, 1059. Sphærophthalma occidentalis, 814, 832, 858. Glassy-winged, 1301. Sphecidæ. Table of genora of, 375. Rapacious, 1059. Sphecius speciosus, 371, 372, 407, 543, 858, 2014. Sphenophori that attack corn. Larval habits of, Ring-banded, 1311. 2030. Spined, 456, 1059, 1311, 1329, 1523. Sphenophorus. Cocklebur, 1301. Solenohia, 38, 943. Corn, 1301. Solenopsis geminata, 2105. Grain, 2363, 2394. xylonii = S. geminata. parvulns, 2394. Solidago. Aphid on, 2205. robustus, 2030, 2119. gall moth, 1059. Grapholitha olivaceana hred from, 2285. sculptilis, 337, 1042, 1301. leaves. Galls on, 1924. zeæ n. sp., 337 = S. sculptilis. nemoralis. Cecidomyia carbonifera on, Sphex ichueumonca, 543. Sphida obliquata, 2357. 1924. Song notes of the periodical Cicada, 2334. Sphingicampa n.g., 40. Sorghum. Nola sorghiella on, 2119. bicolor, 40, 45. Sphingicampa distigma n. sp., 40 = S. hicolor.Sorghum weh-worm, 2119. Sound organs in sphingid pupæ. Probable, 2101. Sphingid pupa. Prohable sound organ in, 2101. South America. Report on cotton crop and its Sphingidæ, 771, 1089, 1277, 1418, 1419, 1761, 1784, 2101. enemies in, 2343. Sphinx. Abhot, 1127, 1248, 1277. American lepidoptera. Notes on, 1784. Achemon, 1127. Entomological ignorance in the, 390. atropos, 2101. carolina = Protoparco carolina. Field for the entomologist in the, 1109. Fruit-culture in the, 2265. catalpæ = Coratomia catalpæ. The imported cabbage-worm in the, 1714. hageni = Ceratomia hageni. Musclo-shape bark-louse on apple trees in Larva of Abbot, 1248, 1277. the, 1774. Aquatic, 1951. Notes from the, 1657. Osage orange, 2119. Pass, Ill. The bag-worm at, 1156. 5-maculata = Protoparce celeus. side of trees. Apple-tree borers on, 634. Satellite, 1127. west. Economic investigations in the, 1864. White-lined morning, 630, 1198, 1301, 1643, Southern buffalo-gnat, 2291, 2394, 2418. cabhage-butterfly, 1127, 2232, 2238. Sphyracephala brevicornis, 299. grass-worm, 1127. Spider egg-nest. Mud-wasp and, 1847. Illinois. Entomological tour in, 372. Egg sacks of some unknown, 1144. Sowing cotton secds in hot-heds, 1772. Ladder, 1299. Spain. Phylloxera congress in, 1906. and nest, 1869. Span-worms. Gooseberry, 1068, 1570. Poisonons, 2306. Sparrow. Anent the English, 1667. Red, 1146, 1225, 1710. as an insect killer. Worthlessness of on roses. Red, 1710. the, 2413. wort owlet-moth, 1301. Spathius trifasciatus n. sp., 1329. Spiders, 217. Spattered-copper underwing, 1301. Grape leaf-folders eaten by, 468. Spearman. Larva of the ten-striped, 866. Jumping, 2302. Ten-striped, 32. Silk, 830. Spocies. Goographical range of, 1614. Trees injured by, 2409. 'Old question of, 2201. Spilochalcis mariæ, 1112. of Otiorhynchidæ injurious to cultiva-Spilonota oculana = Tmetocera ocellana. ted plants, 2117. Spilosoma acræa, 2343.

virginica, 296, 454, 1202, 1307.

Phytophagic, 45, 50.

445

for Minnesota, 1108.

INDEX.

Spindle-worms, 331. State entomologist of Illinois, 327. Spined slug-worm, 1150. soldier-hug, 456, 1059, 1311, 1329, 1523. spider, 813. Spirobolus marginatus, 166, 963, 1025. Spittle insect, 93. Spotted lady-hird, 599. Peliduota, 1301. rove-beetle, 1123. touch-mc-not. Cecidomyia impatientis on, 852. Galls on, 852. rumpet-leaf. Insects associated with, 1385, 1390. Spraying trees for protection against insects. Im proved method of, 2211. Spread of Pieris rapæ into Alahama, 1720. scale insects. Introduction aud, 2232. Spring canker-worm, 1423, 1482, 2238. Howell, M. A. Experience with, 2267. Locust injury next, 1555. tails, 317. Sprinklers and atomizors, 1857. Spruce-borer. Flat-headed, 2267. hud. Tortrix, 2232. worm. Reddish yellow, 2291. Epizeuxis, 2363. SPRUCE. INSECTS AFFECTING. Epizeuxis, 2363. Lophyrus ahietis, 115. Mclanophila, 2267. Nematus integer, 2232. Plume moth, 2363. Therina, 2363. Tortrix fumiferana, 2332. Spruce. Nematus, 2232. plume-moth, 2363. Therina, 2363. tree saw-flies, 115. Squares. Butterfly Iarvæ injurious to cotton, 1872. Squash-borer, 378, 1127. bug, 409, 867, 1059. does not touch the white bush scollop, 825. Glorified, 262. SQUASH. INSECTS AFFECTING. Anasa tristis, 409, 825. Corcus tristis = Anasa tristis. Diabrotica vittata, 355. Epilachna borealis, 125. Melittia ceto, 125, 248, 378. Trochilinm cucurbita = Melittia ceto. Squash-vine borer, 248. insects, 125, 355. Squirrel-bot, 526. Stag-beetle, 957, 1517. Horns of the, 755. Stalk-borer, 1558, 1595, 1646, 2291. Dahlia, 862, 1009. aud aster, 940. Rice, 2119. Tomato, 694, 976. weevil. Potato, 1558. Staphylinidæ wanted. Americau, 1786.

Staphylinus maeulosus, 1123, 1186.

New York, 1863. One day's journal of a, 383. Status and future prospects of silk-culture in the United States, 2336. Steele, J. P., report on cotton insects, 2343. Steganoptycha claypolcana n. sp., 2206. Stick-bug, 448. Stictonotus isosomatis n. sp., 2119.Still they come, 908. Sting? Do locusts, 371. of the seventeen-year, 407 Stinging hug, 778. caterpillars, 1748. larvæ, 760, 811, 1329. Stings of bees, 116. insects, 116, 217. Stiretrus anchorago, 372. fimhriatus = S. anchorago. Stizus hrevipennis n. sp., 375 = Megastizus brevipennis. grandis = Sphecius speciosus. speciosus = Sphecius speciosus. Stock. Insect injurious to live, 2238. Stomach of hlack-hass. Larvæ in, 1792. blue-bird. Ichneumon in, 1878. Larvæ from, 1871. eat-bird. Tipula eggs in, 1735, 1745. lark, rohin, and sun-fish. Insect from, meadow-lark. Boetles in, 1013. robin. Cut-worms from, 1873. rock-hass. Iusects from, 1793. shad. Terrestrial insects in, 1853. Stomoxys calcitrans, 864. Stored corn. Serious injury to, 1683. rice. Insects affecting, 2190. Strachia histrionica = Murgantia histrionica. Straight-horned snout-beetle. Large gray, 1033 Strange bug, 1078. Strawberry heds. White-grub in, 1236. borers, 2385. hugs, 637. crown-horer, 1301. Notes on, 1393. destroyer, 689. Eggs of bugs on, 690. enemy, 342. and grape-vines. Injured, 682. STRAWDERRY. INSECTS AFFECTING. Ægeria impropria, 2385. Agrotis tricosa, 2324. Anarsia liueatella, 2235. Anchylopera fragariæ - Phoxopteris fraga-Anthonomus musculus, 2363 Capsus oblineatus = Lygus pratensis. Chrysomelidæ, 2235. Colaspis flavida, 1904. CorimeIæna pulicaria, 637. Cyclocephala immaculata, 1236. Eccopsis permundana, 2324. Emphytus maculatus = Harpiphorus macula-Graphops, 2229.

STRAWBERRY. INSECTS AFFECTING-Continued. Sunfish. Insects from stomach of, 1926. Sunspots and insect life, 2094. Harpiphorus maculatus, 499, 955, 964, 1570, 1586, Supposed army-worm in New York and the east-2324. Julus sp., 834. ern States, 1990. bark-lice eggs in Missouri, 1084. Lygus pratensis, 682, 2235. cause of yellows in peach trees, 515. Monostegia rosæ, 984. dock. Galls on, 1165. Myriapod, 2235. Paria aterrima, 1904, 2229. eggs of the preying Mantis, 1002. hibernating Aletia chrysalids, 1927. Phoxopteris fragariæ, 499, 984, 1059, 1574, 2324. trout enemy, 1141. Polydesmus serratus, 342. Swallow-tail. Larva of thoas, 1237. sp., 834. Saw-fly, 2418. Philenor, 1127. Thoas, 1268. Selandria rosæ = Monostegia rosæ. Tyloderma fragariæ, 1301, 1393. Swallows, 878, 1502. Strawberry leaf-roller, 984, 1059, 1574. Dipterous larva on, 153. Swarms of butterflies, 406, 1050. A new enemy to, 1904. Pests of the, 2324. lady-birds, 824. saw-fly, 2418. minute flies in rooms, 596. weevil, 2363. that devastate the trans-Mississippi worm, 955, 965, 1570. country. Locust, 1674. and remedy, 1586. Sweet-potato beetles, 681, 1510, 1697. worm, 499. SWEET-POTATO. INSECTS AFFECTING. Streaked cottonwood leaf-beetle, 2291. Cassida bivittata. 681, 1127, 1510. Striped-beetle, 2291. nigripes, 1127, 1510, 1697. blister-beetle, 1059, 1230, 1558. sp., 94. Cassididæ, 993, 1056, 1082, 1127, 1510. bug, 123, 175. Coptocycla aurichalcea, 681, 1127, 1510. cucumber-beetle, 773, 1127, 2238. bug, 148. bivittata = Cassida bivittata. flea-beetle, 2291. guttata, 1127, 1510. Sturnella magna. Contents of the stomach of, 1013. nigripes = Cassida nigripes. Subangular ground beetle, 1059. Gryllus sp., 2384. Subscribers. To our, 1106. Swellings in apple scions, 548. Subterranean mites. Description of new, 1370. roots of Ampelopsis, 1428. Successful management of the most destructive Swinton, A. H., Solar physics and locust multipliorange insects, 2088. cation and migration, 2267. Successor. Mr. Walsh's, 1105. Switzerland. Extermination and restriction of Sucking organs of bees, wasps, and flies, 2182. Phylloxera in, 2207. Sugar-berry. Leaf-galls and caterpillars, 762. SYCAMORE. INSECTS AFFECTING. Orgyia leucostigma on, 762. Cerambycid, 2267. Lachnus platanicola, 2138. cane in Brazil; Branner, J. C. Insects in-Symbols. Scientific, 405. jurious to, 2277. SUGAR CANE. INSECT AFFECTING. Synchlora rubivoraria, 721, 1059. Ligyrus rugiceps, 1794. Synergus albipes = S. lana. Sugar-maple. Beetle in, 1014. lana, 41. borer, 2291. lignicola, 41. Eggs on, 350. mendax n. sp., 41. rhoditiformis n. sp. 41 = S. lignicola. Mite gall on, 1265. tree-borer, 547. Synonyms of parasites, 1932. SUGAR-TREE. INSECTS AFFECTING. Synophrus albipes n. sp. = Synergus lana. Arhopalus speciosus = Plagionotus speciosus. Synopsis of North American Heliothinæ (Review), Plagionotus speciosus, 547. 2178. 25. Sugaring for moths, 1300. Syntomeida sp., 2412. Natural, 2138. Syringa. Citheronia regalis feeding on, 803. Sulphur cure on peach trees, 176. Syrphidæ. Larval habits of, 168. in trees, 98, 228, 931, 1154, 1383. Syrphus-fly. Root-louse, 1059. Sumach-beetle. Jumping, 1363. mellinus, 1813. Eggs in, 118. sp., 1251, 1798. SUMACH. INSECTS AFFECTING. caught by flower of Bidens chrysan-Blepharida rhois, 1363. themoides, 1761. Coccid, 320. Systematic position of the Orthoptera. Packard, Orcholimum sp., 118. A. S., 2267. Xiphidium sp., 118. relations of Platypsyllus as deter-Summer. Anticipated locust injury next, 1615. mined by the larva, 2417. dormancy of butterfly larvæ, 1410. Systechus leucophæus, 2355. larva of Phyciodes nyc-

teis, 1410,

oreas, 1541, 1947, 1959, 2002,

Tabanus atratus, 1127.

Tabanus, sp., 40. Test of machinery for destroying the cotton-Table. On onr, 395, 408, 506, 546, 568, 623, 827, worm. Barnard, W. S., 2253. Tests of silk-fiber from cocoons raised at the De-Tachina aletiæ n. sp., 1712, 2343. partment. McMurtrie, W., 2253. anonyma n. sp., 1311. Tetranychus americanus, 1326. archippivora n.sp., 1301 = Masicera arirritans, 1326. chippivora. telarius, 1146, 1225, 1710. concinnata, 2343. Means against, 508. flies, 1643, 1736. Tetraopes femoratus, 323. fly, 1558. 5-maculatus, 1140. Anonymous, 1423, 1625. tetraophthalmus, 323. tornator = T. tetraophthalmus. Cecropia, 1311. Red-tailed, 1127. Tetrastichus esurus, 1712, 2343. Yellow-tailed, 1127, 1625. productus n. sp., 2332.fraterna, 2343. Tettigonia coagnlata, 1024. phycitæ, 1311. sp., 951. sp., 2378. vitis = Typhlocyba vitis. villica, 2343. Tettigonidæ, 452. Tachinid eggs, 1914. Tettix granulata, 1566. Texas again. Tarantula of, 493. Tachinidæ, 66, 1301. fever. Ticks and, 404. Tæniopteryx fasciata, 249. in 1883. Anderson, E. H. Cotton-worm in Tamarack. Pieris menapia feeding on, 2121. Tamariscus. Nanodes tamarisci on, 1496. south, 2253. Tansy for borers, 1016. spring of 1886. Bruner, L. Locusts in, Tarantula killer. Notes on the, 823. 2382. of Texas, 466, 521, 1178. Tarantula of, 466, 521, 1178. Thalessa, 2286. again, 493. Tarnished plant-bug, 1127, 1219, 2291, 2363. atrata, 338, 477. Tarred paper for fruit trees, 1568. lunator, 1111, 1126, 2350. Tat. Tit for, 432. Thecla peas, 1872. Tawney emperor, 1363. Thelaxes ulmicola = Colopha ulmicola. Tea roses. Failure of, 1708. Thelia bimaculata, 787. Telea polyphemus, 244, 542, 629, 635, 841, 1164, 1311. Theory. Plug-ngly, 233. wanted. New locust, 1532. Caterpillar of, 765. Theridula sphærnla, 2343. Cocoons of, 2181. Therina. Pine, 2363. Telenomus, 2115. bifidus, 2378, 2394. Spruce, 2363. Telephoridæ, 374. Thersilochus conotracheli n. sp., 1301, 1751. Telephorus bilineatus, 1311, 1885. Thick-thighed walking-stick, 1701, 1721. Temnochila virescens = Trogosita virescens. Third report State entomologist of Missouri, 1301. Ten-lined potato beetle, 925. U. S. Entomological Commission, . striped spearman, 32. 2267. Larvæ of, 866. Thirteen-year locust. The periodical Cicada alias Tenacity of life, 1755. the seventeen-year and the, 1159. Tenebrio molitor, 72, 191, 980, 2167. Thistle. Beetles on, 169. obscurus, 191, 2165. THISTLE. INSECTS AFFECTING. Tenebrionidæ, 2105. Diabrotica longicornis, 169. Phyllobrotica longicornis = Diabrotica longi-Tent caterpillar, 642. of the apple tree, 336, 1301, 2238, cornis. Platyptilus carduidactylus, 1059. Eggs of the American, 1329. Thistle-plume, 1059. of the forest, 645, 688, 1181, 1200 Thoas swallow-tail, 1268. Larva of the, 1237. 1301, 1331. THORN. INSECTS AFFECTING. caterpillars, 363. and fall web-worms, 819. Anthonomus cratægi, 376. Tenthredinidæ, 197, 294. Cecidomyid, 376. Descriptions of new, 385. Thorn-leaf gall. Mr. Couper's, 376. Larvæ of, 108. Those centennial insects, 1511. Tenthredo. Venation of, 197. Thousand-legged worm, 193, 236, 261, 834, 1025. Teras cinderella, 1311. Three-banded Spathius, 1329. malivorana, 1311. lined leaf-beetle, 1059, 1558. Willow, 2363. so-called army-worms, 328. Terias, 1711. worms and their work, 1504. Thrips, 1127, 2362. Termes, 2267. Food-habits of, 53, 280, 685, flavipes, 1620, 1729.

sp., 203,

Terrestrial larvæ in stomach of shad, 1853,

vs. bugs, 1103.

Thrips. True and bogus, 203. TOBACCO. INSECTS AFFECTING. of the vine-grower, 53. Crepidodera cucumeris, 1782. Thrushes. Food-habits of, 1726. parvula, 1782. Thyreus abbotii, 763, 1018, 1118, 1127, 1248, 1277. Epitrix cucumeris = Crepidodera cucumeris. Thyridopteryx ephemer:eformis, 182, 206, 266, 271, hirtipennis = Crepidodera parvula. 427, 538, 629, 641, 738, 815, 1036, 1059, 1090, 1156, 1189, Gryllus sp., 2384. 1352, 1424, 1472, 1650, 1658, 2272, 2378, 2379. Macrosila carolina — Protoparce carolina. Thyridopteryx ephemeræformis. Remarks on, 5-maculata = Protoparee celens. Protoparce carolina, 131, 933, 1262, 2238. 2272. Thysania zenobia, 1291. celeus, 131, 771, 885, 933, 1329. Thysanoptera, 174. Sphinx carolina = Protoparce carolina. Thysanura, 2267. 5-maculata = Protoparce celeus. Tibicen cassinii, 1057, 2367. Tobacco moth, 131. septendecim, 51, 189, 213, 370, 407, 474, 527, plants. Flea beetles on young, 1782. 618, 619, 648, 698, 707, 746, 884, 920, 1034, worm, 885, 2238. 1037, 1057, 1059, 1093, 1159, 1311, 1489, 1609, moth, 1059. 1624, 1698, 1737, 1809, 1836, 1971, 1979, 1994, Counterworking the, 1329. 2014, 2112, 2144, 2216, 2305, 2307, 2312, 2314, Tilden tomato and the, 933. 2315, 2318, 2320, 2321, 2329, 2334, 2363, 2367. Tolype velleda, 972, 1773. tredecim, 474, 746, 1034, 1057, 1159, 1609, 1624, Tomato feeding worm, 850. 1809, 1836, 1979, 1994, 1997, 2014, 2112, 2307, fruit worm, 1136. 2312, 2329, 2367. gall. Grape-vine, 1329. Ticks under bark of apple-trees. Seed, 1133. TOMATO. INSECTS AFFECTING. and Texas fever, 404. Doryphora 10-lineata, 21. Tiger beetle. Common, 1763. Gortyna nitela, 694, 734, 976. larva, 719. Heliothis armigera, 1136. Virginia, 1059. Prodenia commelinæ, 850. Protoparce carolina, 933, 1089. beetles, 628, 1643. moth. Chrysalis of the virgin, 674. celeus, 152, 155, 933, 1089, 1661. Isabella, 1311. Sphinx 5-maculata = Protoparce celeus. Tilden tomato and tobacco-worm, 933. Tomato stalk borer, 694, 976. Tile-horned Prionus, 1127. and the tobacco worm. Tilden, 933. Tilia gall, 1116. worm, 152, 1059, 1558, 1661. Odontota rubra on, 1849. again, 1104. Timber borers, 191. Worm eating into green, 734. worm. Parasitized, 155. encourages apple-tree borers, 1441. That venomous, 1089. Insect in, 918. TIMOTHY. INSECTS AFFECTING. worms not poisonous, 771. Hadena devastatrix, 2394. Tomicus pusillus = Pityophthorus minutissimus. Sphenophorus parvulus, 2394. ramulorum = Pityophthorus microgra-Timothy: Webster, F. M. Insects affecting, 2394. phus. Tinea, 1633. Tongue. Butterfly's, 1815. Tineid. Apple-tree, 77, 92. Humming-bird moths caught by the, genus allied to Pronuba. A new, 1830. Tineidæ, 70. Moths and butterflies caught by the, New genus of, 1329, 1336. 1761. Tineids in bee-hives, 252. Too fond of honey, 764. Fossil, 2084. Toothed Dermestes, 2363. Tineina, 1246. Torrubia, 1064. Tingis amorphæ n. sp., 45 = Gargaphia amorphæ. elongata, 1430, 1436. ciliata = Corythuca ciliata. ravenelii, 1803, 1823. tiliæ n. sp., 45 = Gargaphia tiliæ. Tortoise beetle. Black-legged, 1127. Tiphia femorata, 1363. Clubbed, 1227, 1558. inornata, 1363, 2301. Golden, 1127. and Rhipiphorus, 2351. Larva of clubbed, 1238. Unadorned, 1363. Mottled, 1127. Tipula, 947, 1161, 1278, 1368. Pale thighed, 1127. eggs in stomach of cat-bird, 1735, 1745. beetles, 105, 767, 1082, 1127. sp., 512. Tortricid. A pretty new gall-making, 2176. trivittata, 512, 1735. Tortricidæ, 918, 1975. Tit for tat, 432. Description of new, 1969. Tmetocera ocellana, 747. Oviposition of, 1922. Toads, 217, 262. Tortrix cinderella n. sp., 1311 = Teras cinderella. fractivittana = Cacœcia fractivittana. eat worker bees? Do, 544. in gardens, 822. fumiferana, 2232.

rileyana = Cacecia rileyana.

Tortrix. Spruce bud, 2232. Walnut, 1059. Touch-me-not. Cecidomyia impatientis on spotted, 852. Galls on spotted, 852. Tracheæ and double cocoons. Shedding of, 1715. Trade in insects, 1986. Tragidion fulvipenne, 490. Tragocephala viridifasciata = Chortophaga viridifasciata. Transformations of insects, 528. Mite, 1618. of red mites, 1632. the tumble bug, 1386. Trap. Thomas Wiers's apple-worm, 1312. Trapping the carpet beetle, 1752. Treat's insect extinguisher (Review), 531. Tree borers of the family Cossidæ. Bailey, J. S., 2253. cricket, 251, 953, 961, 999. on grape vines. Eggs of, 723. Grapes cut off by, 414. Habits of, 207. Jumping, 1329. in raspberry canes. Eggs of, 1006. Snowy, 2238. Snowy, 1329, 2291. cut-worms, 229, 281. hopper. Buffalo, 415, 1329. injurious to potatoes. Buffalo. 2107. Trees. Apple-tree borers of the south side of, 634. injured by spiders, 2409. for protection against insects. Improved method of spraying, 2211. Shall we scrape our, 1399. Tremex columba, 458, 928, 1126, 2286. Pigeon, 458, 1126. Trenton, N. J. Bennett, T. Report of experiments at, 2344. Trichius delta, 372. Trichobaris trinotata, 401, 1059, 1558, 2172. Trichodectes ovis, 2043. Trichogramma minutum n. sp., 1301. pretiosa n. sp., 1712, 2115, 2343. Trichopsenius depressus, 1729. Trichoptera, 2267. Trifolium insects, 1459. Trimble's insect enemies of fruit, etc. (Review). Trimerotropis cœruleipes n. sp., 1959. latifasciata n. sp., 1959. similis n. sp., 1959.Triodites, 1947. mus, 1959, 2002. Trioza tripunctata, 669. Triphleps insidiosus, 1853, 2048. Triplax thoracica = Tritoma thoracica. Tritoma thoracica, 114. Tritoxa flexa, 225. Triungulin of Meloidæ, 2083. Trochilium acericolum = Ægeria acerni. acerni = Ægeria acerni. caudatum = Alcathee caudatum. cucurbitæ = Melittia ceto. denudatum = Fatua denudata.

29 ENT

Trochilium hospes, 197. sp., 1063. tipuliformis = Ægeria tipuliformis. Trogoderma tarsale, 2167. as a museum pest, 2139. Trogosita mauritanica = Tenebrioides mauritanica. virescens, 721. Trogus obsidianator, 1802. Troilus butterfly caterpillar, 469. Trombidium, 624. developed from Astoma, 2071. giganteum n. sp., 1632. holosericeum, 1470. locustarum n. sp., 146, 728, 1287, 1451, 1521, 1598, 1618, 1632. muscarum n. sp., 1632. sericeum, 1451, 1470, 1598, 1618. telarium = Tetranychus telarius. tiuctorium, 1470. Trout. Death-web of young, 1138, 1160, 1174. cnemy. Supposed, 1141. Food for, 1142. So-called web-worm of young, 1283. True army-worm, 647, 1127. and bogus Yucca moth, 1804. Thrips and bogus Thrips, 203. Trumpet grape-gall, 791, 1116, 1329. leaf. Insects associated with the spotted, 1385, 1390. vine. Clydonopteron tecomæ in seedpods of, 1929. Pyralid infesting seed pods of, 1929. Trupanea apivora = Promachus apivorus. Truths in applied entomology. General, 2291, Trypeta pomonella n. sp., 177, 367, 373, 454, 1320, 1553, 1654, 2119, 2238. solidaginis, 137, 180, 798, 1059. Tryphon atricoxus n. sp., 385. Trypoxylon albitarse, 543. Tulip-tree bark-louse, 271. Tumble-dung. Transformation of the common, Turf web-worm, 2418. Turk and its crescent. The little, 329. Turkey-gnat, 2394. Turnip enemy, 215. Polydesmus complanatus an enemy to, 215. Tusseh silk-worm, 1311. Tussock-moth. Caterpillars of the white-marked, 1227. Eggs of the white-marked, 480, White-marked, 1059, 1363, 2238, 2378. again, 535. Twelve-spotted Diabrotica, 687, 1127, 2418. Twice-stabbed lady-bird, 38, 1329, 1883. Twig-borers, 1052, 1185. girdler, 476, 1938. Twigs amputated by some unknown animal, 442. Berry and cherry, 1503.

Eggs in or on canes and, 1329.

Twigs girdled by some animal, 443. United States. Status and prospects of silkpruners, 288. culture in the, 2336. punctured by periodical Cicada, 1055. Unity of coloration in insects, 50. Two-striped locust, 1423, 2363. Universal remedies, 204, 503, 818. Saperda. Variation in, 1213. More, 278. swect-potato beetle, 1127. remedy. Another, 232. walking-stick, 1298. Unjust accusation, 1537. Tyloderma fragariæ, 1301, 1393. Unknown corn pest, 1073. . larvæ, 853. Typhlocyba aurea n. sp., 22. binotata n. sp., 22. moth, 753. pallidula n. sp., 22. worm, 986. tricineta, 25. Unmasked. A friend, 374. vitis, 203, 686, 1392, 1853. Unnatural secretion of wax, 782. Typhłocybini. New species of, 22. Unsightly galls on cotton-wood, 446. Typhlodromus pyri, 1739, 1759. Untrue. Attractive but, 1644. Tyroglyphus phylloxeræ n. sp., 1363, 1370. Urena Anomis, 2119. siro, 1703. lobata. Anomis erosa feeding on, 2343. Uji parasite, 1311. Uroceridæ injurious to apple-twigs, 893. Uloma impressa, 805. Urocerus flavicornis, 2347. Gall-making Pemphiginæ on, 1653. Uropoda americana n. sp., 1505, 1626.Ulmus. Unadorned Tiphia, 1363. vegetans, 1626. Unfledged locusts. Destruction of young or, Useful Labena, 1423. 1577. lives. Two, 2370. Habits of young or, 1578. A nuisance made, 983. Useless? Is any knowledge, 1135. Unicorn apple-tree caterpillars, 749. prominent, 1280. Ustilago segetum, 1461, 1479. Unimpregnated eggs hatch? Will, 1029. Utah in 1878. Packard, A.S. Notes of a journey Unique and beautiful noctuid, 2189. to, 1959. United States. Agricultural advancement in the, Utilization of ants in horticulture, 2089, 2137. 1750. Vagabond Crambus, 2119. Clothes moths observed in the, Valedictory, 326. Valery Mayet on Phylloxera, 2207. Valuable insecticides. Two, 1742. The cotton-worm in the, 1769. Vanessa antiopa, 540, 907, 1234, 1327, 1609, 1704, 1711. Department of Agriculture. Division of Entomologycomma = Grapta comma. interrogationis = Grapta interrogationis. Bulletin 1, 2164; 2, 2165; 3, 2253; 4, 2277; 6, 2304; Variable molting in Orgyia, 2379. 8, 2315; 10, 2378; 11, 2344; Variation in Anisopteryx, 1540. 12, 2364; 13, 2382; 14, 2388; the two-striped Saperda, 1213. Variegated cut-worm, 1059, 2291. 15, 2389. Entomological Commission. Eccopsis, 2363. Appropriation for, 1843. Varieties. Phytophagic, 45, 50. Bulletin 1, 1577; 2, 1578; Varying Anomala, 2291. Vegetable phenomena, 436. 3, 1736; 6, 2026. Circular 1, 1575; 2, 1576. Vegetal-feeding ground beetles, 1738. Report 1, 1643; 2, 1959; Vegetation caused by locusts. Changes in, 1495. 3, 2267; 4, 2343. Harris. Insects injurious, to (Re-Reports, 1757. view), 568. in Illinois. Insects injurious to, 6, 52. Supplementary instructions to agents of the, 1888. Legislation to control insects injuri-U.S. Entomologist. Report for 1878, 1721. ous to, 1946. Venation of wings of Anisopteryx, 1540. 1881 - '82, 2119. Venomous tomato-worm. That, 1089. for 1883, 2232. Vera Cruz, Mexico. Interesting cotton-worm 1884, 2291. notes from, 1845. 1885, 2363. 1886, 2394. Verified. Predictions, 1494. 1887, 2418. Vertical insect-boxes, 1963. United States. New source of wealth to, 1668. Vesicants. Meloidæ as, 912. Notes on Aphidinæ of the, 27, 1678. Vespa crabro, 616. Psyllidæ of the, 2210. maculata, 543. Vespidæ, 190, 505, 770. settled fact. Hibernation Victoria. Grape Phylloxera in Geelong, 1732. Aletia xylina in the, 2141. Silk-culture in the, 1668, 1945, 2070, Vilfa vaginæflora, 402, 1495, 1538. Vine area of France affected by Phylloxera. Half 2196. industry in the (Review), the, 2020.

Blue caterpillars of the, 1127, 1363.

2268.

Vine-growers. Thrips of, 53. infested with parasites. Hog caterpillar of the, 1247. Insects injurious to the, 2238. loving pomace fly, 2119. Vinegar for insects. Salt and, 937. Virgin tiger-moth. Chrysalis of the, 674. Virginia tiger-beetle, 1059. Vitis lituus, 1116. vinifera. Phylloxera destructive to, 1727. viticola, 1116. V-marked Cacœcia, 2363. Viviparity of a moth, 2153. Volume. Close of the first, 701. Voyle, J. Experiments on orange scale insects, 2164. Report on the effects of cold on the scale insects of the orange in Florida, Waguer, B. Observations on the new crop gallgnat, 2267. Walker, P. Serrell's automatic silk-reel, 2388. Silk-culture, 2418. Walking-stick, 144. Thick-thighed, 1701, 1721. Two-striped, 1298. sticks. Cicadas and, 920. or specters, 1395. Walnut case-bearer, 1311. caterpillars, 1229. Gregarious, 1045. Insect feeding on the sap of black, 1195. WALNUT. INSECTS AFFECTING. Cacœcia rileyana, 1059. Clisiocampa disstria, 363. sylvatica = C. disstria. Datana ministra, 1045, 1229, 2222, 2333. Notodontid, 757. Phycis juglandis, 1311. Psocus venosus, 1195. Walnut Tortrix, 1059. tree. Worm on bark of, 757. and willow. Enemies of the black, 2333. Walsh, B. D. Entomological collection of, 1107, 1203. In memoriam, 1098. Portrait of, 1128. Posthumous paper by, 1344. Successor to, 1105. Walshia amorphella, 1127, 2356. Wanted. Information, 1111. War on corn-worms, 1522. Warbles, 898. Washington Entomological Society. Address as president, 2355. Wasp. Ichneumon-fly mistaken for, 477. and parasites. Mud, 1827. spider-egg nest. Mud, 1847. Wasps, 1736. Digger, 309. and their habits, 375, 543, 615. Social, 616. Sucking organs of, 2182. Water bug, 797, 1191.

Gigantic, 534.

larva, 1205.

Water. Moths attracted by falling, 2103. weevil, 2119. of rice-plant, 1973. Wavy-striped flea-bectle, 348, 564, 2291. Wax-beans. Epilachna corrupta eating, 2135. insect. New, 2119. Unnatural secreticn of, 782. worm, 1059. Waxy secretion on stem of bitter-sweet. White, 1913. Ways of bag-worms, 1424. Wealth to the United States. A new source of, Web-worm. Fall, 1301, 1733, 2238, 2378, 2394. Garden, 2363. on hickory. Fall, 460. Juniper, 1721. Sorghum, 2119. Turf, 2418. of young trout. So-called, 1283. of young trout. Death, 1138, 1160, 1174. Webster, F. M. Insects affecting barley, 2394. buckwheat, 2394. fall-wheat, 2291, 2363, 2394. small grains and grasses, 2394. timothy, 2394. white clover, 2394. Record of experiments on the effect of punctures of hemiptera upon shrubs, fruits, aud grains, 2382. Report on Buffalo gnats, 2388. of experiments at Lafayette, Ind., 2344. upon the season's observations in Indiana, especially upon corn insects, 2418. Weeping lace-wing, 1127. Weevil, 1673. on apple-trees, New York, 1085. Clover, 1777. Honey-locust seed, 1026. Rice, 1683. plant water, 1973. Strawberry, 2363. Weevils. Aniseed vs. grain, 1742. Raspberries destroyed by, 1788. Well-known animals, Little known facts about, Wells. White worms in, 1015. Canker worms at the, 1539. West. Economic investigations in the South and, Genuine army-worm in the, 2009. Hop growing in the, 235, 279. Important observations on the grasshop-

per pest of the, 1571.

Winged pests of the, 2313.

encmies in, 2343.

Locusts in the, 2044.

Western cricket, 1959, 2267.

Indies. Report on the cotton crop and its

Western cricket, Bruner, L. Observations on the, 2267.

Histology of, 1959.

governors on the Rocky Mountain locust, 1557.

grasshopper, 1998.

Missouri. Ravages of young locusts in,

striped cut-worm, 1059.

Westward progress of the imported cabbageworm, 1694, 1721.

What are army-worms? 1400.

becomes of bumble bees? 1032.

Wheat. Beetles supposed to be feeding on, 1746. Bruchus. A new insect in, 19. Cause of smut in, 1461.

cut-worm, 1059.

Cut-worms destroying recently sown, 455. Damage to, 1870.

fly. Companion, 2394. head army-worm, 1570, 2418. Hessian fly in seed, 494.

insects, 1506.

WHEAT. INSECTS AFFECTING.

Aphis avenæ = Nectarophora granaria.

Asopia farinalis, 980.

Blissus leucopterus, 886, 888, 894, 895, 2363.

Brachytarsus variegatus, 1461, 1479.

Bruchus sp., 19.

Calandra remotepunctata, 15.

Calocoris rapidus, 2363.

Cecidomyia destructor, 494, 1506.

tritici = Diplosis tritici.

Cecidomyid, 18.

Chlorops sp., 2060.

Cicadula exitiosa, 1766.

Cut-worm, 1059.

Diedrocephala flaviceps, 1766, 1767, 2363.

Diplosis tritici, 109, 110, 142, 216, 280, 292, 372, 428, 711, 1512, 2238, 2267, 2363.

Dræocoris rapidus = Calocoris rapidus.

Euschistus fissilis, 2363.

Gortyna nitela, 1589, 1870.

Isosoma grande, 2288, 2291, 2363, 2394.

hordei, 2394.

nigrum = I. hordei.

tritici, 2060, 2063, 2123, 2363, 2394.

Jassus sexnotatus, 1766.

sp., 1766.

Leucania albilinea, 1507, 1570.

Longitarsus sp., 636.

Lygus pratensis, 2363.

Meromyza americana, 727, 1058, 1506, 1589, 1848, 1875, 2394.

Nectarophora granaria, 1806.

Noctuid, 455.

Oscinus sp., 2394.

Pyralis farinalis = Asopia farinalis.

Silvanus surinamensis, 1259.

Sitophilus remotepunctata = Calandra remotepunctata.

Tenebrio molitor, 980.

Wire-worm, 2418.

Wheat Isosoma, 2063, 2119.

just before it ripeus, Meromyza americana attacking, 1058.

Wheat. Leaf-hoppers injuring, 1766.

maggots, 727.

midge, 109, 110, 142, 216, 280, 292, 428, 711, 1512, 2238, 2291, 2363.

A new euemy to, 1507.

insect injurious to, 2288.

oats, and rye. Beetles working in, 1259.

plant. Klippart's (Review), 186.

rust and Hessian fly, 1605.

Smut in, 1479.

stalk-worm on Pacific coast, 2123.

stalks. New depredator infesting, 2060, 2063.

straw Isosoma, 2291, 2394.

Larger, 2291, 2394.

Webster, F. M. Insects affecting, 2394.

fall, 2291, 2363, 2394.

wire-worm, 2418.

Worm in joints of, 1848.

worms, 980.

injuring, 1875.

in reference to, 1589.

White ant. Inquilines in galleries of, 1729.

blast, 2119.

clover. Webster, F. M. Insects affecting, 2394.

grub, 68, 410, 1020, 1059, 2238, 2363, 2394.

fungus, 594, 1064, 1430, 1436, 1599, 1803,

1823.

again, 640.

information wanted, 1072.

parasite, 1363.

grubs, 541.

in strawberry-beds, 1236.

heart hickory Gelechia, 2363.

lined morning Sphinx, 630, 1198, 1301, 1643,

marked tussock-moth, 1059, 1363, 2238, 2378.

again, 535.

caterpillar, 1227.

eggs, 480, 600.

oak. Leaf-miner on, 1879.

Woolly gall on, 739. pine plaut-louse, 320.

scale, 97.

trees killed by borers, 1049.

weevil, 781, 2363.

worm, 465, 956.

Abbot's, 1570.

scale on acacias. Large, 1730.

Means against 2374.

waxy secretion on stems of bitter-sweet, 1913.

willow insects, 907.

worm, 917, 1211.

worms in wells, 1015.

Why noxious insects increase upon us, 766.

Wier, D. B. Native plums, 2388.

Wier's apple-worm trap. Thomas, 1312.

trap. The coddling-moth, 1334.

Wild cherry. Caterpillar's nest on, 242.

cherry. Cocoon on, 259.

grape-vine. Conical galls on leaves of, 1077. sage. Galls growing on, 1347.

Will unimpreguated eggs hatch? 1029.

Willow. Enemies of the black walnut and, 2333. gall. Pod-like, 1170.

galls, 46.

of Cecidomyidæ on, 46, 197.

Gracilaria. Purple, 2363.

WILLOW. INSECTS AFFECTING.

Cecidomyia salicis-siliquæ, 1170.

Cecidomyidæ, 46, 197.

Cimbex americana, 1380, 2291, 2333.

laportei = C. americana.

Clostera americana = Ichthyura inclusa.

Colcopteron. 197.

Dipteron, 197.

Gracilaria, 2363.

Hymenopteron, 197.

Ichthyura inclusa, 856.

Nematus ventralis, 907, 917, 1211.

Saperda, 2267.

Tenthredinidæ, 197.

Teras, 2363.

Vanessa antiopa, 907, 1234, 1609.

Willow insects. White, 907.

Teras, 2363.

worm. Large, 1380.

White, 917, 1121.

worms. Gregarious, 856.

Winged pests of the West, 2313.

Phylloxera in California, 1895.

Wire-worms, 224, 892, 932, 1030, 2238.

Means against, 103. in potatoes, 846.

Wits jump together. How great, 567.

W-marked cut-worm, 1059, 2291.

Wonder. Back-rolling, 1363.

Wood-borers. Food habits of, 1902.

Rearing, 1829.

gall on white-oak, 739.

lice on grape-vine roots, 1900.

nymph. Beautiful, 1127, 1363.

Pearl, 1127, 1301, 1363.

Woolly elm-tree lonse, 1059.

lice on the beach, 449.

slug-like worm on apple, 796.

Work in entomology, 202, 418.

Three worms and their, 1504.

Worker bees? Do toads eat, 544.

Workers among hymenoptera, 311.

Works on North American microlepidoptera, 1975.

World. Insect, 1466, 1467. Worm. Apple, 1666.

in apple, 177.

Army, 11, 17, 120, 670, 876, 906, 1127, 1442, 1482, 1551, 1570, 2119, 2239, 2267, 2269.

on bark of walnut tree, 757.

boring into cucumber, 808, 843.

peach, 1182.

in wheat-stalk, 1870.

Canker, 86, 172, 1021, 1066, 1127, 1363, 2012.

in corn, 181.

Cotton, 1127, 1363, 1649, 1702, 1719, 2077, 2119, 2130, 2295.

Currant, 877, 1204.

eating in green tomatoes, 734.

Hundred legged, 219.

infesting meal sacks, 1896.

Worm. In joint of wheat, 1848.

question, 909.

and remedy. Strawberry, 1586.

snake, 1662.

Tobacco, 885, 2238.

Tomato, 152, 1059, 1558, 1661.

An unknown, 986.

Worms. Bee bread devoured by, 1293.

in cabbage, 1915.

Clover, 83, 675, 948, 1132.

in eottonwood, 1459.

Currant, 364, 882, 1696.

on Dutchman's pipe, 1321.

feeding on hawthorn, 1051.

in flonr and rve, 72.

on horse-chestnut. Gregarious, 1192.

How to free wells of, 1015.

injuring wheat, 1875.

in joint wheat, 1848.

under mulch hay, 1161.

in osage orange seed, 597.

in potatoes. Wire, 846.

in wells, 1015.

Wire, 224, 892, 932, 1030, 2238.

and their work. Three, 1504.

Worthlessness of the sparrow as an insect killer, 2413.

Wyckoff's silk-industry in the United States (Re-

view), 2268. Wyoming, Montana, etc., in 1881. Bruner, L. The Rocky Mountain locust in, 2267.

Xanthoptera ridingsii n. sp., 1411 = Exyra ridingsii.

semicrocea, 1385, 1390.

Xiphidium, 118.

Xyleborus cælatus, 2267.

obesus, 1583.

Packard, A. S. Development of, 2267. pyri, 2, 149.

Xyleutes robiniæ = Cossus robiniæ.

Xylina cinerea n.sp., 1301 = Lithophane antennata.

Xylocopa carolina = X. virginica.

virginica, 372, 1111.

Xyloryctes satyrus, 40, 471.

Xylotrechus colonus, 2267.

Yama-mai silkworm, 1311.

Year. Destructive insects of the, 2322.

Entomological notes of the, 2235.

'Insects of the, 2289, 2331.

Is this a grasshopper, 1565.

Yeast ferment in insects. Experiments with, 1930. Yellow-bear. Common, 1301.

fever-fly, 1950.

headed cut-worm, 1059.

locust, 2363.

tailed Tachina-fly, 1127, 1625.

swallow-tail, 361.

Yellows in peach trees. Supposed cause of, 515.

Yersin. Function of the nervous system of articulates, 1959.

Young grasshoppers, 2377.

locusts. Destruction of, 1577.

Ditching for, 1488.

Habits of, 1578.

Young locusts in western Missouri. Ravages of, 1492.

pecan trees girdled, 489. tobacco-plants. Flea-beetle on, 1782. trout. Death-web of, 1138, 1160, 1174.

So-called web-worm of, 1283.

Yucca, 1602, 1603.

angustifoliå fertilized by Pronuba yuccasella, 1804.

borer, 1420, 1482, 1570.

Notes on the, 1465.

Fertilization of, 1329, 1336, 2171.

filamentosa. Prodoxus decipiens ovipositing in, 2049.

Further notes on the pollination of, 2000.

YUCCA. INSECTS AFFECTING.

Elaphidion tectum, 1602.

Megathymus yuccæ, 1420, 1465, 1482, 1570, 1602.

Prodoxus decipiens, 2000, 2049.

Pronuba yuccasella, 1329, 1336, 1337, 1354, 1363, 160%, 2000, 2171, 2391.

Yucca. Insects affecting—Continued. Sycophorus yuccæ, 1602.

Yucca moth, 1363.

Oviposition of the, 1354. Professor Riley and the, 2068.

True and bogus, 1804.
Pollination of, 1603, 1804, 2000, 2371.

Pronuba and its connection with the pollination of, 2391.

Zaræa inflata, 2360.

Zarhipis, 2397.

Further notes on Phengodes and, 2411. Phengodes and, 2408.

Zebra cabbage-worm, 2232.

caterpillar, 179, 1127.

Zeller. Death of Professor, 2179.

Zerene catenaria, 1140.

Zimmermann's flea-beetle, 2291.

Zoo-geographical map of North America. Packard, A, S., 2267.

Zygæna minos, 2224.

. ¢ \ ' • . .





